

## Erratum

# Diastolic function in patients with heart failure with preserved ejection fraction and atrial fibrillation: impact of diabetes: Am J Cardiovasc Dis. 2021; 11(5): 564-575

Ruxandra-Nicoleta Horodinschi<sup>1,2</sup>, Camelia Cristina Diaconu<sup>1,2</sup>

<sup>1</sup>Department of Internal Medicine, University of Medicine and Pharmacy "Carol Davila", Bucharest, Romania;


<sup>2</sup>Internal Medicine Clinic, Clinical Emergency Hospital of Bucharest, Bucharest, Romania

Received November 27, 2021; Accepted December 3, 2021; Epub December 15, 2021; Published December 30, 2021

In this article, there is a mistake in **Figure 1**, about the exact number of patients with HF and AF (253 patients with HF and AF) and the eligible patients with HFpEF and AF included in the statistical analysis after applying the inclusion and exclusion criteria (105 patients with HFpEF and AF). I attach below the **Figure 1** in its final form. We also found another mistake in the Materials and Methods section, the number of recruited patients should be "720", not "702". There are no other changes in the text.

**Address correspondence to:** Dr. Camelia Cristina Diaconu, Department of Internal Medicine, University of Medicine and Pharmacy "Carol Davila", 8 Calea Floreasca, Bucharest 014461, Romania. Tel: +40-726377300; E-mail: [drcameliadiaconu@gmail.com](mailto:drcameliadiaconu@gmail.com)

## Heart failure with preserved ejection fraction and atrial fibrillation



**Figure 1.** Selection of the patients included in the study. Legend: HF, heart failure; AF, atrial fibrillation; LVEF, left ventricular ejection fraction; LV, left ventricular; MI, myocardial infarction; UA, unstable angina; T2DM, type 2 diabetes mellitus; HFpEF, heart failure with preserved ejection fraction; OAD, oral antidiabetic drugs; IR, insulin requiring.