

## Original Article

# IFN- $\gamma$ , IL-17, IL-22<sup>+</sup> CD4<sup>+</sup> subset in patients with hepatitis C virus and correlation with clinical factor

Soolmaz Khansalar<sup>1</sup>, Zahra Faghih<sup>2</sup>, Shaghik Barani<sup>1</sup>, Mehdi Kalani<sup>3</sup>, Mohammad Reza Ataollahi<sup>4</sup>, Zeinab Mohammadi<sup>5</sup>, Sepideh Namdari<sup>1</sup>, Kurosh Kalantar<sup>1,6,7</sup>

<sup>1</sup>Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran; <sup>2</sup>Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran; <sup>3</sup>Department of Immunology, Professor Alborzi Clinical Microbiology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran; <sup>4</sup>Department of Immunology, School of Medicine, Fasa University of Medical Sciences, Fasa, Iran; <sup>5</sup>Department of Radiation Oncology, Memorial Sloan-Kettering Cancer Center, New York, NY, USA; <sup>6</sup>Autoimmune Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran; <sup>7</sup>Department of Bacteriology and Immunology and The Translational Immunology Research Program (TRIMM), The University of Helsinki and HUSLAB, Helsinki University Hospital, Helsinki, Finland

Received December 27, 2023; Accepted February 19, 2024; Epub February 25, 2024; Published February 28, 2024

**Abstract:** Background: CD4<sup>+</sup> T cell responses in HCV infection have a crucial role in the immunopathology of hepatitis C virus (HCV) infection. Our aim was to investigate the frequency of Th1, Th17, and Th22 cells in HCV-infected patients and elucidate their role in the progression of the disease. Methods: Twenty-six HCV-infected patients and 26 healthy individuals were recruited. Peripheral blood mononuclear cells (PBMCs) were stained to separate CD4, IFN- $\gamma$ , IL-17, and IL-22 producing cells using flow cytometry. Results: Results showed that the mean expression of IL-22 in CD4<sup>+</sup> T cells was significantly lower in HCV-infected patients compared to healthy controls. About correlation with clinical factor and T subsets, a negative correlation between the frequency of CD4<sup>+</sup> IFN- $\gamma$ <sup>+</sup> cells and Thyroxine level (T4) was observed in the patients. The data showed a positive link between thyroid-stimulating hormone (TSH), cholesterol levels, and the frequency of Th17 cells. In addition, a positive correlation was seen between serum creatinine level with both Th1 and Th17. Ultimately, it was found that there was a positive link between viral burden and IL-17<sup>+</sup> IL-22<sup>+</sup> cells and a negative correlation between viral load and pure Th22. Conclusions: Our findings indicate that Th22 cells may play a part in the immunopathology of HCV and show the associations between Thelper subsets and the clinical signs of the disease.

**Keywords:** Hepatitis C, Th1, Th17, Th22, inflammation

## Introduction

Hepatitis C is a liver infection that arises from the hepatitis C virus (HCV) categorized as a hepatotropic and non-cytopathic virus that is transmitted through contaminated blood products, sexual activity, and intravenous drug use [1, 2]. HCV infection remains a major global health problem, with around 58 million people of the world population being infected with it, while there is no vaccine for clinical use yet [3]. HCV infection is the prime cause of acute hepatitis, consequently resulting in chronic liver disease, which is characterized by fibrosis and cirrhosis and may ultimately lead to hepatocellular carcinoma (HCC) [4, 5]. In the chronic course

of HCV infection, persistent virus replication in the hepatocytes may promote the stimulation of inflammatory responses leading to progressive fibrosis and damage to the liver [6].

It has been established that during HCV infection, a wide variety of cytokines contribute to viral clearance but also tissue injury [7]. The pro-inflammatory cytokines are fundamental prerequisites for initiating the inflammatory cascades and sustaining the chronic infection, which eventually results in hepatocellular liver injury [8]. Besides, chronic and inflammatory courses of HCV infection might cause an imbalance between the expression of pro-inflammatory and anti-inflammatory cytokines and che-

## T helper subset in hepatitis C virus patients

mokines [9]. Furthermore, a large and growing body of research has emphasized the importance of CD8<sup>+</sup> cytotoxic T lymphocytes (CTLs) and CD4<sup>+</sup> T helper (Th) cells that have influential roles in cell-mediated immunity against HCV infection [10, 11]. Naïve CD4<sup>+</sup> T cells can differentiate into various distinct subtypes, including Th1, Th2, Th17, Th22, and regulatory T cells with definite immunoregulatory functions in both steady and inflammatory states [12]. Among these, Th1 cells are characterized by IFN- $\gamma$  production and robust protection against intracellular pathogens [13]. IFN- $\gamma$  is a pleiotropic molecule related to mediate. Immune responses including pro-apoptotic antiproliferative and antitumor mechanisms which promote the antiviral activity and facilitate macrophage activation [14-16]. Th17 cells produce pro-inflammatory cytokines such as IL-17 (A, F) and IL-22, which promote the recruitment of other immune cells to the inflamed and injured sites [13, 17]. IL-17, also known as IL-17A, activates the neutrophil mobilization and activation. It can mediate protective innate immunity against pathogens and has been shown to be a key cytokine in protecting the host from mucosal infections [18, 19]. Th22 cells, as another helper subset, are specifically involved in producing IL-22 and can modulate the tissue responses to inflammation. IL-22 is recognized as a unique cytokine that contributes critically to the crosstalk between the immune cells and epithelial cells [20].

In this study, we aimed to examine the frequency of peripheral Th1, Th17, and Th22 subsets and the level of their corresponding cytokines, including IFN- $\gamma$ , IL-17, and IL-22, in patients with chronic HCV infection. Furthermore, we investigated whether the alterations in the frequency of Th subsets were associated with the clinical features of the patients, including certain hematological and biochemical findings.

### Materials and methods

#### *Study population*

This study was carried out on 26 patients (24 males and 2 females) who were infected with hepatitis C infection. Our study was conducted before admitting any antiviral or immunomodulatory treatments. The subjects (42.26 $\pm$ 9.8 years old) were recruited from Motahari Clinic affiliated with Shiraz University of Medical Sciences over a one-year period. Twenty-six

age- and gender-matched healthy individuals (42.20 $\pm$ 10 years old) were also included in the study as the control group. All enrolled subjects provided written informed consent according to the Declaration of Helsinki and the research was reviewed and confirmed by the Medical Ethics Committee of Shiraz University of Medical Sciences (IR.SUMS.REC. 1396.S731).


#### *Peripheral blood mononuclear cell (PBMC) isolation*

Five milliliters (ml) of heparinized whole peripheral blood were collected from all healthy controls and patients before receiving any treatment. PBMCs were isolated by Ficoll density gradient centrifugation. The viability of the cells was determined using the trypan blue exclusion test (Sigma-Aldrich, Germany). PBMCs were counted and suspended in 2 ml of complete culture medium (RPMI 1640) supplemented with 100 U/ml penicillin, 100  $\mu$ g/ml streptomycin, 2 mM glutamine, and 10% heat-inactivated fetal calf serum, at a density of 2 $\times$ 10<sup>6</sup> cells/ml. All were purchased from Gibco, USA.

#### *Cell activating and staining*

The PBMCs were stimulated with 50 ng/mL phorbol myristate acetate (PMA) plus ionomycin (1  $\mu$ g/ml) (both from Sigma-Aldrich) in the presence of 0.7  $\mu$ l Brefeldin A, 0.7  $\mu$ l Monensin, and 2  $\mu$ l DNase (BD Biosciences, USA) for 5 hours in a CO<sub>2</sub> incubator [21, 22]. After the incubation period, the cells were harvested and centrifuged at 350 g and 4°C for 10 min. The supernatant was removed, and the pellet was washed with 2 ml of 1 $\times$  phosphate-buffered saline (PBS); then, the cells were fixed with 300  $\mu$ l 1% paraformaldehyde (Sigma-Aldrich) at 4°C for 15 minutes. Following washing with 1 $\times$  PBS, the pellet was diluted in Perm/Wash buffer (Biolegend, USA) and incubated for 15 min in the dark at room temperature for permeabilization. Then, the tube was centrifuged for 10 min at 500 g, and the supernatant was discarded. The cells were transferred into flow cytometry tubes, stained with PerCP/Cyanine5.5 CD4 (RPA-T4) FITC anti-human IFN- $\gamma$  (B27), APC anti-human IL-17A (BL168), and PE anti-human IL-22 antibodies (2G12A41) (all from Biolegend); subsequently, they were incubated for 30 min at 4°C in the dark. After we washed the tubes to remove the extra antibodies, the supernatant was dis-

## T helper subset in hepatitis C virus patients



**Figure 1.** Flow cytometry analysis of different CD4<sup>+</sup> T helper cell subsets. A. Lymphocytes were first gated based on their relative forward and side scatters. B. Cells with high CD4 expression were then gated among the lymphocytes. C-E. The frequency of IFN- $\gamma$ , IL-17, and IL-22-producing CD4<sup>+</sup> T cells was determined in CD4<sup>+</sup> lymphocytes. F-H. The up arrows under the figures show the pure Th1, Th17, and Th22 populations.

carded and 300  $\mu$ l 1% paraformaldehyde was added to the tubes for 15 min at 4°C. Subsequently, the cells were washed and resuspended in 300  $\mu$ l 1 $\times$  PBS. To control the background staining, we used appropriate matched isotype control antibodies, including FITC mouse IgG1, PE mouse IgG2a, PerCP/Cy5.5 mouse IgG1, and APC mouse IgG2b (all from Biolegend). Flow cytometric analysis was carried out using a FACS-Calibur flow cytometer (BD Bioscience), and the obtained data were

further analyzed using FlowJo software 7.6.1 (Tree Star Inc., USA).

### *Gating strategy for evaluating the frequency of Th1, Th17, and Th22 cells*

PBMCs were first divided into two distinct populations of lymphocytes and monocytes based on their relative forward and side scatter (**Figure 1A**). Next, the lymphocytes with a high expression of CD4 were gated (**Figure 1B**), and

## T helper subset in hepatitis C virus patients

**Table 1.** Frequencies of various CD4<sup>+</sup> T helper subsets in untreated HCV-infected patients and controls

Lymphocyte subset	Markers	Untreated HCV-infected patients (Mean ± SD)	Controls (Mean ± SD)	P value
CD4 <sup>+</sup>	CD4 <sup>+</sup>	31.53±14.99	39.61±8.74	0.058
IFN-γ-producing	CD4 <sup>+</sup> IFN-γ <sup>+</sup>	17.43±10.64	16.45±5.96	0.952
IL-17-producing	CD4 <sup>+</sup> IL-17 <sup>+</sup>	1.67±0.85	1.65±0.65	0.680
IL-22-producing	CD4 <sup>+</sup> IL-22 <sup>+</sup>	0.96±0.65	0.83±0.37	0.656
Pure CD4 <sup>+</sup> T helper subsets				
Pure Th1	CD4 <sup>+</sup> IFN-γ <sup>+</sup> IL-17 <sup>-</sup> IL-22 <sup>-</sup>	16.14±10.31	16.66±6.20	0.477
Pure Th17	CD4 <sup>+</sup> IL-17 <sup>+</sup> IFN-γ <sup>-</sup> IL-22 <sup>-</sup>	1.26±0.63	1.22±0.49	0.805
Pure Th22	CD4 <sup>+</sup> IL-22 <sup>+</sup> IFN-γ <sup>-</sup> IL-17 <sup>-</sup>	0.36±0.26	0.31±0.17	0.878

the frequency of different Th subsets was determined based on the expression of IFN-γ (Figure 1C), IL-17 (Figure 1D), and IL-22 (Figure 1E). The IFN-γ, IL-17, and IL-22 single-producing T cells (negative for the two other cytokines) were defined as pure Th1, Th17, and Th22, respectively (Figure 1F-H). The percentages of double-positive IL-17/IFN-γ, IL-22/IFN-γ, and IL17/IL-22 cells were also determined. The frequency of the defined populations was calculated as a percentage of CD4<sup>+</sup> lymphocytes. The geometric mean fluorescent intensity (MFI) of each cytokine was normalized by dividing the MFI of positive cells by the MFI of negative cells and considered as a criterion for its expression level per target cell.

### Statistical analysis

All data were analyzed using SPSS 16.0 software (SPSS Inc., Chicago, IL, USA). The distribution of the data was checked using Kolmogorov-Smirnov test and accordingly non-parametric tests were used to analyze the data. Mann-Whitney U test was carried out to compare the difference between the two groups, along with the Kruskal-Wallis test for multiple comparisons. The Spearman correlation test was run to detect the potential correlations between the variables. The quantitative data were presented as the mean ± standard deviation (SD), and *p*-values less than 0.05 were regarded statistically significant.

### Results

Distribution of CD4<sup>+</sup> T helper subsets expressed IFN-γ, IL-17, and IL-22 in HCV-infected patients and controls. Flow cytometry data indicated that the average frequency of various CD4<sup>+</sup> Th

subsets in HCV-infected patients compared to healthy controls was not significantly different. However, analyzing the frequency of total CD4<sup>+</sup> lymphocytes indicated a trend towards a lower frequency of CD4<sup>+</sup> T cells in HCV-infected patients compared to the healthy controls (*P* = 0.058).

We adopted an approach to identify the pure cells. Pure Th1 (CD4<sup>+</sup> IFN-γ<sup>+</sup> IL-17<sup>-</sup> IL-22<sup>-</sup>), pure Th17 (CD4<sup>+</sup> IL-17<sup>+</sup> IFN-γ<sup>-</sup> IL-22<sup>-</sup>), and pure Th22 (CD4<sup>+</sup> IL-22<sup>+</sup> IFN-γ<sup>-</sup> IL-17<sup>-</sup>) cells in patient and control groups, exclusively expressed only one cytokine by gating cells. They were identified as pure cells. However, they were not significant between the patient and control groups (Table 1). We also examined the cells that expressed two cytokines simultaneously; there were no significant differences in the frequency of double-positive Th cells, including IFN-γ<sup>+</sup> IL-17<sup>+</sup>, IFN-γ<sup>+</sup> IL-22<sup>+</sup>, and IL-17<sup>+</sup> IL-22<sup>+</sup> cells between patients and healthy controls. This data is not shown in the table.

### Mean expression of IFN-γ, IL-17, and IL-22 cytokines in T helper subsets


MFI is the average amount of fluorescence emitted by specific cells in a cell population, defined as a measure of the number of antigens on the surface and within cells in mean or median fluorescence units [23]. We utilized MFI for IFN-γ, IL-17, and IL-22 expression levels in the corresponding T cell subsets (i.e., CD4<sup>+</sup> IFN-γ<sup>+</sup> cells, CD4<sup>+</sup> IL-17<sup>+</sup> cells, and CD4<sup>+</sup> IL-22<sup>+</sup> cells) as well as pure Th subsets (i.e., pure Th1, Th17, and Th22 cells). The comparison of HCV-infected patients and healthy controls demonstrated a lower level of IL-22 in CD4<sup>+</sup> IL-22<sup>+</sup> T cells of HCV-infected patients compared to the

## T helper subset in hepatitis C virus patients

**Table 2.** MFI of IFN- $\gamma$ , IL-17, and IL-22 in Th1, Th17, and Th22 cells of patients and controls

Cytokines	Corresponding lymphocyte subsets	HCV-infected patients (Mean $\pm$ SD)	Controls (Mean $\pm$ SD)	P value
IFN- $\gamma$	CD4 <sup>+</sup> IFN- $\gamma$ <sup>+</sup>	107.76 $\pm$ 111.92	99.71 $\pm$ 46.34	0.286
IL-17	CD4 <sup>+</sup> IL-17 <sup>+</sup>	215.07 $\pm$ 175.58	290.68 $\pm$ 172.94	0.060
IL-22	CD4 <sup>+</sup> IL-22 <sup>+</sup>	6.21 $\pm$ 1.70	7.50 $\pm$ 1.70	0.0008*
MFI of pure CD4 <sup>+</sup> T helper subsets				
Pure Th1	CD4 <sup>+</sup> IFN- $\gamma$ <sup>+</sup> IL-17 <sup>-</sup> IL-22 <sup>-</sup>	86.74 $\pm$ 48.99	90.44 $\pm$ 28.66	0.459
Pure Th17	CD4 <sup>+</sup> IL-17 <sup>+</sup> IFN- $\gamma$ <sup>-</sup> IL-22 <sup>-</sup>	208.95 $\pm$ 167.34	276.90 $\pm$ 182.16	0.105
Pure Th22	CD4 <sup>+</sup> IL-22 <sup>+</sup> IFN- $\gamma$ <sup>-</sup> IL-17 <sup>-</sup>	16.47 $\pm$ 7.08	19.26 $\pm$ 9.57	0.370

\*P<0.05: statistically significant; based on Mann-Whitney U test. MFI: Mean Fluorescence Intensity.



**Figure 2.** Correlations between different T helper subsets and various biochemical characteristics and viral load in untreated HCV-infected patients.

controls ( $P = 0.0008$ ). We also found a trend towards a lower expression level of IL-17 in CD4<sup>+</sup> IL-17<sup>+</sup> lymphocytes of the patients compared to the controls ( $P = 0.060$ ). The expres-

sion level of each cytokine was also analyzed in pure Th subsets, which revealed no significant differences between the patients and controls (**Table 2**).

*Correlation of CD4<sup>+</sup> Th subsets with the hematological and biochemical parameters of the patients with hepatitis C*

Examining the potential correlations between different Th subsets and the clinical parameters of the patients, which were collected from the history of the patients, revealed a negative correlation between the frequency of CD4<sup>+</sup> IFN- $\gamma$ <sup>+</sup> cells and T4 levels ( $R = -0.481$ ,  $P = 0.027$ ) (**Figure 2A**); however, there was a positive correlation between CD4<sup>+</sup> IL-17<sup>+</sup> Th cells and serum TSH levels ( $R = 0.551$ ,  $P = 0.012$ ) (**Figure 2B**). Regarding the serum creatinine level (**Figure 2C** and **2D**), we noticed a positive correlation with the frequency of both pure Th1 cells ( $R = 0.24$ ,  $P = 0.049$ ) and pure Th17 cells ( $R = 0.439$ ,  $P = 0.047$ ). Moreover, the frequency of pure Th1 cells was positively correlated with the platelet count of the patients ( $R = 0.493$ ,  $P = 0.014$ ) (**Figure 2E**). Furthermore, a higher number of Th17 cells were found in patients with higher cholesterol levels ( $R = 0.483$ ,  $P = 0.024$ ) (**Figure**

**2F).** We also observed a negative correlation between the viral load and the frequency of pure Th22 ( $R = -0.512$ ,  $P = 0.036$ ) (**Figure 2G**) and a positive correlation between the viral load and IL-17<sup>+</sup> IL-22<sup>+</sup> cells ( $R = 0.527$ ,  $P = 0.030$ ) (**Figure 2H**).

### Discussion

Hepatitis C is one of the most significant public health challenges and concerns. The therapeutic effectiveness of various anti-viral regimens for HCV eradication depends on the maintenance of a fine balance among various pro- and anti-inflammatory cells and their mediators [8]. The CD4<sup>+</sup> T cell response in HCV infection has a crucial role in viral control, and liver damage can induce HCV infections [7]. To investigate the variations in the frequency of peripheral blood T cell subsets, we examined the frequency of peripheral Th1, Th17, and Th22 cells in untreated HCV-infected patients and healthy controls. In addition, any possible relationships between the Th subsets and clinical laboratory findings of the patients with hepatitis C were evaluated. We aim to elucidate potential associations that may contribute to a better understanding of the complex interplay between the host immune response and the clinical manifestations of HCV infection and also recognition of damage to different body organs caused by viral infection.

Th1 cells are essential for virus clearance in HCV infection and play a crucial role in the immune response to HCV. However, during the clearance of this virus, the Th1 cytokines contribute to necrosis of the hepatocytes. We found no significant differences in the frequency of IFN- $\gamma$ <sup>+</sup> CD4<sup>+</sup> lymphocytes as well as the mean expression of IFN- $\gamma$  in these cells between HCV-infected patients and healthy controls. This observation aligns with the findings of a study that reported no significant alteration in serum IFN- $\gamma$  levels among individuals with HCV infection compared to the healthy control group [8], but contrary to another findings which was reported a higher level of IFN- $\gamma$  in chronic HCV infection compared with healthy controls [24]. A higher expression of IFN- $\gamma$  assessed by enzyme-linked immunosorbent spot assay was also observed in the chronic HCV cases [25].

Regarding IL-17-producing lymphocytes, no significant changes were found in HCV-infected

patients compared with the controls, while previous studies reported an increase in both the frequency of Th17 and the serum level of IL-17 in these patients [26, 27]. Many studies have revealed the detrimental and beneficial impacts of IL-17 on virus-induced pathology. IL-17 function is crucial in fostering viral infections by enhancing virus replication and suppressing antiviral Th1 or CTL immunity to maintain the persistence of HCV in the chronic cases. Moreover, it can inhibit the development of regulatory cells and promote Th2 immune responses, which induce hepatic fibrosis [26, 28].

Th22 cells and IL-22 have been observed to have various functions in the immune system. Several studies have reported an association between increased levels of IL-22 and the severity of hepatic diseases, which can be attributed to the proliferative and anti-apoptotic effects of IL-22 [29, 30]. Detrimental pro-inflammatory effects have been illustrated to deteriorate certain conditions, such as hepatitis B and *Helicobacter pylori* infection [31]; also, previous research has highlighted that IL-22 and IL-22BP have a role in liver diseases with a particular focus on the leading causes of advanced liver failure, namely liver infections, liver damage, and liver malignancies [32].

However, some related studies reported the upregulation of IL-22-producing CD4<sup>+</sup> T cells in patients with acute hepatitis B compared to the controls [31, 33]. Also, examining the serum level of patients with mild/moderate COVID-19 and healthy controls, researcher observed the upregulation of IL-22 cytokines levels in COVID-19 patients [34].

IL-22 also has beneficial anti-inflammatory features that help impede the advancement of infections such as influenza and HIV. The protective role of IL-22 against liver fibrosis and its contribution to hepatic tissue regeneration through prompting the expression of metalloproteinase and inducing the antimicrobial activity have been demonstrated [31, 35]. We found a lower MFI of IL-22 in IL-22<sup>+</sup> CD4<sup>+</sup> T cells in patients with hepatitis C compared to the healthy controls. Our results are in the same line with another study [36]. Given the decreased MFI of IL-22 in the patients with chronic hepatitis C compared with the controls, it can be implied that persistent viral infection leads to the immune exhaustion of certain subsets of the helper cells, Th22 cells in par-

## T helper subset in hepatitis C virus patients

ticular, and reduces the IL-22 production [37]. Another underlying mechanism could be explained by the ability of regulatory T cells to suppress Th22 and Th17 cells as the main sources of IL-22 production, which is mediated by the altered expression of inhibitory receptors including PD-1, CTLA-4, and TIM-3 and regulatory cytokines such as IL-10 [38].

T4 is a hormone that is mainly synthesized in the thyroid follicles and plays an important role in regulating the metabolism. TSH is produced in the brain to stimulate the thyroid gland in order to produce and release more thyroid hormone [39]. Viral diseases like Chronic HBV (Hepatitis B viruses) infections and Covid viruses can be associated with changes in the level of these hormones [40, 41].

With respect to the clinical aspects, we observed a negative correlation between T4 level and the frequency of CD4<sup>+</sup> IFN- $\gamma$ <sup>+</sup> T cells and a positive correlation between TSH level and the percentage of CD4<sup>+</sup> IL-17<sup>+</sup> T cells. The production of IFN- $\gamma$ , as a necro-inflammatory cytokine, by CD4<sup>+</sup> lymphocytes may result in liver injury which consequently reduces the T4 level through the secretion of thyroid hormone-binding globulin (T4-TBG) in the liver [42, 43]. There have been several reports of elevated levels of TSH in patients with chronic hepatitis C, notably those who had not received any treatment; it can be attributed to hypothyroidism following persistent HCV infection [44, 45]. Furthermore, CD4<sup>+</sup> IL-17<sup>+</sup> T cells have a key role in stimulating the production of thyroid autoantibodies and certain Th17-related chemokines involved in the pathogenesis of thyroid diseases [46, 47]. Creatinine is a waste product created in the body from the breakdown of creatine, a chemical that supplies the muscles with energy. Depending on the muscle mass, it is released from the body at a constant rate. Creatinine is primarily filtered from the blood by the kidneys and excreted in the urine [48, 49]. Another important finding of this study was a positive correlation between the creatinine level and the frequency of pure Th1 and Th17 cells in untreated HCV patients. This is in the same line with the results of some studies carried out on renal insufficiency and higher creatinine levels in HCV-infected cases, showing that there is very little scientific understanding of the role of HCV in kidney failure [50]. A study has shown Th1 and Th17 cells are involved in

the pathogenesis of HCV in patients with glomerulopathy [51]. As to our finding about platelets, we observed a positive correlation between PLT and pure Th1, similar to a study in this examination, they found that the pathogenesis of enterovirus 71 (EV71)-caused hand, foot, and mouth disease (HFMD) had a positive correlation between the platelets with the frequency of Th1 cells. They concluded that PLT were responsible for the regulation of pathogenic Th1 cells differentiation and function in the pathogenesis of the disease [52].

Cholesterol homeostasis is dependent on the activation of LXR nuclear receptors. This receptor can have a significant impact on Th17 cell differentiation. HCV is able to affect these nuclear receptors and cholesterol metabolism through direct and indirect mechanisms [53]. Our result from the present study was a direct correlation between the cholesterol level and frequency of pure Th17 cells, which is consistent with previous studies indicating altered lipid metabolism in HCV infection [54].

Interestingly, there was a negative correlation between the viral load and frequency of pure Th22. Several roles have been considered for this cytokine such as anti-oxidation, anti-bacterial anti-viruses, anti-fibrosis, anti-inflammatory and regenerative stimulation in protecting against liver and gut damage [23, 31]. This cytokine has a protective role in the liver and gut epithelium, when its levels are decreased. We can suggest that by decreasing IL-22 the viral load is increased and probably the liver and gut epithelial have been damaged after infections; as a result, the viral load has increased, eventually causing T cell exhaustion by transferring the virus to the blood [55, 56].

Since Th17 cells are mainly the producers of IL-17 and IL-22; also, the cytokines are involved in inflammation and maintenance of gut epithelial barrier function. In addition, a positive correlation with IL-17<sup>+</sup> IL-22<sup>+</sup> T cells and viral load was observed. Thus, with the increase in the viral load, Th17 cells also increased, and it can be mentioned that this subset of cells has been trying to improve the mucosal damage [57, 58].

Taken together, the observed inconsistencies between our results and those of others might be attributed to variations in the sample size and methodology, along with dissimilarities in

## T helper subset in hepatitis C virus patients

the viral load, genotypes, the extent of liver fibrosis and damage, and duration of inflammatory responses. Notwithstanding these limitations, our results provide new insights into the role of different helper subsets and their cytokine profiles in the pathology of HCV infection, which would be a fruitful area for further research.

### Acknowledgements

We would like to express our sincere gratitude in our article to Lida Ahmadi and Leila Kazemi Miraki for their dedicated service at Motahari Clinic. We thank Professor Zahra Amirghofran for her invaluable counseling services. Additionally, we extend my thanks to the patients diagnosed with hepatitis C individuals and normal group the who participated in this project and appreciate the efforts of the Clinical Research Development Center at Namazi Hospital for their valuable contributions to editing the article.

### Disclosure of conflict of interest

None.

**Address correspondence to:** Kurosh Kalantar, Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz 71348-45794, Iran. Tel: +98-7112351575; Fax: +98-7112351575; E-mail: Kalantark@sums.ac.ir

### References

- [1] Schillie S, Wester C, Osborne M, Wesolowski L and Ryerson AB. CDC recommendations for hepatitis C screening among adults - United States, 2020. *MMWR Recomm Rep* 2020; 69: 1-17.
- [2] Michalak TI. HCV lymphotropism and its pathogenic significance. *Hepatitis C-From Infection to Cure* 2018; 45-65.
- [3] Taha G, Ezra L and Abu-Freha N. Hepatitis C elimination: opportunities and challenges in 2023. *Viruses* 2023; 15: 1413.
- [4] Gülceciği DE, Goeser T and Kasper P. Prognostic assessment of liver cirrhosis and its complications: current concepts and future perspectives. *Front Med (Lausanne)* 2023; 10: 1268102.
- [5] Khatun M and Ray RB. Mechanisms underlying hepatitis C virus-associated hepatic fibrosis. *Cells* 2019; 8: 1249.
- [6] Mondelli MU, Ottolini S, Oliviero B, Mantovani S, Cerino A, Mele D and Varchetta S. Hepatitis C virus and the host: a mutual endurance leaving indelible scars in the host's immunity. *Int J Mol Sci* 2023; 25: 268.
- [7] Abouelasrar Salama S, Lavie M, De Buck M, Van Damme J and Struyf S. Cytokines and serum amyloid A in the pathogenesis of hepatitis C virus infection. *Cytokine Growth Factor Rev* 2019; 50: 29-42.
- [8] Neuman MG and Cohen LB. Inflammation and liver cell death in patients with hepatitis C viral infection. *Curr Issues Mol Biol* 2021; 43: 2022-2035.
- [9] Pocino K, Stefanile A, Basile V, Napodano C, D'Ambrosio F, Di Santo R, Callà CAM, Gulli F, Saporito R, Ciasca G, Equitani F, Basile U and Marino M. Cytokines and hepatocellular carcinoma: biomarkers of a deadly embrace. *J Pers Med* 2022; 13: 5.
- [10] Ryscavage P, Hussien S, Seung H and Hynicka L. CD4+ T-cell recovery in HIV/hepatitis C co-infected patients following successful hepatitis C treatment. *HIV Med* 2024; [Epub ahead of print].
- [11] Borhani K, Bamdad T, Hashempour A, Salek Farrokhi A and Moayed J. Comparison of the inhibitory and stimulatory effects of Core and NS3 candidate HCV vaccines on the cellular immune response. *Am J Clin Exp Immunol* 2023; 12: 153-163.
- [12] Kunkl M, Frasca S, Amormino C, Volpe E and Tuosto L. T helper cells: the modulators of inflammation in multiple sclerosis. *Cells* 2020; 9: 482.
- [13] Rožman P and Švajger U. The tolerogenic role of IFN- $\gamma$ . *Cytokine Growth Factor Rev* 2018; 41: 40-53.
- [14] Castro F, Cardoso AP, Gonçalves RM, Serre K and Oliveira MJ. Interferon-gamma at the crossroads of tumor immune surveillance or evasion. *Front Immunol* 2018; 9: 847.
- [15] Ng CT, Fong LY and Abdullah MNH. Interferon-gamma (IFN- $\gamma$ ): Reviewing its mechanisms and signaling pathways on the regulation of endothelial barrier function. *Cytokine* 2023; 166: 156208.
- [16] Pratim Das P and Medhi S. Role of inflammasomes and cytokines in immune dysfunction of liver cirrhosis. *Cytokine* 2023; 170: 156347.
- [17] Cachem FCOF, Dias AS, Monteiro C, Fernandes G, Delphim L, Tavares F, Maciel AMA, Amendola-Pires MM, Brandão-Mello CE, Andrade RM and Bento CAM. Different core-specific T cell subsets are expanded in chronic hepatitis C with advanced liver disease. *Cytokine* 2019; 124: 154456.
- [18] Matsuzaki G and Umemura M. Interleukin-17 family cytokines in protective immunity against infections: role of hematopoietic cell-derived and non-hematopoietic cell-derived interleukin-17s. *Microbiol Immunol* 2018; 62: 1-13.



## T helper subset in hepatitis C virus patients

- [19] Mills KHG. IL-17 and IL-17-producing cells in protection versus pathology. *Nat Rev Immunol* 2023; 23: 38-54.
- [20] Seth P and Dubey S. IL-22 as a target for therapeutic intervention: current knowledge on its role in various diseases. *Cytokine* 2023; 169: 156293.
- [21] Faghieh Z, Erfani N, Haghshenas MR, Safaei A, Talei AR and Ghaderi A. Immune profiles of CD4+ lymphocyte subsets in breast cancer tumor draining lymph nodes. *Immunol Lett* 2014; 158: 57-65.
- [22] Haghshenas MR, Khademi B, Faghieh Z, Ghaderi A and Erfani N. Immune regulatory cells and IL17-producing lymphocytes in patients with benign and malignant salivary gland tumors. *Immunol Lett* 2015; 164: 109-116.
- [23] Tynngård N, Alshamari A, Månsson F and Ramström S. Variation in activation marker expression within the platelet population - a new parameter for evaluation of platelet flow cytometry data. *Platelets* 2022; 33: 1113-1118.
- [24] de Souza-Cruz S, Victória MB, Tarragô AM, da Costa AG, Pimentel JP, Pires EF, Araújo Lde P, Coelho-dos-Reis JG, Gomes Mde S, Amaral LR, Teixeira-Carvalho A, Martins-Filho OA, Victória Fda S and Malheiro A. Liver and blood cytokine microenvironment in HCV patients is associated to liver fibrosis score: a proinflammatory cytokine ensemble orchestrated by TNF and tuned by IL-10. *BMC Microbiol* 2016; 16: 3.
- [25] Dai Z, Zeng W, Li G and Shu X. A conserved domain of HCV E2 glycoprotein interacts with human CD81 and induces interferon-gamma secretion from peripheral blood mononuclear cells. *Front Biosci (Landmark Ed)* 2023; 28: 239.
- [26] Cabral MS, Santos TPS, Santos PL, Schinoni MI, Oliveira IS, Pereira AB, Atta AM and Sousa-Atta MLB. Immune response of Th17-associated cytokines by peripheral blood mononuclear cells from patients with chronic hepatitis C virus infection. *Cytokine* 2018; 102: 200-205.
- [27] Gomaa AF, Wahba MO, Hafez RAEL, Eldaly OM and Badran SG. Assessment of the role of interleukin 17A and interleukin 17F in chronic hepatitis C virus infection in Egyptian patients. *Egypt J Intern Med* 2019; 31: 199-202.
- [28] Ma WT, Yao XT, Peng Q and Chen DK. The protective and pathogenic roles of IL-17 in viral infections: friend or foe? *Open Biol* 2019; 9: 190109.
- [29] Giannou AD, Lücke J, Kleinschmidt D, Shiri AM, Steglich B, Nawrocki M, Zhang T, Zazara DE, Kempinski J, Zhao L, Giannou O, Agalioti T, Brockmann L, Bertram F, Sabihi M, Böttcher M, Ewald F, Schulze K, von Felden J, Machicote A, Maroulis IC, Arck PC, Graß JK, Mercanoglu B, Reeh M, Wolter S, Tachezy M, Seese H, Theodorakopoulou M, Lykoudis PM, Heumann A, Uzunoglu FG, Ghadban T, Mann O, Izbicki JR, Li J, Duprée A, Melling N, Gagliani N and Huber S. A critical role of the IL-22-IL-22 binding protein axis in hepatocellular carcinoma. *Cancers (Basel)* 2022; 14: 6019.
- [30] Wu Y, Min J, Ge C, Shu J, Tian D, Yuan Y and Zhou D. Interleukin 22 in liver injury, inflammation and cancer. *Int J Biol Sci* 2020; 16: 2405-2413.
- [31] Zhang K, Chen L, Zhu C, Zhang M and Liang C. Current knowledge of Th22 cell and IL-22 functions in infectious diseases. *Pathogens* 2023; 12: 176.
- [32] Lücke J, Sabihi M, Zhang T, Bauditz LF, Shiri AM, Giannou AD and Huber S. The good and the bad about separation anxiety: roles of IL-22 and IL-22BP in liver pathologies. *Semin Immunopathol* 2021; 43: 591-607.
- [33] Wang LY, Yang XY, Wu YP and Fan YC. IL-22-producing CD3 + CD8- T cells increase in immune clearance stage of chronic HBV infection and correlate with the response of Peg-interferon treatment. *Clin Immunol* 2023; 250: 109320.
- [34] Majeed AY, Zulkafli NES and Ad'hiah AH. Interleukin-22 and interleukin-33 show up-regulated levels in the serum of patients with mild/moderate Coronavirus disease 2019. *Beni Suef Univ J Basic Appl Sci* 2023; 12: 24.
- [35] Hammerich L and Tacke F. Hepatic inflammatory responses in liver fibrosis. *Nat Rev Gastroenterol Hepatol* 2023; 20: 633-646.
- [36] Baskic D, Vukovic VR, Popovic S, Djurdjevic P, Zaric M, Nikolic I, Zelen I, Mitrovic M, Avramovic D and Mijailovic Z. Cytokine profile in chronic hepatitis C: an observation. *Cytokine* 2017; 96: 185-188.
- [37] Kahan SM, Wherry EJ and Zajac AJ. T cell exhaustion during persistent viral infections. *Virology* 2015; 479-480: 180-193.
- [38] Liang M, Liwen Z, Yun Z, Yanbo D and Jianping C. The imbalance between Foxp3<sup>+</sup>Tregs and Th1/Th17/Th22 cells in patients with newly diagnosed autoimmune hepatitis. *J Immunol Res* 2018; 2018: 3753081.
- [39] Pirahanchi Y, Toro F and Jialal I. Physiology, thyroid stimulating hormone. In: *StatPearls. Treasure Island (FL) ineligible companies. Disclosure: Fadi Toro declares no relevant financial relationships with ineligible companies. Disclosure: Ishwarlal Jialal declares no relevant financial relationships with ineligible companies: StatPearls Publishing Copyright © 2023, StatPearls Publishing LLC.; 2023.*
- [40] Razu MH, Hossain MI, Ahmed ZB, Bhowmik M, Hasan MKE, Kibria MK, Moni DA and Khan M. Study of thyroid function among COVID-19-affected and non-affected people during pre and post-vaccination. *BMC Endocr Disord* 2022; 22: 309.

## T helper subset in hepatitis C virus patients

- [41] Chen JF, Weng WZ, Huang M, Peng XH, Zhang J, Xiong J, He JR, Zhang SQ, Cao HJ, Gao B, Lin DN, Gao J, Gao ZL and Lin BL. The impact of serum thyroid-stimulation hormone levels on the outcome of hepatitis B virus related acute-on-chronic liver failure: an observational study. *BMC Gastroenterol* 2022; 22: 330.
- [42] El-Feki MAF, Abdalla NH, Atta MI and Ibrahim AA. Serum level of thyroid hormones in patients with chronic hepatitis C virus infection. *Open J Endocr Metab Dis* 2016; 6: 126-134.
- [43] Chakravarthy V and Ejaz S. Thyroxine-binding globulin deficiency. In: *StatPearls*. StatPearls Publishing; 2019.
- [44] Batool N, Elahi S, Saleem N and Ashraf A. Thyroid dysfunction in non-interferon treated hepatitis C patients residing in hepatitis endemic area. *Biomed Res Int* 2017; 2017: 2390812.
- [45] Toma L, Zgura A, Isac T, Simu R, Mercan-Stanciu A, Dodot M and Iliescu EL. COVID-19 and the thyroid function in patients with HCV-associated hepatocellular carcinoma. *Acta Endocrinol (Bucharest)* 2022; 18: 392-396.
- [46] Huang Y, Fang S, Li D, Zhou H, Li B and Fan X. The involvement of T cell pathogenesis in thyroid-associated ophthalmopathy. *Eye (Lond)* 2019; 33: 176-182.
- [47] Mele C, Caputo M, Bisceglia A, Samà MT, Zavattaro M, Aimaretti G, Pagano L, Prodham F and Marzullo P. Immunomodulatory effects of vitamin D in thyroid diseases. *Nutrients* 2020; 12: 1444.
- [48] Aujla RS and Patel R. Creatine phosphokinase. In: *StatPearls*. Treasure Island (FL) ineligible companies. Disclosure: Roshan Patel declares no relevant financial relationships with ineligible companies: StatPearls Publishing Copyright © 2024, StatPearls Publishing LLC.; 2024.
- [49] Shahbaz H and Gupta M. Creatinine clearance. In: *StatPearls*. Treasure Island (FL) ineligible companies. Disclosure: Mohit Gupta declares no relevant financial relationships with ineligible companies: StatPearls Publishing Copyright © 2024, StatPearls Publishing LLC.; 2024.
- [50] Barsoum RS, William EA and Khalil SS. Hepatitis C and kidney disease: a narrative review. *J Adv Res* 2017; 8: 113-130.
- [51] Ozkok A and Yildiz A. Hepatitis C virus associated glomerulopathies. *World J Gastroenterol* 2014; 20: 7544-54.
- [52] Li Q, Wang Y, Xue W, Bian Z, Gao Y, Zeng Y, Tang L, Tang T, Tian Y and Guo W. Immunomodulatory effects of platelets on the severity of hand, foot, and mouth disease infected with enterovirus 71. *Pediatr Res* 2021; 89: 814-822.
- [53] Maggio R, Viscomi C, Andreozzi P, D'Ettore G, Viscogliosi G, Barbaro B, Gori M, Vullo V and Balsano C. Normocaloric low cholesterol diet modulates Th17/Treg balance in patients with chronic hepatitis C virus infection. *PLoS One* 2014; 9: e112346.
- [54] Alterations in lipid and lipoprotein levels: helpful or harmful to the heart? *Clin Infect Dis* 2017; 65: 566-567.
- [55] Radaeva S, Sun R, Pan Hn, Hong F and Gao B. Interleukin 22 (IL-22) plays a protective role in T cell-mediated murine hepatitis: IL-22 is a survival factor for hepatocytes via STAT3 activation. *Hepatology* 2004; 39: 1332-1342.
- [56] Mboumba Bouassa RS, Comeau E, Alexandrova Y, Pagliuzza A, Yero A, Samarani S, Needham J, Singer J, Lee T, Bobeuf F, Vertzagias C, Sebastiani G, Margolese S, Mandarino E, Klein MB, Lebouché B, Routy JP, Chomont N, Costiniuk CT and Jenabian MA. Effects of oral cannabinoids on systemic inflammation and viral reservoir markers in people with HIV on antiretroviral therapy: results of the CTN PT028 pilot clinical trial. *Cells* 2023; 12: 1811.
- [57] Hasan MZ, Islam S, Matsumoto K and Kawai T. SARS-CoV-2 infection initiates interleukin-17-enriched transcriptional response in different cells from multiple organs. *Sci Rep* 2021; 11: 16814.
- [58] Walker EM, Slisarenko N, Gerrets GL, Grasperge BF, Mattison JA, Kissinger PJ, Welsh DA, Veazey RS, Jazwinski SM and Rout N. Dysregulation of IL-17/IL-22 effector functions in blood and gut mucosal gamma delta T cells correlates with increase in circulating leaky gut and inflammatory markers during cART-treated chronic SIV infection in macaques. *Front Immunol* 2021; 12: 647398.