

Epigenetic silencing of klotho in gastric cancer

- hepatocellular carcinoma and other digestive tumors. *Hepatology* 2008; 48: 508-518.
- [27] Yu J, Cheng YY, Tao Q, Cheung KF, Lam CN, Geng H, Tian LW, Wong YP, Tong JH, Ying JM, Jin H, To KF, Chan FK and Sung JJ. Methylation of protocadherin 10, a novel tumor suppressor, is associated with poor prognosis in patients with gastric cancer. *Gastroenterology* 2009; 136: 640-651 e641.
- [28] Sanchez-Carbayo M, Socci ND, Lozano J, Saint F and Cordon-Cardo C. Defining molecular profiles of poor outcome in patients with invasive bladder cancer using oligonucleotide microarrays. *J Clin Oncol* 2006; 24: 778-789.
- [29] Radvanyi L, Singh-Sandhu D, Gallichan S, Lovitt C, Pedyczak A, Mallo G, Gish K, Kwok K, Hanna W, Zubovits J, Armes J, Venter D, Hakimi J, Shortreed J, Donovan M, Parrington M, Dunn P, Oomen R, Tartaglia J and Berinstein NL. The gene associated with trichorhinophalangeal syndrome in humans is overexpressed in breast cancer. *Proc Natl Acad Sci U S A* 2005; 102: 11005-11010.
- [30] Karnoub AE, Dash AB, Vo AP, Sullivan A, Brooks MW, Bell GW, Richardson AL, Polyak K, Tubo R and Weinberg RA. Mesenchymal stem cells within tumour stroma promote breast cancer metastasis. *Nature* 2007; 449: 557-563.
- [31] Su LJ, Chang CW, Wu YC, Chen KC, Lin CJ, Liang SC, Lin CH, Whang-Peng J, Hsu SL, Chen CH and Huang CY. Selection of DDX5 as a novel internal control for Q-RT-PCR from microarray data using a block bootstrap resampling scheme. *BMC Genomics* 2007; 8: 140.
- [32] Gumz ML, Zou H, Kreinest PA, Childs C, Belmonte LS, LeGrand SN, Wu KJ, Linton BA, Saha M, Parker AS, Sun LZ, Ahlquist J, Wood JD and Copland JA. Selected methylated protein 1 loss contributes to tumor phenotype of clear cell renal cell carcinoma. *Clin Cancer Res* 2007; 13: 4740-4749.
- [33] Talantov D, Mazumder A, Yu JX, Briggs T, Jiang Y, Backus J, Atkins D and Wang Y. Novel genes associated with malignant melanoma but not benign melanocytic lesions. *Clin Cancer Res* 2005; 11: 7234-7242.
- [34] Gordon GJ, Rockwell GN, Jensen RV, Rheinwald JG, Glickman JN, Aronson JP, Pottorf BJ, Nitz MD, Richards WG, Sugarbaker DJ and Bueno R. Identification of novel candidate oncogenes and tumor suppressors in malignant pleural mesothelioma using large-scale transcriptional profiling. *Am J Pathol* 2005; 166: 1827-1840.
- [35] Pawitan Y, Bjohle J, Amler L, Borg AL, Egyhazi S, Hall P, Han X, Holmberg L, Huang F, Klaar S, Liu ET, Miller L, Nordgren H, Ploner A, Sandelin K, Shaw PM, Smeds J, Skoog L, Wedren S and Bergh J. Gene expression profiling spares early breast cancer patients from adjuvant therapy: derived and validated in two population-based cohorts. *Breast Cancer Res* 2005; 7: R953-964.
- [36] Nutt CL, Mani DR, Betensky RA, Tamayo P, Cairncross JG, Ladd C, Pohl U, Hartmann C, McLaughlin ME, Batchelor TT, Black PM, von Deimling A, Pomeroy SL, Golub TR and Louis DN. Gene expression-based classification of malignant gliomas correlates better with survival than histological classification. *Cancer Res* 2003; 63: 1602-1607.
- [37] Fridman AL, Rosati R, Li Q and Tainsky MA. Epigenetic and functional analysis of IGFBP3 and IGFBPPrP1 in cellular immortalization. *Biochem Biophys Res Commun* 2007; 357: 785-791.
- [38] Ibanez de Sacaca I, Cortes-Sempere M, Moratilla A, Machado-Pinilla R, Rodriguez-Ranjo V, Manguan-Garcia C, Cejas P, Lopez-Rios P, Paz-Ares L, de CastroCarpeno J, Nistal M, Beldarrainista C and Perona R. IGFBP-3 hypomethylation-derived deficiency mediates cisplatin resistance in non-small-cell lung cancer. *Oncogene* 2010; 29: 1681-1690.
- [39] Smith P, Nicholson LJ, Syed N, Payne A, Hiller L, Garrone O, Occelli M, Gasco M and Crook T. Epigenetic inactivation implies independent functions for insulin-like growth factor binding protein (IGFBP)-related protein 1 and the related IGFBPL1 in inhibiting breast cancer phenotypes. *Clin Cancer Res* 2007; 13: 4061-4068.
- [40] Wiley A, Katsaros D, Fracchioli S and Yu H. Methylation of the insulin-like growth factor binding protein-3 gene and prognosis of epithelial ovarian cancer. *Int J Gynecol Cancer* 2006; 16: 210-218.
- [41] Chang YS, Wang L, Liu D, Mao L, Hong WK, Khuri FR and Lee HY. Correlation between insulin-like growth factor-binding protein-3 promoter methylation and prognosis of patients with stage I non-small cell lung cancer. *Clin Cancer Res* 2002; 8: 3669-3675.