

Erratum

Hispidulin prevents hypoxia-induced epithelial-mesenchymal transition in human colon carcinoma cells: Am J Cancer Res. 2015; 5(3): 1047-1061

Jing Xie^{1*}, Hui Gao^{1*}, Jianjun Peng^{2*}, Yantao Han^{1*}, Xuehong Chen¹, Qixiao Jiang¹, Chunbo Wang¹

¹Medical College, Qingdao University, Qingdao 266071, Shandong, China; ²College of Life Sciences, Chongqing Normal University, Chongqing 401331, China. *Equal contributors.

Received February 27, 2020; Accepted March 2, 2020; Epub April 1, 2020; Published April 15, 2020

Recently, we rechecked all our papers and realized that some data in **Figures 2** and **3** were wrongly presented. The data presented came from the wrong folder. Therefore, we would like to make corrections. Hope you can help me with this process. The revised **Figures 2** and **3** are presented as following:

Address correspondence to: Dr. Jianjun Peng, College of Life Sciences, Chongqing Normal University, Chongqing 401331, China. E-mail: jianjunpeng@126.com; Dr. Yantao Han, Medical College, Qingdao University, Qingdao 266071, Shandong, China. E-mail: hanyt19@126.com

Hispidulin prevents hypoxia-induced EMT in cancer cells




Figure 2. Hispidulin suppresses hypoxia-induced migration (A) and invasion (B) in HT-29 cells. HT-29 cells growing in hypoxia were treated with hispidulin (25 μ M) for 24 hours. Cell migration and invasion were assessed by wound scratch and Transwell assays, respectively. The results represent mean \pm SD from three independent experiments. *P < 0.05 vs. control, ^P < 0.05 vs. hypoxia.

Hispidulin prevents hypoxia-induced EMT in cancer cells




Figure 3. Hispidulin prevents hypoxia-induced EMT. HT-29 cells growing in hypoxia were treated with hispidulin (25 μ M) for 24 hours. A. Effect of hispidulin on cell morphology. The images represent three independent experiments. B. The effect of hispidulin on the expression of E-cadherin examined by fluorescence staining, western blotting and qRT-PCR. The images and blot are representative of three independent experiments. C. The effect of hispidulin on the expressions of N-cadherin, vimentin, fibronectin and α -SMA determined by western blotting and qRT-PCR. The immunoblot represents three independent experiments. Data are presented as mean \pm SD. *P < 0.05 vs. control, ^P < 0.05 vs. hypoxia.