

Erratum

Heat shock protein 70-2 (HSP70-2) a novel cancer testis antigen that promotes growth of ovarian cancer: Am J Cancer Res. 2017; 7(6): 1252-1269

Namita Gupta¹, Nirmala Jagadish¹, Avadhesh Suroli², Anil Suri¹

¹Cancer Microarray, Genes and Proteins Laboratory, National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi 110067, India; ²Molecular Biophysics Unit, Indian Institute of Science, Bangalore 560012, India

Received April 29, 2020; Accepted May 4, 2020; Epub June 1, 2020; Published June 15, 2020

In the paper of AJCR0041068R1, a mistake has been found that in the published version of **Figure 5, Figure 5B** panel at 0 hour of HSP70-2 shRNA3 treatment was inadvertently presented with same image for 0 hour of HSP70-2 shRNA4 treatment during figure assembly during manuscript submission. We have accordingly corrected and replaced the image in **Figure 5B** panel at 0 hour for HSP70-2 shRNA4 in corrected **Figure 5**. Change in representative image does not affect the interpretation in **Figure 5**. The corrected **Figure 5** is enclosed. The figure legend is correct as published and is

also shown for reference below. The error has no bearings on the interpretation of the results, nor do they influence the conclusions of the work.

Address correspondence to: Anil Suri, Convener, Cancer Research Program, Cancer Microarray, Genes and Proteins Laboratory, National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi 110067, India. Tel: 91-11-26703-700; Fax: 91-11-26-1621-25, 91-11-26742125; E-mail: anil@nii.res.in

HSP70-2 promotes ovarian cancer cellular growth and invasion




Figure 5. HSP70-2 ablation inhibits cellular motility. A. Phase contrast images and histogram show the significant difference in migration or invasion abilities through insert membrane and matrigel in shRNA3 and shRNA4 transfected cells respectively as compared to NC shRNA transfected cells. B. Representative images show delayed wound healing in shRNA3 and shRNA4 transfected cells as compared to NC shRNA transfected cells from 12-48 hours. Histogram depicts relative wound healing index at different time points. C. Western blot analysis shows the upregulation of epithelial markers, while downregulation of mesenchymal markers. Original magnification $\times 100$, objective $\times 10$. The experiments were performed three independent times in triplicates. Data are represented as mean \pm standard error of the mean. * $P < 0.05$, ** $P < 0.001$, *** $P < 0.0001$.