

Erratum

Muscle weakness caused by cancer and chemotherapy is associated with loss of motor unit connectivity: Am J Cancer Res. 2021; 11(6): 2990-3001

Joshua R Huot^{1,2}, Fabrizio Pin², Andrea Bonetto^{1,2,3,4,5}

¹Department of Surgery, ²Department of Anatomy, Cell Biology & Physiology, ³Department of Otolaryngology-Head & Neck Surgery, ⁴Indiana Center for Musculoskeletal Health, ⁵Simon Comprehensive Cancer Center, Indiana University School of Medicine, Indianapolis, IN, USA

Received February 11, 2022; Accepted February 17, 2022; Epub March 15, 2022; Published March 30, 2022

In this article, a mistake has been found in Materials and Methods, Animals section, the correct dosing for the folfiri *in vivo* administration is: 50 mg/kg 5-fluorouracil, 90 mg/kg leucovorin, 24 mg/kg CPT-11; 2×/week.

of Medicine, 980 W Walnut Street, R3-C522, Indianapolis, IN 46202, USA. Tel: 317-278-0302; Fax: 317-274-8046; E-mail: jrhuot@iu.edu; fpin@iu.edu; abonetto@iu.edu

Address correspondence to: Dr. Andrea Bonetto, Department of Surgery, Indiana University School