

## Original Article

# Effect of sunitinib derivatives on glioblastoma single-cell migration and 3D cell cultures

Girstaute Dabkeviciute<sup>1,3</sup>, Elias Maccioni<sup>2</sup>, Vilma Petrikaite<sup>1,3</sup>

<sup>1</sup>*Institute of Biotechnology, Life Sciences Center, Vilnius University, Sauletekio al. 7, LT-10257, Vilnius, Lithuania;*

<sup>2</sup>*Department of Life and Environmental Sciences, University of Cagliari, Via Ospedale 72, 09124 Cagliari, Italy;*

<sup>3</sup>*Laboratory of Drug Targets Histopathology, Institute of Cardiology, Lithuanian University of Health Sciences, Sukileliu pr. 13, LT-50162, Kaunas, Lithuania*

Received January 30, 2023; Accepted March 14, 2023; Epub April 15, 2023; Published April 30, 2023

**Abstract:** This study aimed to evaluate the anticancer activity of 16 new sunitinib derivatives in brain cancer cells (2D model) and spheroids (3D model). The effect on cell viability was determined by the MTT assay. Single-cell migration assay was performed to examine the effect of selected compounds on individual cell migration. The activity of compounds in 3D cell cultures was examined by measuring the size change of spheroids formed using the Hanging drop method. The viability of brain cancer (U-87MG and A-172) cells was most reduced by compound EMAC4001. EMAC4001 showed the strongest effect on U-87MG cell migration, and EMAC4007 was the most active in the A-172 cell line. Only sunitinib had a statistically significant impact on spheroid growth at 100 nM and 500 nM concentrations in the U87-MG cell line and EMAC4007 had a statistically significant impact on A-172 spheroid growth at 100 nM and 500 nM concentrations, similarly to sunitinib.

**Keywords:** Sunitinib analogs, glioblastoma, tyrosine kinase inhibitors, spheroids, cell migration

## Introduction

Glioblastoma (GBM), also called grade IV astrocytoma, is the most common and most malignant cancer of the central nervous system in adults [1]. The GBM and low-grade gliomas mainly differ in growth rate, angiogenesis, and necrosis [2]. The GBM prognosis is associated with poor quality of life: necrosis-induced changes in the nervous tissue cause behavioral, emotional, and cognitive disorders, which lead to patients' mental disability and lack of independence [3]. The median survival time for GBM patients is about 14 to 16 months, and there are still extremely limited opportunities for conventional therapy [4]. Therefore, to this day, GBM is considered an incurable type of cancer.

Astrocytoma develops from irregular, star-shaped glial cells called astrocytes, characterized by the ability to form pseudopodia, which allow GBM cells to spread deeper into healthy brain tissues while complicating the determina-

tion of the exact border between normal and malignant brain sites, making complete tumor resection impossible and GBM recurrence inevitable [5, 6]. There are various commercially available glioma cell lines. However, more elongated cells like U-87MG and A-172 are characterized by a higher invasiveness level [7]. Infiltrative brain tumors are associated with the disintegration of the normal astrocytic-endothelial links in the brain-blood barrier, but increased permeability of such barrier causes an increase of expression of various membrane transporters and pumps that limit drugs entry to tumor parenchyma making GBM hardly reachable for therapeutics. For instance, temozolomide (TMZ), the first-line agent for GBM treatment, has shown limited efficacy in GBM patients due to developed resistance related to overexpression of ATP-binding cassette drug efflux transporters and it was found that maximum TMZ concentration in the brain reaches only up to 20% of serum concentration, suggesting that research for new methods capable

## Sunitinib derivatives affect cell migration and spheroid growth

of improving GBM therapy and prognosis is crucial [8-10].

A relatively new approach to treat GBM is targeted therapy that involves small molecular drugs targeting cell surface receptors called receptor tyrosine kinases (RTKs), which play a significant role in most cellular processes by initiating intracellular signal transduction through two major pathways Ras/MAPK/ERK and Ras/PI3K/AKT [11]. Studies have shown that invasive and angiogenic tumor behavior is related to the activation of RTKs and growth factors, which are overexpressed in GBM as a result of various genetic alterations [12]. Small molecular inhibitors have been developed to block the active site of the kinase and thereby prevent phosphorylation resulting in inhibition of migration and apoptosis of cancer cells [13]. In GBM increased kinase activity is commonly detected in EGFR, PDGFR and VEGFR families based on tumor subtype and region. EGFR is known to be more often present in the classical subtype of GBM and is usually associated with highly proliferative properties [14, 15]. A few inhibitors targeting EGFR, such as erlotinib and lapatinib, were tested in clinical trials, unfortunately, compounds did not have significant effect for GBM patients suggesting that single-target or dual agents might not be sufficient in blocking tumor intracellular pathways and overcoming drug resistance [16, 17]. Other important targets for kinase inhibitors are PDGFR and VEGFR, as they promote invasive growth and angiogenesis of GBM [12, 18]. It was found that PDGFRs and their ligands are co-expressed in high-grade gliomas proposing an autocrine loop, not common for normal brain tissue, thereby making specific targeting possible [19]. VEGF is considered to be a critical mediator of new blood vessels formation in GBM and is known to be released as a response to hypoxia [20]. Additionally, upregulation of its receptors is common in mesenchymal subtype of GBM, which is associated with the lowest survival rates of all four subtypes [14]. Studies have demonstrated VEGFR and its ligands involvement not only in endothelial cells but also in GBM cells chemotaxis by activation of ERK1/2 pathway [21]. Moreover, sunitinib, a multi-targeted kinase inhibitor, reduced U87 and M059K glioma cells migration by inhibiting VEGFR-integrin $\beta$ (3) complex formation *in vitro* [22]. After encouraging results of *in vitro* and *in vivo* research, sunitinib has been tested as a

potent antiangiogenic agent for GBM patients in clinical trial, however, regardless the high activity in renal cell carcinoma and gastrointestinal stromal tumors, compound did not have sufficient effect in GBM treatment [23-25].

As GBM still remains incurable, it is necessary to search for new compounds that could overcome the present challenges of the treatment and could be used individually or in combination with conventional therapy. Because of high GBM vascularization level, most of targeted therapies that are being tested for GBM patients in clinical trials include antiangiogenic agents inhibiting VEGFR and PDGFR as main targets [26-29]. One of them is sunitinib, an indolin-2-one derivative inhibiting more than eight kinases with highest activity against VEGFR-1,-2,-3, PDGFR- $\alpha$  and PDGFR- $\beta$  [30]. Since it was determined that indolin-2-one is essential for antiangiogenic properties, new compounds possessing this fragment have been developed to enhance anticancer activity. The chemical structure of sunitinib consists of 5-fluoroindolin-2-one ring, pyrrole group and amino side chain (**Figure 1**). Substitutions in indolin-2-one core and modifications of pyrrole have shown to have a significant effect on anticancer activity, VEGFR-2 inhibition and toxicity [31, 32]. Therefore, isatin/dihydropyrazole hybrids (EMAC) have been synthesized, which are suggested as more active sunitinib analogs [33].


EMAC derivatives can be categorized into two major groups: compounds possessing 2-naphthyl or 2-thiophenyl residues at C-3 position of dihydropyrazole. Additionally, new analogs possess indolin-2-one, a fragment of sunitinib, also referred as isatin (indoline-2,3-dione), which has different substitutes at C-5 and C-7 positions. We hypothesize that new sunitinib derivatives might possess stronger tyrosine kinase inhibitory properties than sunitinib and be beneficial in GBM treatment. Hence, the aim our study is to evaluate the effect of new analogs on human brain cancer cell viability, migration, and spheroid growth.

### Materials and methods

#### Materials

EMAC derivatives were synthesized by Maccioni E. group (University of Cagliari, Italy) and pub-

## Sunitinib derivatives affect cell migration and spheroid growth



**Figure 1.** Tyrosine kinase inhibitor sunitinib and EMAC derivatives.

lished earlier [33]. Sunitinib malate was purchased from Sigma Aldrich (St. Louis, MO, USA).

### Cell culturing

The human brain cancer U-87MG and A-172 cell lines were obtained from the American Type Culture Collection (ATCC, Manassas, USA). Glioma cells were cultured in DMEM Glutamax medium (Gibco, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (Gibco) and 10,000 U/mL penicillin, and 10 mg/mL streptomycin (Gibco) at 37°C with 5% CO<sub>2</sub> in a humidified atmosphere. Cell cultures were grown to 70% confluency, trypsinized with 0.125% TrypLE Express solution (Gibco) before passage and used until passage 20.

### Cell viability assay


Cell viability was tested using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT, Sigma-Aldrich Co., ≥ 97%) reduction assay. U-87MG and A-172 were seeded into 96-well plates (Corning) in a volume of 100 μL (5 × 10<sup>3</sup> U-87MG cells/well or 4 × 10<sup>3</sup> A-172 cells/well). After 24 hours incubation, cells were treated with 100 μL of different concentrations of sunitinib analogs and sunitinib itself. Medium without cells was used as a positive control and medium with 0.5% DMSO

(Sigma-Aldrich Co., ≥ 99%) was considered as a negative control. After 72 hours, cells were incubated with 0, 5 mg/mL of MTT solution for 3 hours. The formed formazan crystals were dissolved in DMSO. The absorbance was measured spectrophotometrically at wavelengths of 570 and 630 nm. Half maximal effective concentration (EC<sub>50</sub>) values were calculated using the Hill equation.

### Cell migration assay

Cell migration was tested using the single-cell tracking method. U-87MG and A-172 glioma cells were seeded into collagen (Gibco) coated imaging dishes (IBIDI, Martinsried, Germany) with numbered grids in a volume of 2 mL (5 × 10<sup>3</sup> cells/mL). After 24 hours incubation, cells were treated with sunitinib analogs and sunitinib at the concentrations equal to EC<sub>50</sub> values, determined as described previously. After treatment with tested compounds, photos of gridded area were taken every hour by inverted microscope (Olympus), then processed using *ImageJ* software (National Institutes of Health, USA). Coordinates of cells were estimated in the X and Y axes, then distance travelled per hour was calculated using the Pythagoras theorem equation and presented as average cell velocity.

## Sunitinib derivatives affect cell migration and spheroid growth



**Figure 2.** The effect of sunitinib analogs on human brain cancer cell viability. U-87MG and A-172 cells were treated with each compound at 1  $\mu$ M concentration for 72 h and cell viability was determined by MTT assay. The asterisks (\*) indicate  $P < 0.05$ , compared to sunitinib,  $n = 3$ . Abbreviations: SNT, sunitinib.

### Spheroid growth

Spheroids were formed from U-87MG and A-172 glioma cells by the Hanging drop method. The cell suspension ( $5 \times 10^5$  cells/mL) was diluted with a medium containing 0.2% methylcellulose (average  $M_w \sim 90,000$ , Sigma-Aldrich Co.) and drops of 20  $\mu$ L ( $1 \times 10^3$  U-87MG cells/drop or  $2 \times 10^3$  A-172 cells/drop) were placed on the lid of the Petri dish (Corning). The number of cells was selected based on preliminary experiments performed at the beginning of the study. After 24 hours incubation, formed spheroids were placed into 96-well plate (Corning) coated with 2% agarose (Sigma-Aldrich Co.) and treated with 100 nM and 500 nM solutions of tested compounds. After 4 days, the medium containing solutions of tested compounds was changed by the fresh one. Images of spheroids were taken by inverted microscope (Olympus) every 24 hours for 9 days and processed with the *ImageJ* software (National Institutes of Health).

### Statistical analysis

Experiments were repeated three times, calculating the mean and standard deviation. The normality of the distribution was checked using the Shapiro-Wilk test. To determine the significant differences between means of normally distributed data, the Student's t-test was performed. The analysis of abnormally distributed data was conducted using Kruskal-Wallis test. The data of migration and spheroid growth analysis is displayed in box plots in which the vertical line indicates the median and the cross

indicates the mean. The level of statistical significance was set at  $P < 0.05$ . The data was processed using *Microsoft Office Excel 2020 (Version 2211)* and *IBM SPSS Statistics (Version 20)*.

## Results

### *Sunitinib analogs effect on brain cancer cells viability*

7 out of 16 new sunitinib analogs (compounds EMAC4001, 4002, 4005, 4015, 4018, 4019) reduced U87-MG cell viability stronger than A172 cell line. Out of all 16 sunitinib analogs, EMAC4001, 4006 and 4007 were the most active ones in both tested glioblastoma cell lines and reduced cell viability up to 44% (**Figure 2**). Based on their potent cytotoxic effect against brain cancer cells, EMAC4001, 4006 and 4007 were selected for further research.


After screening for potential novel sunitinib analogs,  $EC_{50}$  values of EMAC4001, EMAC4006 and EMAC4007 were calculated. Selected compounds showed from 18 to 69 times stronger effect on U87-MG cells viability and from 25 to 95 times stronger cytotoxic effect in A172 cell line, compared to sunitinib. The most active compound in both cell lines after 72 hours was EMAC4001, containing 5-chloroisatin fragment ( $EC_{50}$  values in U87-MG and A172 cell lines were  $81.7 \pm 6.0$  nM and  $72.3 \pm 6.0$  nM, respectively) (**Figure 3**).

### *Effect of sunitinib analogs on cell migration*


For single-cell migration experiment, the compounds were tested at the concentrations equal to their estimated  $EC_{50}$  values. To evaluate biological response of glioma cells to new compounds, the average cells velocity was determined and compared among control and experimental groups.

EMAC4001 and EMAC4006 statistically significantly inhibited the migration of single U-87MG cells. The most active compound was EMAC4001, which reduced cell velocity 2 times, compared to the control, and was 28%

## Sunitinib derivatives affect cell migration and spheroid growth



**Figure 3.** A. EC<sub>50</sub> values of sunitinib analogs. Medium without cells was used as a positive control and cells with 0.5% DMSO served as a negative control. The asterisks (\*) indicate P < 0.05, compared to sunitinib. B. Sunitinib and EMAC4001 effect on U-87MG cell viability after 72 h of incubation by MTT assay. Data points are experimental values (averages of three repeats) while the line is a fit of the standard inhibition model with the Hill coefficient of 6.5 (EMAC4001) and 1.5 (sunitinib). Abbreviations: SNT, sunitinib.



**Figure 4.** Effect of sunitinib analogs on (A) U-87MG cell and (B) A-172 cell migration. Cells were treated with compounds at their EC<sub>50</sub> values for 8 hours. The asterisks (\*) indicate P < 0.05 and the double asterisk (\*\*) indicates P < 0.01, compared to the control. Abbreviations: SNT, sunitinib.

more active than sunitinib. Regarding the migration of A172 cells, it was found that all tested compounds showed statistically significant inhibition of cell movement from 2 to 2.5 times, compared to the control. The most active compound in A172 cell line was compound EMAC4007 (Figure 4). Over 6 hours, the average U-87MG cell velocity increased steadily in the control group (Figure 5A) and in sunitinib group cell movement started decreasing after 3 hours of treatment (Figure 5B). After treatment with EMAC4001 and EMAC4006, U-87MG cell movement maintained reduced for the 6 hours (Figure 5C and 5D). No movement trends were observed in cell migration after treatment of EMAC4007 (Figure 5E). A-172 cells had simi-


lar migration routine over six hours (Data not shown).

### Effect of sunitinib analogs on spheroid growth

In 3 days both cell lines were able to form compact spheroids with a diameter ranging from 260 µm to 290 µm. U-87MG spheroids were more irregularly shaped and slightly larger in the start of the experiment (day 0), however, after the treatment A172 spheroids entered the log phase and surpassed U-87MG spheroids in size in 3 days (Figures 6 and 7).

After treatment with 100 nM solutions of tested compounds, U-87MG spheroids continued to grow after day 5 of the experiment, with the

## Sunitinib derivatives affect cell migration and spheroid growth



**Figure 5.** U-87MG cell migration at different time points in control group (A) and after treatment of sunitinib (B), EMAC4001 (C), EMAC4006 (D) and EMAC4007 (E). The asterisks (\*) indicate  $P < 0.05$ . Abbreviations: SNT, sunitinib.

sunitinib group as an exception, as it showed steady decrease in spheroid size after day 2. A lower concentration was sufficient to reduce the size of the U-87MG spheroids from 9.6 to 34.2%, however only in sunitinib group the size change was statistically significant compared to the control. Contrary to the results in 2D models, all compounds except EMAC4001 had significant effect on U-87MG spheroid growth at higher concentration (Figure 6). As in migration studies, compound EMAC4007 has shown a selective effect on the A172 cell line. It was the only compound that had statistically significant impact on glioma spheroid growth at lower concentration, similarly to sunitinib (Figure 6).


### Discussion

Our results showed that sunitinib analogs EMAC4001, EMAC4006, EMAC4007 possess stronger anticancer effect on tested GBM cell lines (U-87MG and A-172) in monolayer cells, compared to sunitinib. Chemical structure of all three compounds consists of 2-naphthyl substituent in position 3 of the dihydropyrazole ring and halogen or methoxy group in position 5 of isatin, suggesting these chemical features are beneficial for the antiproliferative activity of GBM cells. New substituents significantly increased cytotoxicity compared to the effect of sunitinib and  $EC_{50}$  values of tested analogs were up to 250 times lower in U87-MG cell line


and up to 600 times lower in A172 cell line, compared to temozolomide, the main agent of GBM chemotherapy [34].

*In vitro* studies have shown that presence of chlorine in isatin fragment usually results in higher activity of different hybrid molecules against cancer cells [35], as halogen improves permeability to lipid membranes, increasing cellular concentration of compound. Nevertheless, optimizing the position of the substituents is essential for stronger activity. Studies performed on various compounds containing an isatin core have shown that inclusion of substituents at C-5 and C-7 positions of isatin is a promising direction for the development of hybrid molecules [36, 37]. In our case, modifications in C-5 position had clearly greater impact on the cytotoxicity against GBM cells, as EMAC4002 bearing chlorine at C-7 almost completely lost its activity in both cell lines compared to EMAC4001 containing chlorine at C-5. Additionally, inclusion of bromine at C-7 (EMAC4000) was not sufficient to achieve a strong cytotoxic effect on glioma cells, even though, bromo-substituents has shown to be beneficial for different isatin derivatives activity against myeloma and breast cancer cells *in vitro* studies [38, 39]. In our case, presence of iodine at C-5 was more favorable as EMAC4006 showed strong anticancer activity in 2D cell cultures.

## Sunitinib derivatives affect cell migration and spheroid growth



**Figure 6.** Effect of sunitinib analogs on U-87MG spheroid growth at (A, B) 100 nM and (C, D) 500 nM concentrations. The asterisks (\*) indicate  $P < 0.05$  and the double asterisk (\*\*) indicates  $P < 0.01$ , compared to the control. Scale bar = 100  $\mu\text{m}$ . Abbreviations: SNT, sunitinib.



**Figure 7.** Effect of sunitinib analogs on A-172 spheroid growth at (A, B) 100 nM and (C, D) 500 nM concentrations. The asterisks (\*) indicate  $P < 0.05$ , compared to the control. Scale bar = 100  $\mu\text{m}$ . Abbreviations: SNT, sunitinib.

Hypervascularization of GBM depends on endothelial cells migration, which is proven to be regulated by VEGFRs, PDGFRs and their growth

factors [40-42]. Like sunitinib, our tested derivatives contain flat aromatic group essential for binding to the ATP-binding pocket in kinases

[31, 33], thus we hypothesized that hybrid molecules with lipophilic substituents in isatin might have stronger kinase inhibitory effect in GBM cells, compared to sunitinib. Our results showed that addition of chlorine at C-5 position resulted in strong inhibitory effect on cell migration of EMAC4001, which surpassed sunitinib in U-87MG cell line and had similar effect to sunitinib in A-172 cell line. Iodine substituent in EMAC4006 showed advantage in reducing U-87 single cell movement, compared to sunitinib and EMAC4007. EMAC4007 containing methoxy substituent showed reduced activity in U-87MG, however it was the most active compound in A172 cell line. It should be noted that average U-87MG cell velocity remained the same for 6 hours after treatment of EMAC4001 and EMAC4006, while treatment with EMAC4007 resulted in variation in cell velocity. Interestingly, EMAC derivatives had faster effect on U-87MG cell movement, as sunitinib reduced migration only after 3 hours. As it was established that analogs had different effect on single cells movement based on the cell line, it is possible that compounds have different selectivity for kinases expressed in tested glioma cells. There is evidence that U-87MG and A-172 cells express both VEGF receptors, VEGFR-1 and VEGFR-2 [21], however PDGFR- $\alpha$  expression is associated with U-87MG cells and PDGFR- $\beta$  is upregulated in A-172 cells [43].

As expected, new derivatives showed encouraging results in human brain cancer 2D models, however, their effect on 3D structures may be described as inconsistent. EMAC4001 outperformed sunitinib in inhibiting tested GBM cells viability and migration suggesting more potent effect on RTKs essential for GBM development. Unfortunately, new sunitinib derivative did not show a significant effect in GBM spheroids meaning further investigation of permeability is needed to understand the reason for failure. Although, it is not uncommon for new molecules to lose potential activity in 3D cultures as it is proven cancer cells monolayer is more sensitive to traditional chemotherapy [44]. Hypoxia is one of the main features of tumor which is highly associated with activation of different signaling pathways leading to resistance to therapy and has been observed in 200-300  $\mu\text{m}$  diameter spheroids meaning compounds that are active in 3D models are more likely to be active clinically [44, 45].

EMAC4007 was the only compound which not only surpassed sunitinib in cell viability and migration testing, but also was similarly active in 3D cultures of A-172 cells. We discovered that halogen substituents contribute to activity in 2D cultures, but methoxy group is more beneficial in 3D cultures of GBM and can be associated to selectivity for A-172 cell line. However, considering high GBM heterogeneity, the anticancer effect of EMAC4007 alone is likely to be insufficient, thus further studies of new sunitinib derivative effect in combinations with other therapeutic agents are needed.

### Conclusions

Sunitinib analogs possess the anticancer activity in brain cancer U-87MG and A172 cell lines. Compounds had a stronger effect on brain cancer cell viability and migration based on the cell line. However, none of the new tested compounds were more active in 3D cell cultures than sunitinib.

### Disclosure of conflict of interest

None.

**Address correspondence to:** Vilma Petrikaite, Laboratory of Drug Targets Histopathology, Institute of Cardiology, Lithuanian University of Health Sciences, Sukileliu pr. 13, LT-50162, Kaunas, Lithuania. Tel: +370-686-29383; E-mail: vilma.petrikaite@ismuni.lt

### References

- [1] Skalli O, Wilhelmsson U, Orndahl C, Fekete B, Malmgren K, Rydenhag B and Pekny M. Astrocytoma grade IV (glioblastoma multiforme) displays 3 subtypes with unique expression profiles of intermediate filament proteins. *Hum Pathol* 2013; 44: 2081-8.
- [2] Hambardzumyan D and Bergers G. Glioblastoma: defining tumor niches. *Trends Cancer* 2015; 1: 252-65.
- [3] Hanif F, Muzaffar K, Perveen K, Malhi SM and Simjee SU. Glioblastoma multiforme: a review of its epidemiology and pathogenesis through clinical presentation and treatment. *Asian Pac J Cancer Prev* 2017; 18: 3-9.
- [4] Domingo-Musibay E and Galanis E. What next for newly diagnosed glioblastoma? *Future Oncol* 2015; 11: 3273-83.
- [5] Davis ME. Glioblastoma: overview of disease and treatment. *Clin J Oncol Nurs* 2016; 20: S2-8.



## Sunitinib derivatives affect cell migration and spheroid growth

- [6] Brandao M, Simon T, Critchley G and Giamas G. Astrocytes, the rising stars of the glioblastoma microenvironment. *Glia* 2019; 67: 779-90.
- [7] Louca M, Stylianou A, Minia A, Pliaka V, Alexopoulos LG, Gkretsi V and Stylianopoulos T. Ras suppressor-1 (RSU-1) promotes cell invasion in aggressive glioma cells and inhibits it in non-aggressive cells through STAT6 phospho-regulation. *Sci Rep* 2019; 9: 7782.
- [8] Nduom EK, Yang C, Merrill MJ, Zhuang Z and Lonser RR. Characterization of the blood-brain barrier of metastatic and primary malignant neoplasms. *J Neurosurg* 2013; 119: 427-33.
- [9] Noch EK, Ramakrishna R and Magge R. Challenges in the treatment of glioblastoma: multi-system mechanisms of therapeutic resistance. *World Neurosurg* 2018; 116: 505-17.
- [10] Jackson S, Weingart J, Nduom EK, Harfi TT, George RT, McAreavey D, Ye X, Anders NM, Peer C, Figg WD, Gilbert M, Rudek MA and Grossman SA. The effect of an adenosine A2A agonist on intra-tumoral concentrations of temozolomide in patients with recurrent glioblastoma. *Fluids Barriers CNS* 2018; 15: 2.
- [11] Pearson JRD and Regad T. Targeting cellular pathways in glioblastoma multiforme. *Signal Transduct Target Ther* 2017; 2: 17040.
- [12] Sciacaluga M, D'Alessandro G, Pagani F, Ferrara G, Lopez N, Warr T, Gorello P, Porzia A, Mainiero F, Santoro A, Esposito V, Cantore G, Castigli E and Limatola C. Functional cross talk between CXCR4 and PDGFR on glioblastoma cells is essential for migration. *PLoS One* 2013; 8: e73426.
- [13] Pottier C, Fresnais M, Gilon M, Jérusalem G, Longuespée R and Sounni NE. Tyrosine kinase inhibitors in cancer: breakthrough and challenges of targeted therapy. *Cancers (Basel)* 2020; 12: 731.
- [14] Zhang P, Xia Q, Liu L, Li S and Dong L. Current opinion on molecular characterization for GBM classification in guiding clinical diagnosis, prognosis, and therapy. *Front Mol Biosci* 2020; 7: 562798.
- [15] Cantanhede IG and de Oliveira JRM. PDGF family expression in glioblastoma multiforme: data compilation from ivy glioblastoma atlas project database. *Sci Rep* 2017; 7: 15271.
- [16] Thiessen B, Stewart C, Tsao M, Kamel-Reid S, Schaiquevich P, Mason W, Easaw J, Belanger K, Forsyth P, McIntosh L and Eisenhauer E. A phase I/II trial of GW572016 (lapatinib) in recurrent glioblastoma multiforme: clinical outcomes, pharmacokinetics and molecular correlation. *Cancer Chemother Pharmacol* 2010; 65: 353-61.
- [17] Raizer JJ, Abrey LE, Lassman AB, Chang SM, Lamborn KR, Kuhn JG, Yung WK, Gilbert MR, Aldape KA, Wen PY, Fine HA, Mehta M, Dean-gelis LM, Lieberman F, Cloughesy TF, Robins HI, Dancey J and Prados MD; North American Brain Tumor Consortium. A phase II trial of erlotinib in patients with recurrent malignant gliomas and nonprogressive glioblastoma multiforme postirradiation therapy. *Neuro Oncol* 2010; 12: 95-103.
- [18] Jaal J, Kase M, Minajeva A, Saretok M, Adamson A, Junninen J, Metsaots T, Jõgi T, Joonsalu M, Vardja M and Asser T. VEGFR-2 expression in glioblastoma multiforme depends on inflammatory tumor microenvironment. *Int J Inflamm* 2015; 2015: 385030.
- [19] Cao Y. Multifarious functions of PDGFs and PDGFRs in tumor growth and metastasis. *Trends Mol Med* 2013; 19: 460-73.
- [20] Krcek R, Matschke V, Theis V, Adamietz IA, Bühler H and Theiss C. Vascular endothelial growth factor, irradiation, and axitinib have diverse effects on motility and proliferation of glioblastoma multiforme cells. *Front Oncol* 2017; 7: 182.
- [21] Atzori MG, Tentori L, Ruffini F, Ceci C, Lisi L, Bonanno E, Scimeca M, Eskilsson E, Daubon T, Miletic H, Ricci Vitiani L, Pallini R, Navarra P, Bjerkvig R, D'Atri S, Lacal PM and Graziani G. The anti-vascular endothelial growth factor receptor-1 monoclonal antibody D16F7 inhibits invasiveness of human glioblastoma and glioblastoma stem cells. *J Exp Clin Cancer Res* 2017; 36: 106.
- [22] Dimitropoulos K, Giannopoulou E, Argyriou AA, Zolota V, Petsas T, Tsiata E and Kalofonos HP. The effects of anti-VEGFR and anti-EGFR agents on glioma cell migration through implication of growth factors with integrins. *Anticancer Res* 2010; 30: 4987-92.
- [23] Grisanti S, Ferrari VD, Buglione M, Agazzi GM, Liserre R, Poliani L, Buttolo L, Gipponi S, Pederisini R, Consoli F, Panciani P, Roca E, Spena G, Triggiani L and Berruti A; Gruppo Neuro-Onco-logico Bresciano. Second line treatment of recurrent glioblastoma with sunitinib: results of a phase II study and systematic review of literature. *J Neurosurg Sci* 2019; 63: 458-67.
- [24] Kollmannsberger C, Soulieres D, Wong R, Scalera A, Gaspo R and Bjarnason G. Sunitinib therapy for metastatic renal cell carcinoma: recommendations for management of side effects. *Can Urol Assoc J* 2007; 1 Suppl: S41-54.
- [25] Mulet-Margalef N and Garcia-del-Muro X. Sunitinib in the treatment of gastrointestinal stromal tumor: patient selection and perspectives. *Onco Targets Ther* 2016; 9: 7573-82.
- [26] van Linde ME. A phase II/III study of high-dose, intermittent sunitinib in patients with recurrent glioblastoma multiforme: the STELLAR study [Internet]. [clinicaltrials.gov](https://clinicaltrials.gov); 2021 [cited 2022 Apr 26]. Report No.: NCT03025893. Avail-

## Sunitinib derivatives affect cell migration and spheroid growth

- able from: <https://clinicaltrials.gov/ct2/show/NCT03025893>.
- [27] National Cancer Institute (NCI). A randomized phase 2 trial of cediranib and olaparib compared to bevacizumab in patients with recurrent glioblastoma who have not received prior VEGF therapy [Internet]. [clinicaltrials.gov](https://clinicaltrials.gov/ct2/show/NCT02974621); 2022 [cited 2022 Apr 26]. Report No.: NCT02974621. Available from: <https://clinicaltrials.gov/ct2/show/NCT02974621>.
- [28] Wu H. Clinical study of apatinib combined with temozolomide in the treatment of uncontrolled or repeated high-grade gliomas [Internet]. [clinicaltrials.gov](https://clinicaltrials.gov/ct2/show/NCT04253873); 2020 [cited 2022 Apr 26]. Report No.: NCT04253873. Available from: <https://clinicaltrials.gov/ct2/show/NCT04253873>.
- [29] Tao R. Phase II clinical trials on anlotinib for the treatment of recurrent glioblastoma [Internet]. [clinicaltrials.gov](https://clinicaltrials.gov/ct2/show/NCT04004975); 2019 [cited 2022 Apr 27]. Report No.: NCT04004975. Available from: <https://clinicaltrials.gov/ct2/show/NCT04004975>.
- [30] Kumar R, Crouthamel MC, Rominger DH, Gontarek RR, Tummino PJ, Levin RA and King AG. Myelosuppression and kinase selectivity of multikinase angiogenesis inhibitors. *Br J Cancer* 2009; 101: 1717-23.
- [31] AboulMagd AM and Abdelwahab NS. Analysis of sunitinib malate, a multi-targeted tyrosine kinase inhibitor: a critical review. *Microchem J* 2021; 163: 105926.
- [32] Yousefian M and Ghodsi R. Structure-activity relationship studies of indolin-2-one derivatives as vascular endothelial growth factor receptor inhibitors and anticancer agents. *Arch Pharm (Weinheim)* 2020; 353: e2000022.
- [33] Meleddu R, Petrikaite V, Distinto S, Arridu A, Angius R, Serusi L, Škarnulytė L, Endriulaitytė U, Paškevičiu Tė M, Cottiglia F, Gaspari M, Taverna D, Deplano S, Fois B and Maccioni E. Investigating the anticancer activity of isatin/dihydropyrazole hybrids. *ACS Med Chem Lett* 2018; 10: 571-6.
- [34] Karpel-Massler G, Westhoff MA, Kast RE, Dwucet A, Nonnenmacher L, Wirtz CR, Debatin KM and Halatsch ME. Artesunate enhances the antiproliferative effect of temozolomide on U87MG and A172 glioblastoma cell lines. *Anticancer Agents Med Chem* 2014; 14: 313-8.
- [35] Ding Z, Zhou M and Zeng C. Recent advances in isatin hybrids as potential anticancer agents. *Arch Pharm (Weinheim)* 2020; 353: e1900367.
- [36] Vine KL, Locke JM, Ranson M, Benkendorff K, Pyne SG and Bremner JB. In vitro cytotoxicity evaluation of some substituted isatin derivatives. *Bioorg Med Chem* 2007; 15: 931-8.
- [37] Hou Y, Shang C, Wang H and Yun J. Isatin-azole hybrids and their anticancer activities. *Arch Pharm (Weinheim)* 2020; 353: e1900272.
- [38] Panga S, Podila N and Veeresham C. Design, synthesis, characterization, and in vitro evaluation of isatin-pomalidomide hybrids for cytotoxicity against multiple myeloma cell lines. *J Heterocycl Chem* 2018; 55.
- [39] Kumar S, Gu L, Palma G, Kaur M, Singh A, Singh P and Kumar V. Design, synthesis, antiproliferative evaluation and docking studies of 1H-1,2,3-triazole tethered ospemifene-isatin conjugates as selective estrogen receptor modulators. *New J Chem* 2018; 42.
- [40] Wang X, Bove AM, Simone G and Ma B. Molecular bases of VEGFR-2-mediated physiological function and pathological role. *Front Cell Dev Biol* 2020; 8: 599281.
- [41] Horikawa S, Ishii Y, Hamashima T, Yamamoto S, Mori H, Fujimori T, Shen J, Inoue R, Nishizono H, Itoh H, Majima M, Abraham D, Miyawaki T and Sasahara M. PDGFR $\alpha$  plays a crucial role in connective tissue remodeling. *Sci Rep* 2015; 5: 17948.
- [42] Weddell JC, Chen S and Imoukhuede PI. VEGFR1 promotes cell migration and proliferation through PLC $\gamma$  and PI3K pathways. *NPJ Syst Biol Appl* 2017; 4: 1.
- [43] Carrasco-Garcia E, Martinez-Lacaci I, Mayor-López L, Tristante E, Carballo-Santana M, García-Morales P, Ventero Martin MP, Fuentes-Baile M, Rodríguez-Lescure Á and Saceda M. PDGFR and IGF-1R inhibitors induce a G2/M arrest and subsequent cell death in human glioblastoma cell lines. *Cells* 2018; 7: 131.
- [44] Däster S, Amatruda N, Calabrese D, Ivanek R, Turrini E, Droesser RA, Zajac P, Fimognari C, Spagnoli GC, Iezzi G, Mele V and Muraro MG. Induction of hypoxia and necrosis in multicellular tumor spheroids is associated with resistance to chemotherapy treatment. *Oncotarget* 2017; 8: 1725-36.
- [45] Jing X, Yang F, Shao C, Wei K, Xie M, Shen H and Shu Y. Role of hypoxia in cancer therapy by regulating the tumor microenvironment. *Mol Cancer* 2019; 18: 157.