

Review Article

Hearing loss during chemotherapy: prevalence, mechanisms, and protection

Lexie Kessler^{1*}, Chail Koo^{1*}, Claus-Peter Richter^{1,2,3,4}, Xiaodong Tan^{1,2}

¹Department of Otolaryngology-Head and Neck Surgery, Feinberg School of Medicine, Northwestern University, Chicago, Illinois 60611, USA; ²Hugh Knowles Center for Clinical and Basic Science in Hearing and Its Disorders, Northwestern University, Evanston, Illinois 60208, USA; ³Department of Biomedical Engineering, Northwestern University, Evanston, Illinois 60208, USA; ⁴Department of Communication Sciences and Disorders, Northwestern University, Evanston, Illinois 60208, USA. *Equal contributors and co-first authors.

Received May 15, 2024; Accepted August 23, 2024; Epub September 25, 2024; Published September 30, 2024

Abstract: Ototoxicity is an often-underestimated sequela for cancer patients undergoing chemotherapy, with an incidence rate exceeding 50%, affecting approximately 4 million individuals worldwide each year. Despite the nearly 2,000 publications on chemotherapy-related ototoxicity in the past decade, the understanding of its prevalence, mechanisms, and preventative or therapeutic measures remains ambiguous and subject to debate. To date, only one drug, sodium thiosulfate, has gained FDA approval for treating ototoxicity in chemotherapy. However, its utilization is restricted. This review aims to offer clinicians and researchers a comprehensive perspective by thoroughly and carefully reviewing available data and current evidence. Chemotherapy-induced ototoxicity is characterized by four primary symptoms: hearing loss, tinnitus, vertigo, and dizziness, originating from both auditory and vestibular systems. Hearing loss is the predominant symptom. Amongst over 700 chemotherapeutic agents documented in various databases, only seven are reported to induce hearing loss. While the molecular mechanisms of the hearing loss caused by the two platinum-based drugs are extensively explored, the pathways behind the action of the other five drugs are primarily speculative, rooted in their therapeutic properties and side effects. Cisplatin attracts the majority of attention among these drugs, encompassing around two-thirds of the literature regarding ototoxicity in chemotherapy. Cisplatin ototoxicity chiefly manifests through the loss of outer hair cells, possibly resulting from damages directly by cisplatin uptake or secondary effects on the stria vascularis. Both direct and indirect influences contribute to cisplatin ototoxicity, while it is still debated which path is dominant or where the primary target of cisplatin is located. Candidates for hearing protection against cisplatin ototoxicity are also discussed, with novel strategies and methods showing promise on the horizon.

Keywords: Inner ear, cochlea, cisplatin, ototoxicity, hearing protection

Introduction

Antineoplastic or chemotherapeutic drugs for cancer treatment can also harm normal tissues and cells, leading to various adverse effects [1]. Ototoxicity is a notable sequela of cancer treatment, impacting hearing and balance with symptoms including hearing loss, vertigo, dizziness, and tinnitus [2-5]. The severity of these symptoms varies and is influenced by factors including the age of the patient, the specific chemotherapeutic agent used, dosage, and administration method [1, 4, 5]. Although ototoxicity in chemotherapy is not life-threatening, its consequences, such as communication dif-

iculties, social isolation, depression, and fatigue, can significantly impair the quality of life [6, 7]. Furthermore, hearing loss has been suggested as a significant modifiable risk factor for dementia [8]. In pediatric patients, hearing loss is even more devastating as it can delay the development of speech and language abilities, communication skills, and impede cognitive maturation [9].

To determine the ototoxic potential and mechanisms of chemotherapeutic drugs and shed light on emerging treatment compounds and approaches, we performed a comprehensive review of literature and databases on cancer

Hearing loss during chemotherapy

Table 1. Ototoxic chemotherapeutic drugs and their incidence of symptoms

Drug name	Cisplatin	Carboplatin	Vinblastine	Vincristine	Dasatinib	isotretinoin	Tretinoin	
Drug Class	Alkylating agent		Plant alkaloid	Vinca alkaloid	Tyrosine kinase inhibitor	Retinoid	Retinoid	
Antineoplastic Mechanism	DNA binding and cross-linking Protein cross-linking		microtubule inhibitor		Src kinase inhibitor	microtubule inhibitor		
Ototoxic Mechanism	ROS overload, proinflammatory cytokine production		Possible synergy with other drugs	Unclear. Possibly associated with their antitumor activity				
Incidence rate	Hearing loss	31%	12%	10-29%	No data	0.1-1%	No data	6%
	Tinnitus	31%	12%	No data	No data	1-10%	No data	No data
	Dizziness	No data	No data	Rare	No data	1-10%	No data	20%
	Vertigo	No data	No data	Rare	No data	0.1-1%	No data	No data

treatment drugs. A PubMed search revealed nearly 2,000 publications on chemotherapy-associated ototoxicity in the last decade. Among these publications, two-thirds are related to platinum-based agents used in 10-20% of the chemotherapy regimens. However, the mechanisms underlying the ototoxicity of chemotherapeutic drugs remain unclear, and effective strategies to mitigate this side effect are still in development. This review discusses the tentative mechanisms for ototoxicity and advances in hearing protection during cancer treatment, highlighting promising drugs and methodologies recently developed. Our objective is to provide clinicians who treat cancer with a foundational understanding of ototoxicity in chemotherapy. We also offer researchers insight into in-depth mechanisms and potential innovative protective strategies against it. Currently, sodium thiosulfate (STS) is the only FDA-approved drug to mitigate hearing loss during cancer treatment in pediatric patients [10].

Chemotherapeutic drugs affecting the inner ear

To identify compounds/drugs with potentially damaging effects on the inner ear during cancer therapy, we searched the online databases from the websites of the National Cancer Institute (NCI) [2], Chemocare [3], and Beaumont [4] using the search terms hearing loss, tinnitus, dizziness, and vertigo. Out of the approximately 700 compounds used in cancer treatment, only seven have listed hearing loss as a side effect (Table 1) [2-4].

Ninety-seven drugs reported dizziness as a side effect but not hearing loss or tinnitus (Table 2). While dizziness may originate from

the inner ear, it can also have other origins, such as the central nervous system. Since the mechanisms underlying dizziness are multifactorial, and our primary focus is on hearing loss, drugs causing only dizziness but not any other symptoms of ototoxicity are not further explored in this review. Some other drugs, such as doxorubicin (showing ototoxic effects in animal studies [11], but clinical evidence is missing) and nitrogen mustard (strongly restricted use as a chemical weapon), are also excluded.

Platinum-based drugs. The two platinum-based alkylating agents, cisplatin (also referred to as cis-diamminedichloroplatinum (II), CDDP, and platinol) and carboplatin (paraplatin), are the most commonly reported compounds that cause hearing loss. According to the National Cancer Institute (NCI), cisplatin or similar platinum-based drugs are used in 10-20% of cancer chemotherapy regimens [12]. The most common indications include testicular, ovarian, cervical, bladder, and head and neck cancers. Multiple well-documented significant sequelae are nausea, vomiting, liver damage, kidney failure, hearing loss, tinnitus, and vertigo [13-17]. The prevalence of hearing loss during chemotherapy with cisplatin can be as high as 60-80% [14, 18]. Based on the NCI estimated cancer patients of 2018, cisplatin may cause approximately 100-300 thousand new cases of ototoxicity annually in the United States. Oxaliplatin, a third-generation platinum-based chemotherapy drug, has not been considered ototoxic based on a clinical trial with 18 patients [19]. However, case reports showed that the drug might cause hearing loss [20-23].

Vinblastine (velban, alkaban-AQ) and Vincristine are plant alkaloids listed as ototoxic drugs,

Hearing loss during chemotherapy

Table 2. Chemotherapeutic drugs causing only dizziness but not other symptoms of ototoxicity

Dizziness (incidence rate > 30%)		
<i>Generic/Other Name</i>	<i>Brand Name</i>	<i>Mechanism of Action</i>
Axicabtagene Ciloleuceel	Yescarta	chimeric antigen receptor T-cell immunotherapy agent, binds CD19 B-lymphocyte antigen
Entrectinib	Rozlytrek	protein-tyrosine kinase inhibitor, inhibits tropomyosin receptor tyrosine kinases - inhibits high affinity nerve growth factor receptor, BDNF/NT-3 growth factor receptors, NT-3 growth factor receptor, proto-oncogene tyrosine-protein kinase ROS, tyrosine-protein kinase JAK2
Tisagenlecleucel	Kymriah	chimeric antigen receptor T-cell immunotherapy agent, binds CD19-expressing cells and promotes T-cell expansion, activation, target cell elimination
Larotrectinib	Vitrakvi	tropomyosin receptor kinase (TRK) inhibitor, inhibits TRKA, TRKB, TRKC preventing neurotrophin-Trk interaction and Trk activation inducing apoptosis and inhibition of cell growth
Dizziness (incidence rate 10-29%)		
<i>Generic/Other Name</i>	<i>Brand Name</i>	<i>Mechanism of Action</i>
13-cis-Retinoic Acid	Accutane, Isotretinoin	retinoid, acts on nuclear receptors RAR or RXR
5-Azacitidine, Azacitidine	Vidaza, Onureg	antimetabolite and demethylating agent
Abemaciclib	Verzenio	cyclin-dependent kinase inhibitor (CDK4 and CDK6), arrests G1 to S phase
Brentuximab vedotin	Adcetris	CD30-direct antibody drug conjugate (monoclonal antibody that disrupts microtubules)
Ado-Trastuzumab Emtansin	Kadcyla	Anti-HER2 monoclonal antibody combined with microtubule inhibitor DM1 (maytansine derivative)
Anagrelide	Agrylin	phospholipase A2 inhibitor (prevents maturation of megakaryocytes)
Hydrocortisone, Hydrocortone Phosphate, Ala-Cort, Cortisone, Hydrocortisone Sodium Succinate, Hydrocortisone Sodium Phosphate	Solu-Cortef, Hydrocort Acetate, Lanacort	glucocorticosteroid
Aldesleukin, Interleukin-2, IL-2	Proleukin	cytokine, increases production of T lymphocytes and NK cells and improves function of lymphokine-activated killer cells and tumor-infiltrating lymphocytes
Alemtuzumab	Campath	CD52 monoclonal antibody
All-Trans Retinoic Acid, Tretinoin	Vesanoid	Retinoid, acts on nuclear receptors RAR or RXR
Interferon Alfa, alpha interferon, IFN-alpha	Intron A, Roferon-A	cytokine and biologic response modifier
Altretamine, Hexamethylmelamine, HMM	Hexalen	alkylating agent, hydrazine and triazine
Amifostine	Ethylol	chemoprotective agent, deactivates harmful components of chemotherapy drugs; scavenger, binds free radicals produced by cisplatin or radiation therapy
Aminoglutethimide	Cytadren	adrenal cortex corticosteroid production inhibitor, decreases production of estrogens and androgens
Nilutamide	Nilandron, Anandron	antiandrogen, blocks androgen/testosterone receptors
Apalutamide	Erleada	antiandrogen, blocks androgen/testosterone receptors
Arabinosylcytosine, Cytarabine, Ara-C	Cytosar-U	antimetabolite, inhibits DNA polymerase beta; cross-linking/alkylation of DNA, blocks G1/S
Nelarabine	Arranon	antimetabolite, adenosine deaminase inhibitor, incorporates into and destabilizes DNA, inhibits DNA polymerase alpha catalytic subunit; S phase-specific arrest
Arsenic Trioxide	Trisenox	not well understood, DNA fragmentation in leukemia cells; also damages or degrades the fusion protein PML-RAR
Avapritinib	Ayvakit	tyrosine kinase inhibitor, small molecule inhibitor of platelet-derived growth factor receptor alpha (PDGFR-A), targets PDGFRA and PDGFRA D842 mutants as well as multiple KIT mutations

Hearing loss during chemotherapy

Bevacizumab	Avastin, Mvasi, Zirabrev	targets and inhibits VEGF preventing angiogenesis
Belinostat	Beleodaq	histone deacetylase inhibitor
Carmustine, BCNU	BiCNU, Gliadel wafer	alkylating agent, nitrosurea, cross-links DNA and RNA
Binimetinib	Mektovi	oral MEK inhibitor, targets MEK1 and MEK2 protein kinase; usually given with a BRAF kinase inhibitor
Blinatumomab	Blincyto	bispecific T-cell engager monoclonal antibody, induces T-cells to bind CD19 on surface of B-cell leukemia or lymphoma cells
Bortezomib	Velcade	proteasome inhibitor, inhibits 26S proteasome, inhibits proteasome subunit beta type-5 and type-1; cell arrest in G2-M phase; in multiple myeloma, works by blocking adhesion molecule activation
Bosutinib	Bosulif	kinase inhibitor, targets ABL and SRC kinases
Encorafenib	Braftovi	BRAF kinase inhibitor
Busulfan	Busulfex, Myleran	alkylating agent, alkylsulfonate
Cabozantinib	Cometriq, Cabometyx	oral receptor tyrosine kinase inhibitor (RET, MET, VEGF), blocks cell division pathways
Carfilzomib	Kyprolis	tetrapeptide epoxyketone proteasome inhibitor, irreversibly binds to N-terminal threonine-containing active sites of 20S proteasome
Crizotinib	Xalkori capsules	oral receptor tyrosine kinase inhibitor, inhibits ALK, hepatocyte growth factor receptor (HGFR, c-Met) and receptor d'origine natais (RON); blocks cell division pathway
Decitabine	Dacogen	antimetabolite and demethylating agent, restores function of tumor suppressor genes and cytotoxic effect on rapidly dividing cells
Daratumumab and Hyaluronidase	Darzalex Faspro	CD38 monoclonal antibody (present on myeloma cells), IgG1k human monoclonal antibody binds CD38 and induces apoptosis through mediated cross linking and immune mediated tumor cell lysis through complement dependent cytotoxicity, antibody mediated cytotoxicity, and antibody dependent cellular phagocytosis; hyaluronidase helps with absorption into blood
Daunorubicin and Cytarabine (Liposomal)	Vyxeos	Daunorubicin is an anthracycline, intercalates between DNA base pairs inhibiting DNA synthesis and DNA-dependent RNA synthesis; Cytarabine is an antimetabolite (S phase); liposome helps with drug distribution and lengthens time of effect of the drug allowing for extended treatment effect
Glasdegib	Daurismo	hedgehog pathway inhibitor, inhibits increases in tumor size and decrease the amount of CD45+/CD33+ cells in the bone marrow; binds and inhibits Smoothened (SMO) receptor
Dexamethasone, Dexamethasone Sodium Phosphate, Dexamethasone Acetate	Decadron, Dexasone, Diodex, Hexadrol, Maxidex	glucocorticosteroid
Prednisolone	Delta-Cortef, Orapred, PEDIAPRED, Prelone	glucocorticosteroid
Prednisone	Deltasone, Liquid Pred, Meticorten, Orasone	glucocorticosteroid
Denileukin Diftitox	Ontak	biologic response modifier agent, a fusion protein (combination of diphtheria toxin and IL-2), selectively delivers the cell-killing activity of diphtheria toxin to targeted cells; binds to lymphoma cells that express high affinity IL-2 receptor (IL-2 part of fusion protein binds cell surface) and halts protein synthesis
Dexrazoxane	Zincard	chemoprotectant agent, binds free radicals formed by doxorubicin; extravasation antidote, binds chemotherapy drug that leaked from vein preventing damage to surrounding tissue

Hearing loss during chemotherapy

Methylprednisolone	Duralone, Medrol, Medralone, M-Prednisol, Solu-Medrol	glucocorticosteroid
Eculizumab	Soliris	monoclonal antibody, binds C5 complement protein preventing formation of MAC, prevents hemolysis and stabilizes hemoglobin
Eltrombopag	Promacta	colony stimulating factor, thrombopoietic agent, growth factor that stimulates platelet production by binding to and activating the thrombopoietin (TPO) receptor; a thrombopoietin nonpeptide agonist
Tagraxofusp-erzs	Elzonris	biologic response modulator and cytokine, combination of recombinant human IL-3 and truncated diphtheria toxin, binds CD123 (alpha chain of IL-3 receptor) and delivers diphtheria toxin to cells, blocks protein synthesis; binds ADP-ribosylation factor-like protein 2
Fam-trastuzumab deruxtecan-nxki	Enhertu	anti-HER2 monoclonal antibody (anti-HER2 IgG1) combined with a topoisomerase I inhibitor; antibody attached to chemotherapy, allows selective delivery into HER2 overexpressing cells, DNA damage
Enzalutamide	Xtandi	antiandrogen (second generation), blocks androgen/testosterone receptors
Toremifene	Fareston	anti-estrogen, estrogen receptor antagonist, blocks estrogen binding and uptake into cells
Gilteritinib	Xospata	protein-tyrosine kinase inhibitor, inhibits FLT3 receptor, serotonin receptors, TP53, and ALK tyrosine kinase receptor
Trastuzumab	Herceptin (Biosimilars: Herzuma, Kanjinti, Ogivri, Ontruzant)	HER2/neu receptor monoclonal antibody
Ibritumomab, Ibritumomab Tiuxetan	Zevalin	CD20 monoclonal antibody linked with Yttrium-90 (radioactive substance), directly delivers radiation to CD20+ cells
Ibrutinib	Imbruvica	binds to and inhibits the bruton's tyrosine kinase (BTK) signaling molecule of the B-cell receptor signaling complex
Ponatinib	Iclusig	tyrosine kinase inhibitor
IL-11, Oprelvekin, Interleukin-11	Neumega	biologic response modifier and cytokine, stimulates production, maturation and activation of platelets
Talimogene Laherparepvec, T-VEC	Imlygic	genetically modified weakened form of live HSV (oncolytic), replicates within tumors and produces GM-CSF to promote anti-tumor immune response
Interferon Alfa-2b (PEG Conjugate), PEG Interferon	PEG-Intron	biologic response modifier and cytokine, activates human type 1 interferon causing them to dimerize which activates JAK/STAT pathway
Ruxolitinib	Jakafi	oral receptor tyrosine kinase inhibitor, inhibits JAK1 and JAK2
Pembrolizumab	Keytruda	highly selective humanized monoclonal IgG4 antibody directed against the PD-1 receptor on cell surface, prevents binding and activation of PD-L1 and PD-L2 which activates T-cell mediated immune response against tumor cells
Lenalidomide	Revlimid	immunomodulatory agent and antiangiogenic agent, inhibits protein cereblon and TNF ligand superfamily member 11, antagonizes cadherin-5, negative modulator of prostaglandin G/H synthase 2
Lenvatinib	Lenvima	oral receptor tyrosine kinase inhibitor, inhibits VEGF, VEGFR, FGF, PDGFR alpha, KIT and RET
Lorlatinib	Lorbrena	reversible tyrosine kinase inhibitor, blocks abnormal ALK protein
Luspatercept	Reblozyl	recombinant fusion protein, hematopoiesis agent (contains modified form of the extracellular domain of human activin receptor), binds and inhibits transforming growth factor beta super family molecules increasing expression of blood cell precursors

Hearing loss during chemotherapy

Olaparib	Lynparza	poly (ADP-ribose) polymerase (PARP) enzyme inhibitor (PARP1, PARP2, PARP3), induces synthetic lethality in BRCA1/2 deficient tumor cells through formation of double-strand DNA breaks
Procarbazine	Matulane	alkylating agent, hydrazine and triazine
Midostaurin	Rydapt	tyrosine kinase inhibitor, inhibits FLT3 inhibiting leukemic cell production
Niraparib	Zejula	PARP inhibitor, highly selective for PARP1 and PARP2 resulting in DNA damage and apoptosis, induces cytotoxicity in tumor cell lines w/ and w/o BRCA1/2 deficiencies
Romiplostim	Nplate	biologic response modifier, colony stimulating factor, promotes platelet production via the thrombopoietin receptor
Pertuzumab	Perjeta	HER2 monoclonal antibody (binds different area of HER2 protein than trastuzumab)
Pomalidomide	Pomalyst	thalidomide analogue, inhibits protein cereblon, TNF and prostaglandin G/H synthase 2
Sipuleucel-T	Provenge	autologous cellular immunotherapy, selectively targets prostatic acid phosphatase (PAP), a PSA
Rucaparib	Rubraca	PARP inhibitor (PARP1, PARP2, PARP3), increases formation of PARP-DNA complexes; cytotoxicity in BRCA1/2 deficient tumor cell lines and other DNA repair genes
Sacituzumab Govitecan-hziy	Trodelyv	trop-2-directed antibody-drug conjugate combined with a topoisomerase I inhibitor (SN-38) attached by a linker, binds trop-2-expressing cancer cells and is internalized, SN-38 is released in cancer cell by breaking the linker
Sunitinib, SU11248	Sutent	receptor protein-tyrosine kinase inhibitor, inhibits VEGF
Talazoparib	Talzenna	PARP inhibitor (PARP1, PARP2), strong catalytic inhibition and a PARP-trapping potential
Temozolomide	Temodar	alkylating agent, hydrazine and triazine (similar to dacarbazine, acts as a pro-drug)
Vorinostat	Zolinza	not fully understood, histone deacetylase inhibitor, inhibits HDAC2, HDAC2, HDAC3, HDAC6
Zoledronic Acid	Zometa	bisphosphonate, decreases osteoclast actions on bone

albeit with little evidence in the literature. As a microtubule inhibitor, vinblastine-induced hearing loss was observed in two case reports of patients with Hodgkin's lymphoma who underwent combined chemotherapy, including doxorubicin, bleomycin, vinblastine, and dacarbazine (ABVD) [24, 25]. In one published case, vinblastine was given in a combination treatment with vincristine. It has been discussed that the observed cochlear damage [24] originated from vinblastine. However, it is also possible that the damage might have been caused by vincristine, a commonly used vinca alkaloid similar to vinblastine [26-28].

Dasatinib (Sprycel), a tyrosine kinase inhibitor targeting multiple cancer cells, is also reported to cause hearing loss in case reports but the incidence is rare (0.1-1%). Instead, more frequent side effects of dasatinib are tinnitus and dizziness.

Isotretinoin (cis-retinoic acid) and tretinoin (all-trans retinoic acid) are retinoids that can slow down the growth of cancer cells. Occasional case reports indicate that both can induce hearing loss [29, 30], while detailed studies on isotretinoin's and tretinoin's ototoxicity are still missing. Furthermore, isotretinoin is mainly used to treat severe acne. In two clinic studies with cohorts of acne patients, isotretinoin treatment leads to transient hearing improvement instead [31, 32]. These controversial results suggest their influence on the auditory system, while more studies are needed to determine the mechanisms.

Mechanisms of hearing loss in chemotherapy

The working theory for the mechanism of platinum-based drugs is related to the generation of reactive oxygen species (ROS) and DNA damage. However, their ototoxicity is mainly associated with excessive ROS generation [33, 34], while DNA damage [35, 36] likely plays a minor role. However, the mechanisms of other ototoxic chemotherapeutic drugs listed in **Table 1** remain largely unclear because the studies are rare. Some studies have speculated that it is associated with their antitumor activity [24]. Due to the availability of literature, this review focuses mainly on the ototoxicity of platinum compounds, particularly cisplatin.

Inner ear structures targeted by cisplatin

Structural and functional features of the cochlea: The cochlea comprises three fluid-filled tubes, scala tympani, scala media, and scala vestibuli (**Figure 1**). Scala vestibuli and scala media are separated by Reissner's membrane, while scala media and scala tympani by the basilar membrane and reticular lamina. The organ of Corti is located on the basilar membrane. It contains two types of hair cells: the inner hair cells (IHCs) and outer hair cells (OHCs). IHCs and OHCs have hair bundles, or stereocilia, with ion channels responsible for transforming sound-induced vibrations of soft tissue structures of the inner ear into action potentials. The channels are named mechano-electrical transduction (MET) channels, of which the core domain is recently identified as transmembrane channel-like protein 1 [37]. Opening of MET channels in OHCs results in a depolarizing current enhancing the sound-induced vibrations of soft tissue structures by changing the stiffness and length of the OHCs. In IHCs, the opening of the MET channels releases neurotransmitters and generates action potentials. The MET current is primarily carried by calcium and potassium ions [38]. Please refer to [39-41] for recent reviews on the MET channels.

The driving force for the MET current is the voltage difference between the endocochlear potential (EP) of about 80 mV [42, 43] and the resting potential of the hair cells. Intracochlear ion homeostasis, combined with the selective permeability for different ions across the boundaries of the scalae and active ion transport across stria vascularis, generates the EP. The perilymphatic ion concentration, high in sodium (~150 mM) and low in potassium (~5 mM), is similar to the ion concentration of the extracellular fluids. This differs from the endolymphatic ion concentration, low in sodium (~15 mM) and high in potassium (~140 mM), which is comparable with the ion composition of the cytoplasm [44]. The ion homeostasis of the endolymph is primarily controlled by the stria vascularis (SV), a structure lining the lateral wall of the scala media. The marginal cells in the SV regulate the potassium concentration [45], and the intermediate cells form tight junctions with basal cells to separate the en-

Hearing loss during chemotherapy




Figure 1. Schematic image showing the basic cochlear structures and the targets of cisplatin ototoxicity. Cochlear duct forms three scalae, scala vestibuli, scala media, and scala tympani, separated by the basilar membrane (BM) and Reissner's membrane (RM), respectively. Cells affected by cisplatin include the outer hair cells (OHC) and inner hair cells (IHC) in the organ of Corti (OC), basal cells (BC), intermedial cells (IC), and marginal cells (MC) in the stria vascularis (SV), and the spiral ganglion neurons (SGN) in the modiolus. Whether OHC loss is initiated directly through cisplatin uptake or indirectly through the drop of endocochlear potential (EP) following SV damage is still debating. Refer to the text for more details. TM: tectorial membrane; RM: Reissner's membrane. All images are created with BioRender.com.

dolymph from the surrounding perilymph (Figure 1, for recent reviews, see [46, 47]).

This blood-labyrinthine barrier (BLB) is similar to the blood-brain barrier (BBB) in terms of cellular and molecular basis [48], which is only permeable to some small molecules (up to 500 KDa) [49]. Unfortunately, this permeability to the BLB includes most ototoxic drugs, while it excludes most otoprotective agents proven effective *in vitro* [50, 51]. Furthermore, once inside the cochlea, the outflow of the ototoxic drugs is also difficult.

Path to hearing loss in chemotherapy: Cisplatin ototoxicity leads to structural changes, including shrinkage and inflammation of SV, loss of

OHCs and IHCs, morphological changes of the stereocilia bundles, IHC synaptopathy, changes of supporting cells, and loss of spiral ganglion neurons (SGNs) (for reviews, see [15, 52-56]). Recent cellular and molecular biology studies also revealed additional cells and structures affected by cisplatin ototoxicity, such as the spiral ligament [57, 58], spiral limbus, spiral modiolar veins and lacunae [15, 59], and pericytes in the SV [49, 60]. Functional changes include decreased EP, threshold elevation of distortion product otoacoustic emissions (DPOAEs), auditory brainstem responses (ABRs), and compound action potentials (CAPs) [61-63]. DPOAE magnitudes and ABR wave-I amplitudes are also decreased. These changes are closely re-

lated to the cellular processes, including ROS generation, inflammation, and apoptosis [64]. Meanwhile, the inflammatory and apoptotic mechanisms of cisplatin ototoxicity are highly interconnected, which leads to a vicious cycle of inflammation, ROS production, nuclear and mitochondrial DNA damage, ER stress, and cell death, which will be discussed later.

Following cisplatin therapy, OHC loss is usually the most prominent and severe cochlear damage [46] and is likely the cause of permanent hearing loss. How OHCs are affected by cisplatin remains unclear (**Figure 1**). While early studies suggest direct damage of the OHCs following the uptake of cisplatin via different ion channels [65] or transporters [66, 67], conclusive experimental evidence of whether platinum exists in the hair cells is still missing. OHC loss might also originate from changes in the EP resulting from compromised SV function [46, 68]. This view is further supported by the following findings: (1) The highest platinum accumulation is found in the SV [20, 68, 69]. Solid evidence comes from Cunningham's group, which showed that platinum mainly accumulates in SV after chemotherapy, using the inductively coupled plasma mass spectrometry visualization technique [68]. (2) The immediate decrease of the EP after cisplatin treatment suggests the involvement of the stria vascularis in cisplatin ototoxicity [46]. (3) Platinum-DNA adduct was observed in SV marginal cells as early as 8 hours after cisplatin treatment, while ROS accumulation was not shown even after 48 hours [69]. This finding suggests that ROS accumulation might be a secondary effect of SV damage. (4) A recent study shows in cell culture that the pericytes in SV, which are also critical for EP, are the targets of cisplatin and may account for BLB breakdown in CIHL [60].

Cisplatin uptake by cochlear cells

The first step for cisplatin ototoxicity is its uptake by cochlear cells, which may occur, like in other tissues, through the organic cation transporter 2 (OCT2) [67], copper transporter 1 (CTR1) [66, 70], and LDL receptor-related protein 2 (LRP2) [71]. The high expression of CTR1 and OCT2 in IHCs, OHCs, spiral ganglion neurons (SGN), and SV [66] supports this view (for review, see [56, 66, 72]). The interaction and

uptake of cisplatin via mechano-transducer (MET) channels of the hair cells have also been studied [65, 73]. One early study shows that cisplatin blocks MET channels in chicken cochlear hair cells as an acute effect [73]. However, the paper has not demonstrated whether cisplatin could pass the MET channels. A study on the zebrafish lateral line organ shows that functionally intact MET channels are required for the toxicity of cisplatin to the hair cells. Hair cell death is prevented when the MET channels are non-functional through chemical blockage or mutation [65]. Interestingly, the same publication also suggests that the roles of OCT2 and CTR1 for cisplatin uptake are insignificant in these cells [65]. A more recent study on murine cochlear hair cell explants, however, shows that both MET channel and OCT are involved in the uptake of fluorescent dye-conjugated cisplatin [74]. In addition to transporters and MET channels, cisplatin may also enter cells through passive diffusion [75], as shown in the digestive system of rats [76] and a cochlear-derived cell line OC-k3 [77]. In the *in vitro* studies on OC-k3 cells, cisplatin enters the cells via first-order kinetics without saturation before the induction of cell death. Once inside the cell, cisplatin undergoes an aquation reaction and hydrolyzes as water ligands displace chloride ligands [75]. The ability of this activated and positively charged aqua-cisplatin compound to passively diffuse back across the plasma membrane is significantly decreased. The drug is then trapped intracellularly, leading to unimpeded damage [72, 78, 79].

ROS generation and its central role in cisplatin ototoxicity

Platinum has a high affinity for sulfur ligands [80]. Once it enters the cochlea, the highly reactive aquated form of cisplatin binds to both DNA and proteins [79], which triggers a series of signaling pathways (**Figure 2**). Cisplatin may induce NADPH oxidase 3 (NOX3) directly [81] and/or via cisplatin activated transient receptor potential cation channel subfamily V member 1 (TRPV1) channel, followed by calcium influx [82]. Cisplatin-activated NOX3 leads to the generation of ROS, specifically superoxide ($O_2^{\cdot-}$) [81]. $O_2^{\cdot-}$ can induce mitochondrial translocation of the B-cell lymphoma 2 gene (Bcl2) associated X (Bax), leading to mitochondrial-

Hearing loss during chemotherapy




Figure 2. Schematic image showing intracellular signaling pathways associated with cisplatin ototoxicity in the cochlea. Different cell death pathways, including apoptosis (either mitochondria dependent or independent) and non-apoptotic cell death pathways are involved and interconnected. Refer to the text for details.

dependent apoptosis (further discussed below) [72, 83]. Another study demonstrates that NOX3-dependent ROS generation, rather than cisplatin itself, activates TRPV1 channels [84] (for review, see [72, 85]). Nevertheless, both NOX3 induction and TRPV1 activation by cisplatin can result in ROS generation and calcium influx. In addition, reducing TRPV1 expression [84] and knockout of NOX3 [82] both attenuate cisplatin-induced hearing loss (CIHL), during which both SGNs and OHCs are protected [82]. ROS is also a potent inducer of NOX3, which is over 50 times more abundant in the inner ear compared to other tissues [81] (for review, see [79]). It leads to a vicious cycle of ROS-induced TRPV1 and NOX3 activation, causing further calcium influx, ROS production, and TRPV1 and NOX3 activation [84].

ROS is detoxified by an intracellular antioxidant defense system composed of antioxidant enzymes, including superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GSH.Px), and glutathione reductase (GR) [13, 56, 72]. This antioxidant defense system can be rapidly overwhelmed by cisplatin [80, 86].

The reactive platinum moiety of aqua-cisplatin reacts with intracellular oxygen molecules, generating $O_2^{\cdot-}$ [56]. SOD converts $O_2^{\cdot-}$ to hydrogen peroxide (H_2O_2), which CAT further catalyzes to produce water and oxygen [56, 72]. In this process, GSH.Px converts glutathione from the reduced form (GSH) to the oxidized form (GSSG) during the conversion of H_2O_2 to water. GR catalyzes the conversion of GSSG back to GSH using NADPH as a cofactor [56, 72]. Cisplatin can covalently bind to the thiol group of these antioxidant enzymes, leading to their inactivation [56]. This reactive cisplatin compound also directly binds to GSH, leading to its excretion or conversion back to GSSG [55]. NADPH depletion downregulates GR antioxidant activity, which further decreases GSH levels. Depletion of GSH downregulates GSH.Px activity [56]. Most experimental evidence shows that antioxidant enzymes are reduced by cisplatin, some to the extent of 50-70% [55, 86, 87] (for reviews, see [56, 72, 88, 89]). However, other studies indicate that SOD and CAT activities can be increased instead [55, 79, 90]. Together, antioxidant depletion and ROS over-

load trigger a series of interconnected cell death pathways involving DNA damage, lipid peroxidation, protein oxidation and enzyme inactivation, ion channel expression changes, endoplasmic reticulum (ER) stress, and inflammation [56, 59, 72, 83]. The cascade of ROS signaling involves processes in different compartments of the cells, including the cytoplasm, nucleus, and mitochondria (**Figure 2**). Different cell death processes are involved, including mitochondrial-dependent and mitochondrial-independent apoptosis, necroptosis, autophagy, ferroptosis, etc. (**Figure 2**).

Mitochondrial-dependent apoptosis associated with cisplatin ototoxicity

Mitochondrial-dependent apoptosis is characterized by releasing pro-apoptotic factors from the mitochondria to cytosol. This process can be triggered by, and is associated with, different factors and pathways during cisplatin treatment, including lipid peroxidation, DNA damage, signal transducer and activator of transcription 1 (STAT1), p53 activation, Bcl2/Bax activation, cyclin/cyclin-dependent kinase (CDK) activation, Cytochrome C, Caspases, etc. In the cochlea, mitochondrial dysfunction and apoptosis due to ROS overload and disruption of intracellular redox homeostasis result in the loss of OHCs and other cells [91, 92] (for review, see [13]) (**Figure 2**). Most of the factors and pathways related to mitochondrial-dependent apoptosis have been studied. They are shown to be involved in cisplatin ototoxicity. First, cisplatin can integrate into DNA in the nucleus via adduct formation [13, 16, 69, 93], leading to cross-linking and damage [93]. DNA damage activates p53, initiating the intrinsic mitochondrial apoptosis pathway [94]. P53 increases the expression of the pro-apoptotic molecule Bax [78], which is translocated to the mitochondria, where it permeates the outer mitochondrial membrane. Mitochondrial membrane permeabilization leads to the loss of the mitochondrial membrane potential, mitochondrial ROS (mtROS) production, and release of Cytochrome C and mtROS from the mitochondria into the cytoplasm [95, 96]. Second, both NOX3-dependent ROS production [57, 79, 85, 88, 97] and ROS-mediated activation of extracellular signal-regulated kinases 1 (ERK1) [59, 64] activate STAT1 signaling. STAT1 triggers p53 activation, leading to the above-described

mitochondrial translocation of Bax and downstream mitochondrial-dependent apoptosis in the cochlea [57, 85]. STAT1 activity also links the apoptotic and inflammatory pathways involved in cisplatin ototoxicity [64], which will be discussed later. Third, multiple factors, including cisplatin itself, ROS, lipid peroxidation, calcium influx, and CDKs, can act directly on mitochondria, causing an increase in its permeabilization and deterioration of its function. Cisplatin can also form mitochondrial DNA (mtDNA) adducts [98-101], which is likely the main factor in cisplatin-induced ototoxicity [15]. The platination of mtDNA leads to mtROS production and accumulation, and in turn, the damage of mtDNA, proteins, and lipids within the mitochondrial membrane. Cyclin A is another critical player in mtROS production, which can be upregulated by cisplatin and activates CDK2 kinase, consequently facilitating mtROS production [102]. Fourth, cytochrome C release and the activation of caspase-9 and caspase-3 mark the final stage of mitochondrial-dependent apoptosis, which is also observed in the cochlea [72] (**Figure 2**). As an essential component of the mitochondrial electron transport chain, cytochrome C is one of the apoptotic protease-activating factors, which activate caspase 9. Caspase 9 activation subsequently activates caspase 3, causing the fragmentation of chromosomal DNA through the cleavage of its substrates [103].

Mitochondrial-independent apoptosis pathways in cisplatin ototoxicity

Apoptosis can also be induced in the cytoplasm through ROS overload, ER stress [104], inflammation, and lipid peroxidation [79].

ER stress: ER stress results from oxidative damage, intracellular calcium imbalance, and protein damage [83]. It is involved in cisplatin ototoxicity through caspase-12 activation, located on the ER plasma membrane, and auto-cleaves in response to ER stress. Caspase-12 then activates caspase-9, which activates caspase-3, leading to apoptosis in a mitochondrial-independent manner [97, 98, 105]. This ER-specific apoptosis pathway is also closely linked to the mitochondrial-dependent path due to the simultaneous activation of C/EBP homologous protein (CHOP) [104]. CHOP plays an important role in ER stress-induced apopto-

sis while regulating Bcl2 family expression. Decreased Bcl2 enables increased Bax activity, leading to the release of apoptotic active substances from mitochondria to the cytoplasm [106], i.e., mitochondrial-dependent apoptosis [79, 104] (**Figure 2**).

Inflammation: Inflammation is the body's defense mechanism in response to harmful stimuli, such as damaged cells, which also plays a vital role in inducing apoptosis. Inflammatory signaling pathways, most commonly the nuclear-factor kappa B (NF- κ B), mitogen-activated protein kinases (MAPKs), and JAK-STAT pathways, are also involved in cisplatin ototoxicity. NF- κ B is a transcription factor mediating inflammatory responses by regulating the expression of various pro-inflammatory genes, such as tumor necrosis factor- α (TNF- α). In cisplatin ototoxicity, NF- κ B can be activated directly by ROS [107] or toll-like receptor 4 (TLR4) [108], leading to its translocation to the nucleus and production of inflammatory cytokines [80, 92]. TLR4 is a transmembrane protein that plays a fundamental role in pathogen recognition and activation of innate immunity. TLR4 can also be activated by cisplatin, which triggers the activation of proinflammatory cytokines such as interleukin-6 (IL-6), interleukin-8 (IL-8), TNF- α , and NF- κ B [108]. Translocation of NF- κ B to the nucleus, mediated by TNF- α and IL-6, further induces the *de novo* synthesis of TNF- α , IL-6, IL-1 β , and inducible nitric oxide synthase (iNOS), and activates caspase-3 in a mitochondrial-independent manner [78, 79]. Nevertheless, NF- κ B translocation can occur much earlier (1-2 hours) than maximal ROS generation (1 day), suggesting an early involvement of the inflammatory pathways or even an upstream event of ROS formation [15, 59, 79, 109, 110]. NF- κ B [111] (and also STAT1 [85]) induces an increased expression of iNOS, which produces nitric oxide (NO) (for reviews, see [78, 79]). NO reacts with O₂⁻ to form peroxynitrite (ONOO⁻), a highly reactive oxidizing molecule that damages proteins [112]. Peroxynitrite induces protein peroxidation via nitration of tyrosine residues (nitrotyrosine) [112, 113], which alters protein configuration and function (for reviews, see [13, 15, 72]). In addition, expression of TNF- α also further activate NF- κ B [80, 92] as well as the extrinsic apoptosis pathway by binding to TNF receptors (TNFR). It leads to caspase-8 activation, which

in turn activates caspase-3, leading to mitochondrial-independent apoptosis (**Figure 2**) [71, 93], which is also known as the death receptor pathway [42] or extrinsic apoptosis [94].

Other than the NF- κ B pathways, early phosphorylation of two well-characterized MAPK families, ERKs and the c-Jun N-terminal kinases (JNKs), is activated by cisplatin treatment in two cochlear-derived cell lines - House Ear Institute-Organ of Corti 1 (HEI-OC1) cells [109] and OC-k3 cells [77] (**Figure 1**). Early activation of JNK may play a minor role in cisplatin-induced ototoxicity [109] or may assist in DNA repair in response to cisplatin-DNA adducts [114]. The activation of MAPK/ERK also facilitates secretion of the pre-existing TNF- α , IL-1 β , and IL-6 [77] (**Figure 2**), which in turn activates the translocation of NF- κ B to the nucleus [109]. Nevertheless, most of the studies mentioned above are performed on cell cultures, although the role of the MAPK family in cisplatin toxicity is well documented [115]. It is also worth noting that most of the research on the molecular mechanism of cisplatin ototoxicity included in this review has been conducted using *in vitro* cell cultures or *ex vivo* cochlear explants. A recent study has highlighted that the molecular pathways implicated in cisplatin ototoxicity may differ between the *in vitro/ex vivo* and the *in vivo* settings [116].

JAK-STAT pathway activation allows the transfer of signals from the receptors to the nucleus. It thus involves a repertoire of processes, such as apoptosis and tissue repair, through a cytokine-membrane receptor-JAK-STAT cascade [117]. In CIHL, the JAK-STAT pathway regulates cell death and inflammatory responses by activating the expression of inflammatory cytokines such as cyclooxygenase-2 (Cox-2) and TNF- α . Suppression of the JAK-STAT pathway with an adenosine A1 receptor (A1AR) agonist decreases cisplatin-induced apoptosis of OHCs [118]. Among STAT proteins (STAT1-6), STAT1 is likely the most relevant to CIHL because it is known to directly induce apoptosis and p53-mediated apoptosis [64, 85]. In rat models, STAT1 signaling is linked with ROS, causing cisplatin-induced cochlear cell apoptosis [85]. Pre-treatment with STAT1 siRNA (48 hours before) [85] or oral uptake of an inhibitor of STAT1 signaling (45 minutes before) [64] both

protect against CIHL. STAT3 and STAT6 are also important players in CIHL [119]. STAT6 works through inflammatory cytokines IL-4 and IL-13, and knockout of STAT6 protects against CIHL [119].

Lipid peroxidation: Downstream effects of $O_2^{\cdot-}$ can also cause lipid peroxidation. Catalyzed by SOD, $O_2^{\cdot-}$ is converted to hydrogen peroxide, which is catalyzed by iron to form hydroxyl free radicals [15]. These highly reactive ROS react with polyunsaturated fatty acids in cellular membranes, producing the highly toxic aldehydes, 4-hydroxynonenal (4-HNE) and malondialdehyde (MDA) [72]. Antioxidant enzyme depletion has been linked to increased levels of MDA, indicating high lipid peroxidation [80]. This inverse relationship between glutathione and MDA activity (low glutathione levels and high MDA levels) has been shown *in vivo* using cisplatin-treated rats [86]. Lipid peroxidation, especially the production of 4-HNE, induces calcium influx into the cells [15, 88]. ROS further enhances calcium influx. An early study found that $O_2^{\cdot-}$, but not H_2O_2 , increases the intracellular calcium concentration via transmembrane influx in OHCs of guinea pig cochleae. Researchers suggested that $O_2^{\cdot-}$ stimulates voltage-sensitive calcium channels. However, they did not rule out the possibility of increased calcium permeability caused by lipid peroxidation [120]. According to a review article, ROS can open the ER calcium channel, ryanodine receptor, L-type, T-type, and the TRPV1 plasma membrane calcium channels [79]. A study by Yoshida et al. found that intracellular NO and H_2O_2 open TRPV1 channels and facilitate calcium influx [121]. In an earlier study, inhibition of T-type calcium channels did not inhibit ROS generation. Still, it significantly inhibited lipid peroxidation, mitochondrial membrane permeabilization, and cytochrome c release in HEI-OC1 cells and rat organ of Corti explants treated with cisplatin [96]. The findings suggest that calcium influx via T-type calcium channels occurs downstream of ROS and plays a major role in intrinsic apoptosis of cochlear cells under cisplatin-induced stress conditions. Nevertheless, increased cytosolic calcium causes mitochondrial membrane permeabilization and loss of membrane potential, initiating the intrinsic mitochondrial-dependent apoptosis pathway [122].

Other cell death pathways

Necroptosis: In addition to activation of the extrinsic apoptosis pathways, TNF- α induces cell death via necroptosis pathway activation in cisplatin-induced ototoxicity [116, 123] (for reviews, see [59, 124]). Studies indicate a dose-dependent effect on cisplatin-induced cell death using HEI-OC1 cells [103] and OC-k3 cells [77]. Specifically, apoptosis occurs at lower doses, while necroptosis occurs at higher doses. As opposed to the organized breakdown of cells during apoptotic cell death, necroptosis results from cellular and organelle membrane permeation, releasing intracellular substances and exacerbating inflammation [122]. Activation of caspase-8 inactivates receptor-interacting serine/threonine-protein kinase 1 (RIPK1) and RIPK3, which induce activation of the extrinsic mitochondrial-independent apoptosis pathway as described above. Inactivation of caspase-8 leads to RIPK3 activation, shifting death receptor-mediated cell death from apoptotic to necroptotic pathways [124]. In necroptosis, TNF- α binds its receptor 1 (TNFR1) and, without caspase-8, leads to the formation of the RIPK1/RIPK3 complex, also known as necrosome [123, 124]. Downstream effects of necrosomes involve the activation of Mixed Lineage Kinase Domain-Like Pseudokinase (MLKL), which induces calcium influx via Transient Receptor Potential Cation channel, subfamily M, member 7 (TRPM7) channels. It creates pores in the plasma membrane, leading to leakage of substances, cell lysis, and cell death [125].

Autophagy: As a protective mechanism of cells, autophagy removes specific cellular structures or components, such as damaged organelles. Under stress conditions such as cisplatin treatment, autophagy can also lead to cell death [126]. The interest in the involvement of autophagy in CIHL has increased rapidly in recent years. Increased expression of three autophagic mediators has been observed in CIHL, including Beclin-1 (the initial autophagy promoter), microtubule-associated protein light chain 3 II (LC3-II), and mitochondrial-bound [127] nucleotide-binding domain and leucine-rich-repeat-containing family member X1 (NL-RX1) [59]. Autophagy signaling involves specific factors and interconnections with other cell death pathways. First, intracellular organelle or

protein damage leads to the activation of AMP-activated protein kinase, which inhibits the mammalian target of rapamycin (mTOR), leading to autophagy [128] and to the activation of phosphoinositide 3-kinase (PI3K), inducing Beclin-1 formation of the phagosome [129]. Second, beclin-1 activates the conversion of unlipidated LC3-I to lipidated LC3-II, which is required for complete autophagosome formation and autophagosome-lysosome fusion [126, 129]. Third, overexpression of NLRX1 correlates with the accumulation of autophagosomes and acceleration of autophagic cell death [83, 126]. In addition, NLRX1 overexpression also accelerates mitochondrial-dependent apoptosis in cisplatin treatment, evidenced by increased Bax, caspase-3, and ROS levels [127]. The interactions between autophagy and apoptosis pathways are also shown in mitochondrial autophagy or mitophagy. Activation of mitophagy can negatively regulate cisplatin-induced apoptosis in hair cells and SGNs [130] and vice versa [131], suggesting a protective role of mitophagy in CIHL. Nevertheless, questions still exist about whether autophagy plays a protective role or induces cell death in CIHL, and the arguments are far from settled (for reviews, see [59, 132, 133]).

Ferroptosis: Ferroptosis is another form of non-apoptotic programmed cell death, which requires irons generated by processes such as lipid peroxidation and autophagy in response to stress [134, 135]. Hallmarks of ferroptosis, such as lipid peroxidation and impaired antioxidant capacity, have been observed in cochleae in mice after cisplatin treatment, suggesting its potential roles in CIHL [135]. A recent study showed that transferrin 1, a marker of ferroptosis, is increased in OHCs but not IHCs, SV, or supporting cells after cisplatin treatment. RNA sequencing completed in the same study showed that the expression of the ferroptosis-related gene is upregulated in CIHL [136]. Suppressing ferroptosis using ferrostatin-1 reduces CIHL by protecting cochlear hair cells [135, 137, 138] while facilitating ferroptosis by blocking lipid repair function leads to an exacerbated breakdown of mitochondrial membrane potential in cultured HEI-OC1 cells. Ferrostatin-1 also protects hearing against CIHL in mice and rescues OHCs in a knockout mouse model lacking a key regulator for ferroptosis [136]. However, ferroptosis is highly dependent

on mitochondrial function and interconnected with other signaling pathways, including lipid peroxidation and autophagy. Further studies are needed to address the potential interference of ferroptosis with the antitumor effects in chemotherapy involving cisplatin. More discussion regarding the roles of ferroptosis in CIHL and the underlying mechanism can be found in some recent studies [139, 140] and reviews [89].

Hearing protection in chemotherapy

Current studies on hearing protection in chemotherapy are mainly focused on cisplatin ototoxicity. Among the four ototoxic chemotherapeutic drugs listed in **Table 1**, vinblastine and dasatinib have rarely been studied in hearing protection, while the studies on carboplatin are almost all associated with cisplatin [141, 142]. Since the damage to the cochlea is largely irreversible and regeneration has not been successful yet, intervention before and during chemotherapy to protect hearing is critical. However, preventing cisplatin ototoxicity using drug therapies is facing severe challenges, including interference with the therapeutic effects of cisplatin, bioavailability, and side effects. Concerns regarding interference with the antitumor effects of cisplatin lie in the fact that interventions with antioxidants that reduce cisplatin ototoxicity may also affect the outcomes of the cancer therapy by deactivating cisplatin [78, 143, 144] and by protecting tumor cells [145] (for review, see [78, 146]). Challenges to bioavailability include the permeability of otoprotective drugs through BLB [50, 51] and cell membranes. Side effects are manifested by the toxicity of hearing protective drugs to other tissues. An example is amifostine [147], for which high drug doses are required for treatment because of the drug's impermeability of the BLB *in vivo*.

Drug delivery strategies

Local drug administration: The challenges mentioned above can be addressed through trans-tympanic delivery [148], which has proven to be successful for some candidate drugs for hearing protection. Local drug administration through trans-tympanic injection has been developed in early studies to treat inner ear disorders such as Ménière's disease and sudden

hearing loss using steroids (for review, see [149]). It is also adopted to treat cisplatin ototoxicity with antioxidants to avoid interfering with the antitumor effects [143, 148, 150, 151]. Depending on the site, two types of local drug administration are usually used: intratympanic (middle ear and round window) and intra-cochlear/labyrinthine (perilymph). Intra-tympanic injection, sometimes combined with a small tube through tympanic annulus for multiple doses, allows localized and high-dose treatment, as shown in some clinical trials treating cisplatin ototoxicity [152]. Intra-cochlear/labyrinthine delivery further renders the benefits of passing through the BLB, sustained treatment, and (semi-) dose control when combined with chronic implantation of an osmotic pump [153]. Local administration is shown to be effective for many drugs without interfering with the antitumor effect of cisplatin [154-156]. Concerns for local drug administration include potential pain during myringotomy and damage to the tympanic membrane, middle ear structures, and round window. Intra-cochlear/labyrinthine injection may even cause permanent damage to the inner ear, which may prevent its potential application in clinics. A more detailed introduction of the local delivery techniques and their application in cisplatin ototoxicity [79] can be found in research papers and reviews [157-161].

Other drug administration: Some other strategies have been developed to increase the efficiency of hearing protection and reduce the side effects and risk of counteracting the antitumor effect of cisplatin. Hydrogel has been used for intra-tympanic drug delivery to improve drug sustainability on the round window by reducing drainage through the eustachian tube [162-164]. However, subsequent conductive hearing loss posed by the residue on the round window and middle ear ossicles might be a concern. Nanoparticles, with a diameter of less than 1 μm , are another strategy to increase the efficacy of local drug delivery for treating cisplatin ototoxicity [165]. A few groups have studied the influence of particle size and various transportation vehicles [166]. The paper by Yu et al. details various nanoparticles and hydrogel preparations for improved drug delivery to the inner ear and cochlea manipulations [79]. As stated in the paper, inadequate cellular uptake of nanoparticles limits its use. However, it can

be reduced with various endogenous (pH, redox, and enzymes) and exogenous (heat, ultrasound, light, and magnetic field) measures. Delayed or staggered drug delivery is also used to reduce the possibility of interference with the antitumor effect of cisplatin. Nevertheless, showing accurate drug concentration in the cochlea will help to determine the mechanism and effective doses of the drugs. In this regard, proof of the existence of the drugs in the cochlea by measuring the concentration from the extraction of cochlear fluids is a gold standard for candidates claiming hearing protection [167-169].

Antioxidants as traditional candidates for hearing protection against cisplatin ototoxicity

Since ROS plays a central role in cisplatin ototoxicity, various exogenous antioxidants that work as free radical scavengers have been tested for hearing protection in chemotherapy in early studies [51, 170-172]. A common feature of antioxidants is the presence of a thiol, which can act as a substrate for redox reactions. Some antioxidants have been tested in clinical practice, such as N-acetyl-L-cysteine (NAC) [172], D-methionine [173, 174], STS [155, 163], and amifostine [147, 175] (for reviews, see [13, 72, 78, 142, 147, 176]). A list of antioxidants tested for their hearing protective effects against cisplatin ototoxicity is presented in **Table 3**. Some well-known and promising candidates and their roles are shown in **Figure 3** and described below.

STS: STS has been traditionally used to treat metal poisoning. It can form strong complexes with metal ions, including platinum, thus deactivating cisplatin. Its antioxidant property comes from a reduced sulfur residue. The hearing protective effect of STS has been proven in both animal studies [172, 177] and clinical trials [178, 179]. Local delivery through intra-cochlear injection [148, 152] or delayed administration [177] is required to treat cisplatin ototoxicity. Although the effect of local delivery is controversial [148, 152], delayed administration of STS by 4-8 hours is proven to be protective to hearing [178, 179], while it does not interfere with the therapeutic effect of cisplatin [180]. In these clinical studies, the incidence of cisplatin ototoxicity decreased by about half, and no significant difference in overall or event-

Hearing loss during chemotherapy

Table 3. Antioxidant drug candidates for protection against CIHL

Drug	Main mechanism of oto-protection	Experiment model	Drug dose	Route	Outcome
STS (sodium thiosulfate) Branded as Nithiodote & PedMark	By forming complexes with platinum & antioxidant activity (PMID: 25994788)	Phase III clinical trial on hepatoblastoma in children (n = 109) (PMID: 29924955)	Cisplatin (80 mg/m ²) per dose over 6 doses	IV	Reduction in hearing loss grade 1-4 by 48%.
			STS (20 g/m ²) 6 hours after each cisplatin treatment	IV	
		Phase III Clinical trial on children (n = 104) with cancers (PMID: 27914822)	Cisplatin (totaling 200 mg/m ²) STS (16 g/m ²) 6 hours after each cisplatin treatment	IV IV	
NAC (N-acetyl-L-cysteine) Branded as Acetadote, Fluiimucil & Mucomyst	Antioxidant with a thiol donating hydrogen (PMID: 29429900)	Rat cochlea cell culture treated with NAC (PMID: 35346799)	Cisplatin (50 uM) treatment to cell cultures in the background of NAC at 37 °C for 48 hours	Cell culture	Hair cell loss prevented, seen through Ab staining.
		Rat treated with NAC (PMID: 15219317)	Cisplatin (6 mg/kg) NAC (400 mg/kg) 15 minutes before cisplatin	IA IV	Hearing protected at 4 kHz (~10 dB), 8 kHz (~20 dB), 12 kHz (~22 dB), 16 kHz (~18 dB) by change in threshold.
		Phase I clinical trial on children (n = 52) with cancers (PMID: 37134194)	Cisplatin (totaling 200 mg/m ²) NAC (6 g) 4 hours after each cisplatin treatment	IV IV	Reduction in the risk of SIOP ≥ 2 hearing loss post-chemotherapy (OR = 0.13, CI 0.021-0.847, P = 0.033).
		Double blind clinical trial on cancer patients (n = 114) (PMID:29993216)	Cisplatin (not stated) 0.4-0.8 ml NAC (10%)	Not stated Intratympanic	No significant changes in auditory thresholds.
D-methionine (D-met)	D-methionine can donate a cysteine which can act as an antioxidant (PMID: 16366723)	Mice treated with D-methionine (PMID: 8951454)	Cisplatin (16 mg/kg) D-met (300 mg/kg)	IP IP	Hearing protected at 1, 4, 8, 14 kHz.
		Chinchilla treated with D-met (PMID: 12087338)	Cisplatin (125 µg) D-met (4 µg) 30 minutes before cisplatin treatment	Intracochlear Intracochlear	Hearing protected at 8, 16 kHz.
		Rats with MTLn3 breast cancer cells (PMID: 11405249)	Cisplatin (5 mg/kg) per dose for 3 doses with 72 hours interval L-met (300 mg/kg) 30 minutes before each cisplatin injection	IP IP	Hearing protected at 1, 2, 4, 8, 16, 18 kHz. OHC protected.
		Phase II clinical trial in cancer patients (n = 27) (PMID: 34622731)	Cisplatin totaling (264 mg/m ²) D-met (100 mg/kg) oral dose 1 hour before cisplatin injection	IV Oral	Hearing protected in the left ear only at 11.2 kHz by a mean difference of 22.97 dB.

Hearing loss during chemotherapy

Amifostine (WR-2771) Branded as Ethiol	Antioxidant activity through WR-1065 (PMID: 11201306)	Scid mice model of human ovarian cancer with implanted tumor cells (PMID: 9389935)	Paclitaxel (27 mg/kg) over 5 doses	IP	Amifostine improved animal survival. In cell cultures, amifostine protected normal cells from paclitaxel, while cytotoxicity was increased in malignant cells.	
			Amifostine (200 mg/kg) over 5 doses	IP		
		Hamsters treated with amifostine (PMID: 15185124)	Cisplatin (15 mg/kg) over 5 doses	IP		Protected hearing at 8 kHz (~15 db), 16 kHz (~20 db), 20 kHz (~18 db) by ABR threshold shift.
			Amifostine (40 mg/kg) over 5 doses 30 minutes before cisplatin injection	IP		
		Non-randomized clinical study with cancer patients (n = 62) (PMID: 18669462)	Cisplatin (300 mg/m ²) over 4 doses	IV		Amifostine caused ~22% reduction in the probability of requiring hearing aid. Amifostine related adverse reaction in 19% patients.
			Amifostine (600 mg/m ²) 3 hours before and immediately before cisplatin injection	IV		
		A randomized clinical study (n = 25) investigating medulloblastoma in children (PMID: 15999362)	Cisplatin (40 mg/m ²) per day for 5 days	IV		Amifostine did not offer otoprotection against cisplatin combined with etoposide and bleomycin.
			Amifostine (825 mg/m ²) per day for 5 days 30 minutes before cisplatin	IV		
			Etoposide (100 mg/m ²) per day for 5 days	IV		
			Bleomycin (15 IU/m ²) per day for 5 days	IV		
Tauroursodeoxycholic acid (TUDCA)	HO1 and SOD2 mediated antioxidant activity (PMID: 32061715)	Rats treated with TUDCA (PMID: 32061715)	Cisplatin (5 mg/kg) per day for 3 days	IP	Protected hearing at 4, 8, 16, 32, 40 kHz.	
			TUDCA (100 mg/mL) 1 hour before cisplatin treatment	IP		
TUDCA	HO1 and SOD2 mediated antioxidant activity of TUDCA (PMID: 32061715)	Rats treated with TUDCA (PMID: 33631298)	Cisplatin (4.6 mg/kg) per day for 3 days	IP	Protected hearing at 8, 16, 24, 32, 40 kHz.	
			TUDCA (500 mg/kg) 1 day before cisplatin treatment	Subcutaneous		
Ebselen	Antioxidant activity acting as a GPx mimic	Mice treated with ebselen (PMID: 19286452)	Cisplatin (16 mg/kg)	IP	Protected hearing at 4, 8, 16, 32 kHz.	
			Ebselen (16 mg/kg)	IP		
		Rat treated with ebselen (PMID: 21804453)	Cisplatin (14 mg/kg)	IV	Did not protect hearing.	
			Ebselen (12 mg/kg) 1 hour before cisplatin treatment	IP		
		Rat treated with ebselen and allopurinol (PMID: 15721563)	Cisplatin (16 mg/kg)	IP	IP ebselen + allopurinol protected hearing at 8, 16, 24 kHz.	
Ebselen (8 mg/kg) 1 hour before cisplatin treatment	IP/oral gavage					
	Allopurinol (8 mg/kg) 1 hour before cisplatin treatment	IP/oral gavage				

Hearing loss during chemotherapy




Figure 3. Schematic image showing key targeting cellular factors and signaling steps for preventative strategies against cisplatin ototoxicity. Cisplatin induces DNA damage and ROS generation. Multiple processes and cytokines also contribute to ROS generation along the apoptosis and other (such as inflammation) signaling pathways. Proposed agents (in purple) target different sites in these processes. Refer to the text for details and abbreviations.

free survival was observed between the treated group and the cisplatin-alone group [178, 179]. In 2022, the FDA approved STS for protective treatment against cisplatin ototoxicity in pediatric patients with localized, non-metastatic solid tumors. It is the first and the only approved drug to treat cisplatin ototoxicity in clinics.

NAC: NAC is a precursor to GSH, a ubiquitous thiol-containing antioxidant and a widely used cytoprotectant with a promising hearing protective effect [181-184]. As a weak free radical scavenger, NAC may protect against cisplatin ototoxicity by induction of endogenous GSH and blocking apoptosis through the caspase signaling pathway [181, 185]. Pre-treatment of NAC (400 mg/kg) improves cisplatin-induced hearing threshold shift in rats [186], although the results are slightly inconsistent with an organo-culture study [185]. Nevertheless, since NAC also protects against cisplatin cytotoxicity [181], a local or staggered administration is required for its otoprotection [186, 187]. In clinical studies using intra-tympanic injection, NAC failed to prevent hearing loss during cisplatin treatment, although it was suggested for

some patients [154, 184]. In a recent phase I clinical study with children and adolescents newly diagnosed with nonmetastatic tumors delayed NAC application through intravenous (i.v.) injection (4 hours after cisplatin injection, peak concentration of NAC up to 450 mg/kg) was effective for preventing CIHL without adverse events [188]. These studies indicate that NAC might be another promising drug for treating cisplatin ototoxicity in clinics.

D-methionine: D-methionine is the enantiomeric counterpart of L-methionine, a naturally occurring L-alpha-amino acid. The antioxidant property of methionine is due to methionine being oxidized into methionine sulfoxide, which inhibits ROS. It also deactivates cisplatin by forming an inactive complex with it [78]. D-methionine is one of the earliest antioxidants tested in the protection against CIHL [189] and has proven to be effective both in animal studies [152, 189, 190] and in humans [174]. Since it is known that D-methionine interferes with the therapeutic effects of cisplatin, local, delayed, or staggered delivery should be considered in clinical trials. The in-human study, completed in India in 2022, used oral

Hearing loss during chemotherapy

uptake of D-methionine (100 mg/kg) one hour before cisplatin treatment. The study did not show adverse events significantly different from the placebo group [174]. However, the study analyzed data from only 27 of the 50 enrolled participants. Furthermore, the study did not gather data on whether D-methionine affected cisplatin cytotoxicity in tumors. Overall, more careful examinations of the research and further larger scale clinical trials might be necessary to assess the influence of D-methionine on the antitumor efficacy of cisplatin.

Amifostine: Amifostine is used in chemotherapy and radiotherapy as a cytoprotective adjuvant for reducing renal toxicity [191]. Alkaline phosphatase can dephosphorylate it into an active free thiol metabolite that functions as an antioxidant [192]. Its cytoprotective action is believed to be achieved through scavenging free radicals, stabilizing DNA, and upregulating p53 [176], thereby selectively protecting normal tissues due to their high alkaline phosphatase levels [193]. Moreover, amifostine does not affect the antitumor activity of platinum drugs [192]. In animal research, multiple high doses of amifostine, given before cisplatin, protect hearing in hamsters (40 mg/kg/day) [194] and guinea pigs (100 mg/kg/day) [195]. However, lower doses (18 mg/kg/day) fail to achieve this effect [196]. Clinical studies on amifostine's ability to protect against cisplatin ototoxicity have yielded mixed results. For instance, one study with 97 pediatric patients with medulloblastoma indicates a reduced need for hearing aids following amifostine treatment (600 mg/m²) [147]. In contrast, another study with 11 patients shows no protective effect (1000 mg/m²) [197]. A follow-up clinical trial with 379 patients reveals that amifostine protects from cisplatin ototoxicity only in average-risk but not in high-risk medulloblastoma patients [198]. In another study with 25 cases of pediatric germ cell tumor, amifostine treatment (825 mg/m²) does not confer otoprotection compared to historical controls [175]. These varying outcomes may be attributed to differences in treatment regimens, sample sizes, age groups, and cancer types across studies. As a note, two retrospective evaluations of the efficacy of amifostine in preventing cisplatin ototoxicity have been either inconclusive [199] or insignificant [200], possibly due to similar complexities. While high doses of

amifostine are crucial for the treatment, they are associated with adverse events, including neurotoxicity (in animal studies) [194], hypocalcemia, hypotension, and nausea and vomiting (in clinical settings) [147]. Complex administration protocols, careful medical attention, and sustained monitoring of vital signs are required to ensure patient safety during amifostine treatment [201]. A more detailed review can be found in [78].

Coenzyme Q10: Coenzyme Q10 is a crucial enzyme for electron transport in the mitochondrial respiratory chain. It also works as an anti-inflammatory and antioxidant agent in dietary supplements [202]. Co-application of coenzyme Q10 and multivitamins was tested for protection against CO in both an animal model [203] and a pilot case-control clinical trial [204]. In the animal study, Q10 terclatrate (500 mg/kg) and vitamin supplements (vitamin E and B12) given before cisplatin (4.6 mg/kg/day for three days) treatment protected against CIHL in rats. In the pilot study on cancer patients treated with a coenzyme Q10 plus dietary multivitamin, the incidence of hearing disorders and tinnitus induced by cisplatin was reduced significantly. A significant difference in threshold shift only occurred at 8 kHz [204]. However, a fully powered clinical study has not been completed, even six years after the pre-clinical animal study.

Candidates/approaches targeting specific pathways/factors

Candidates targeting more specific cell signaling pathways in cisplatin ototoxicity are also often proposed and tested for their hearing protective effects. An early attempt is dexamethasone, a glucocorticosteroid that can downregulate proinflammatory cytokines, inhibit apoptosis, and upregulate antioxidant enzymes [78]. While animal studies yielded promising results [150, 151, 162], dexamethasone failed in most clinical trials [154, 156], although attempts were made using nanoparticles [205] or poloxamer hydrogels [162] to increase local drug concentration in the cochlea. A detailed review regarding the protective effect of dexamethasone against cisplatin ototoxicity can be found in [78], with no further discussion in this review. As the understanding of cisplatin ototoxicity signaling pathways grows, more spe-

cific targets on factors/cytokines in metabolic [203, 204], inflammatory [206, 207], or apoptosis signaling pathways [102, 208, 209] have been identified with promising results. Some novel candidates that have been tested are listed in **Table 4**, and their potential mechanism is shown in **Figure 3**. These candidates include neurotrophins [210], hormones [154], molecules involved in endogenous metabolism [204], modulators of cell signaling pathways [102], etc.

Epigenetic modulators: The significance of epigenetic modulation in inner ear development, damage, and protection was noticed over a decade ago [211-213], while it has garnered more attention only in recent years [214-216]. Histone deacetylases (HDACs) and their inhibitors (HDACis) regulate transcription and control cell cycle through modulation of histone acetylation. HDACis, such as sodium butyrate [212] and sulforaphane [215], are protective against cisplatin ototoxicity in animal models (guinea pigs and rats). The mechanism might be related to the pro-survival pathway activation, as shown in the study on HDAC inhibition against kanamycin and furosemide-induced hearing loss [214]. In addition, both sodium butyrate [217] and sulforaphane [218] can induce the expression of an antioxidant-responsive gene, Nrf2, which may also account for their hearing protective effects [219]. HDACs have antitumor activity and are well tolerated. They are ideal candidates for treating cisplatin ototoxicity. However, their mechanisms of action are not fully understood, and explanations rely on indirect evidence only. More studies regarding their specificity are needed before testing them in clinical trials.

Other promising epigenetic modifications have been reported recently. They are still conducted in cell cultures and animals. A recent study showed that a DNA methyltransferase inhibitor, RG108, alleviates CIHL in a mouse model [216]. Inhibition of DNA methylation by RG108 upregulates BCL-2 and downregulates mitochondrial-dependent apoptosis pathway factors BAX and BCL2 Associated Agonist of Cell Death (BAD) in HEI-OC1 cell cultures. Another histone methyltransferase inhibitor, BIX01294, has been shown to protect against CIHL in a mouse model established through the combination of i.p. furosemide and subcutaneous cis-

platin delivery [220]. The mechanism, verified primarily on cell cultures, is associated with activating the autophagy pathway through the Forkhead box G1 gene, a critical regulator for morphogenesis of the mammalian inner ear during development. Nevertheless, further studies to verify these results and mechanisms are necessary, and their interference with the antitumor effects of cisplatin and other adverse effects are still unclear.

Statins: Statins are widely used as anti-inflammatory agents to control lipid peroxidation. They are proposed to protect hearing against various insults (noise [221], drug [222], and age [223]). For a review, see [224]. Statins are also used in an ongoing clinical trial against sudden hearing loss (ClinicalTrials.gov ID: NCT04826237). Two recent studies showed that statins protect against CIHL in animals (lovastatin [206]) and humans (atorvastatin [207]). The non-randomized, retrospective clinical study executed by Cunningham's group showed that atorvastatin (with unknown doses) decreased the incidence of CIHL by 53% without affecting the three-year survival rates in head and neck cancer patients [207]. The same group will perform a randomized phase III clinical study (NCT04915183) on atorvastatin (20 mg) against CIHL. In this clinical trial, 186 patients with squamous cell carcinoma of the head and neck will be involved. The study aims to assess hearing using audiograms before and 2-4 months after cisplatin treatment and compare outcomes between atorvastatin users and controls. CTCAE criteria will be used for data analysis, and reduced bias will be expected. Although it might not be a major concern, sequelae of statins include muscle pain, liver damage, increased blood sugar, and memory loss.

Protein kinase inhibitors: Protein kinase phosphorylation cascades regulate multiple signal transduction pathways, including inflammation, apoptosis, and proliferation. Cytokine inhibitors can reduce cytokine synthesis and concentration and interfere with the interaction between cytokines and their receptors. Certain kinase inhibitors have been studied for their potential to protect against CIHL. Flunarizine, known initially as a calcium channel blocker, could reduce cisplatin-induced inflammatory cytokines (TNF- α , IL-1 β , NF- κ B seen through IHC) in

Hearing loss during chemotherapy

Table 4. Non-antioxidant drug candidates for protection against CIHL

Drug	Main mechanism of oto-protection	Experiment model	Drug dose	Route	Outcome
Dexa-methasone Branded as DexPak & Decadron	Inhibiting inflammatory cytokines (PMID: 9486420)	Mice treated with dexamethasone (PMC2720789)	Cisplatin (14 mg/kg) treatment after the first dexamethasone injection	IP	Protected hearing at 16 kHz.
		Guinea pigs treated with dexamethasone (PMID: 21521888)	Dexamethasone (24 mg/ml) per day over 5 days	Intratympanic	Not protective.
			Cisplatin (12 mg/kg)	IP	
			Dexamethasone (6 mg) treatment on the day before and on the day of cisplatin treatment	Intratympanic	
Coenzyme Q10 Branded as ubiquinone & CoQ10	Anti-inflammatory dietary supplements (PMID: 35326965)	Clinical trial on cancer patients (n = 26) on dexamethasone (PMID: 24618499)	Cisplatin (total 400 mg) Dexamethasone (~8.5 mg)	IP Intratympanic	Protected hearing at 4-8 kHz.
		Rats treated with Coenzyme Q10 and vitamins (PMID: 27632426)	Cisplatin (4.6 mg/kg) per day for 3 days	IP	Protected hearing at 2, 4, 8, 16, 32 kHz.
			Q10 terclatrate (500 mg/kg)	Oral	
			Acuval 400 (100 mg/kg)	Oral	
Statins Branded as Lipitor & Mevacor	Inhibiting inflammatory cytokines (PMID: 35125240).	Clinical trial on cancer patients (n = 26) being treated with Coenzyme Q10 and vitamins (PMID: 28239674)	Cisplatin (100 mg/m ²) every 21 days Acuval Audio (1.8 g) per day starting 7 days before the first cisplatin treatment	IV Oral	Protected hearing at 8 kHz.
		Mice treated with lovastatin (PMID: 32062294)	Cisplatin (3 mg/kg) per day for 4 days, followed by 10 days of recovery. 3 cycles (total 36 mg/kg)	IP	Protected hearing at 20, 40 kHz.
			Lovastatin (40 mg/kg) daily from 3 days before cisplatin treatment	Oral gavage	
			Clinical trial on head and neck cancer patients (n = 277) on atorvastatin (PMID: 33393488)	Cisplatin (~80 mg/m ²) every 3 weeks. Median cumulative cisplatin was 200 mg/m ² . Atorvastatin (10~80 mg)	
HDACi Branded as Buphenyl & Avmacol	Epigenetically downregulating genes linked to apoptosis (PMID: 23558232)	Guinea pigs treated with sodium butyrate (PMID: 16467722)	Cisplatin (14 mg/kg)	IP	Protected hearing at 3.5~20 kHz, statistical significance not shown.
		Rats treated with sulforaphane (PMID: 34344210)	Sodium butyrate (1.2 mg/kg) per day for 7 days before and 5 days after cisplatin treatment	IP	
			Cisplatin (7 mg/kg) twice a day for 7 days	IP	Protected hearing at 4, 8, 16, 24, 32 kHz. OHC partially protected.
			SFN (30 mg/kg) per day for 7 days	IP	
STAT inhibitors Branded as Nifuroxazide etc	Anti-inflammatory and anti-apoptosis	Mice treated with RG108 (PMID: 35530135)	Cisplatin (30 umol/L) RG108 (100 umol/L) 2 hours before cisplatin treatment	IP IP	Protected hearing at 8, 16, 24, 32 kHz. OHC protected seen through IHC.
		Rats treated with STAT1 siRNA (PMID: 21776018)	Cisplatin (11 mg/kg)	IP	Protected hearing at 8, 16, 32 kHz. OHC protected seen through electron microscopy.
			STAT1 siRNA (0.5 µg) 48 hours before cisplatin treatment	Intratympanic	
			Rats treated with STAT1 inhibitor EGCG (PMID: 28703809)	Cisplatin (11 mg/kg) EGCG (100 mg/kg) 45 minutes before cisplatin + 3 more post-cisplatin treatments	
STAT6 -/- mice (PMID: 21321603)	Cisplatin (4 mg/kg) per day for 4 days	IP	Hearing protection at 4, 16, and 32 kHz.		

Hearing loss during chemotherapy

Cytokine inhibitors Branded as Basiliximab, daclizumab	Inhibiting inflammatory cytokines	Mice treated with flunarizine (PMID: 18584244)	Cisplatin (4 mg/kg)	IP	Reduction of cochlear TNF- α and IL-1 β . No ABR recording.
		Rats treated with etanercept (PMID: 28730299)	Sibelium (143 ug/kg) Cisplatin (16 mg/kg) Etanercept (6 mg/kg) 24 hours before cisplatin treatment	Oral IP IP	
Honokiol Branded as Honobsolete	SIRT3-mediated antioxidant activity (PMID: 21172655)	Mice treated with honokiol (PMID: 33415008)	Cisplatin (20 mg/kg) Honokiol (20 mg/kg) treatment 1 hour before cisplatin treatment	IP	Protected hearing at 4.5, 6, 8, 10 kHz on day 3. Protected hearing at 4-32 kHz. Cisplatin cytotoxicity not affected.

Hearing loss during chemotherapy

mouse cochleae [110]. The reduction of cytokines was through the activation of Nrf2/HO-1 signaling. However, no results on ABR threshold changes were shown in the paper. Multiple inhibitors for CDK2 (kenpaullone and two unrevealed compounds) protect hearing against CIHL and noise-induced hearing loss [102, 208, 225], suggesting that CDK2 is an important target for hearing protection. Dabrafenib, a BRAF (a member of Raf family kinases) inhibitor, is also protective against CIHL and noise-induced hearing loss when combined with the CDK2 inhibitor AZD5438 [209]. However, the influence of cisplatin on the antitumor effect must be verified in tumor-bearing mice before its potential application in clinical practice.

Polyphenols: Polyphenols are a diverse group of naturally occurring compounds abundant in plants. Many polyphenols are natural antioxidants and anti-inflammatory agents that protect against various stress conditions such as UV light radiation and microbial infections. The roles of polyphenols in hearing protection against different insults are recognized, and the mechanisms studied most recently (for review, see [47, 226]). Epicatechin protected hearing in rats against CIHL by inhibiting ERK signaling in an early study [227, 228]. However, the protective effect of epicatechin needs further verification with cochlear whole-mount samples and frequency-specific ABR data. Resveratrol, a polyphenol abundant in fruits and red wine, was also shown to protect against CIHL in guinea pigs [229] and rats [230] at 10 and 0.1-0.5 mg/kg/day, respectively. In contrast, high doses (50, 10, and 1 mg/kg/day) were toxic to the cochlea in rats [230, 231]. The hearing protection mechanism is associated with activating the anti-oxidative response and reducing inflammatory responses mediated by NF- κ B, IL6, and IL1 β [231]. Curcumin, a bright yellow pigment extracted from the plant turmeric, is also shown to protect against CIHL in rats at 200 mg/kg [232, 233]. The mechanism is associated with decreased lipid peroxidation, potentially through the Nrf2 pathway [232]. Although arguably [234], curcumin is reactive, unstable, and non-bioavailable, which has shown to be unsuccessful in any clinical trial [235]. In addition, the interaction with the anti-tumor function of cisplatin is yet to be investigated. Luteolin is a flavone derived from celery, green pepper, and chamomile with anti-inflam-

matory, antioxidant, and anticarcinogenic properties. In a recent study, luteolin showed promising protection against CIHL by inhibiting ferroptosis [136]. The authors comprehensively studied the involvement of Gpx4, the key regulator for ferroptosis in CIHL. Luteolin protection through alleviation of ferroptosis was proven in cell cultures, cochlear explants cultures, and *in vivo* Gpx4 knockout mice levels [136]. Honokiol, a lignan extracted from the plant Magnolia, is a multifunctional small polyphenol with both antitumor effects that synergize with cisplatin [236-238] and protective effect to various tissues and organs against oxidative stress, including the brain [239-241], heart [242-244], kidney [245], liver [246], etc. In a recent study from our lab, honokiol has shown strong hearing protective effects against cisplatin ototoxicity in tumor-bearing mice undergoing chemotherapy [247]. Honokiol also increases animal survival without interfering with cisplatin's antitumor effects. The mechanism is associated with activating sirtuin 3 (SIRT3), a member of the sirtuin NAD⁺-dependent deacetylase family, critical regulators of the intrinsic anti-ROS systems.

Sirtuins comprise a highly conserved NAD⁺-dependent deacetylase family, all key regulators of the intrinsic anti-ROS systems [248-251]. Seven members (SIRT1-7) of the sirtuin family are expressed in the cytoplasm (SIRT1&2), mitochondria (SIRT3-5), and the nucleus (SIRT1, 2, 6, 7). The involvement of sirtuins (mostly SIRT3) in hearing protection has been shown in noise-induced [252, 253], drug-induced [254], and age-related hearing loss [255, 256]. Activation of sirtuins (mostly SIRT1) by polyphenols has also been reported in resveratrol [257], curcumin [258], and epicatechin [259]. Sirtuins might represent novel targets for hearing protection against cisplatin ototoxicity, and polyphenols are promising candidates. For example, honokiol does not interfere with the therapeutic effects of cisplatin, as shown in our study [247] and other publications [237, 238, 260, 261]. It can pass the BBB [262, 263] and diffuse across plasma and mitochondrial membranes *in vivo* [243, 263]. Although bioavailability is still a concern for most polyphenols because of their insolubility and instability, different ways for assisting the delivery of polyphenols, such as liposomal [264-266], amorphous solid dispersion [267, 268], and bio-

degradable microsphere [269], have made significant progress in recent years.

Summary and outlook

This paper sought to comprehensively review the current understanding and research on ototoxicity in chemotherapy (particularly with cisplatin) and explore strategies and compounds for hearing protection. Using symptoms including hearing loss, tinnitus, vertigo, and dizziness, we identified potential ototoxic drugs, among which four are specifically causing hearing loss. While the ototoxic mechanisms of cisplatin are extensively studied, other agents like isotretinoin, vinblastine, and dasatinib remain less understood.

The primary target of cisplatin in the cochlea is still debatable, although OHC loss is the most commonly observed change in cisplatin ototoxicity. Damage of the SV might be the leading cause, which affects the EP and may result in cisplatin influx into the scala media. These changes induce a series cascade of cellular processes, eventually leading to cell death. ROS generation is the predominant factor in this process, and its subsequent signaling pathways are discussed.

The review emphasizes the urgent need for effective otoprotective strategies that can mitigate the adverse effects of chemotherapy on hearing without compromising its anticancer efficacy. Various antioxidants, such as STS, NAC, and D-methionine, have been explored for their potential to counteract the ROS-mediated ototoxicity of cisplatin. STS, in particular, has gained FDA approval for reducing hearing loss in pediatric cancer patients receiving cisplatin. However, the challenge of ensuring that these agents do not interfere with the chemotherapeutic efficacy of cisplatin remains a critical consideration in their clinical application.

Emerging research has focused on more specific molecular targets and pathways involved in cisplatin-induced hearing loss. The review highlights the exploration of various compounds, including statins, protein kinase inhibitors, and polyphenols like epicatechin, resveratrol, curcumin, luteolin, and honokiol, for their protective effects against CIHL. These agents, often possessing antioxidant and anti-inflammatory properties, offer new avenues for oto-

protection, with some showing promise in pre-clinical and clinical studies. Particularly, honokiol has been noted for its synergistic antitumor effects with cisplatin and its ability to protect against ototoxicity, potentially through activating the sirtuin pathway.

The review also discusses innovative drug delivery methods, such as local administration via trans-tympanic injection and advanced formulations like nanoparticles and hydrogels, to enhance the efficacy of otoprotective agents and minimize their systemic side effects. These strategies aim to improve the bioavailability of protective agents in the cochlea and reduce the risk of interfering with the systemic anticancer activity of chemotherapeutic drugs.

In conclusion, while significant advances have been made in understanding the mechanisms of chemotherapy-induced ototoxicity and identifying potential otoprotective strategies, further research is needed to develop safe and effective treatments that can be integrated into standard oncology care. The balance between preserving hearing and maintaining the anti-neoplastic effectiveness of chemotherapy presents a complex challenge that future studies must address to improve the quality of life for cancer patients undergoing treatment.

Acknowledgements

This work is supported by grants from the NIH R01-DC019434, Hearing Health Foundation, and American Hearing Research Foundation awarded to XT.

Disclosure of conflict of interest

None.

Abbreviations

ABR, Auditory Brainstem Response; ABVD, Doxorubicin, Bleomycin, Vinblastine, Dacarbazine; AMPK, AMP-activated Protein Kinase; ARHL, Age-Related Hearing Loss; Bax, Bcl-2-Associated X Protein; Bcl2, B-cell Lymphoma 2; BBB, Blood-Brain Barrier; BLB, Blood-Labyrinthine Barrier; CAP, Compound Action Potential; CAT, Catalase; CDK, Cyclin-Dependent Kinase; CHOP, C/EBP Homologous Protein; CIHL, Cisplatin-Induced Hearing Loss; CTR1, Copper Transporter 1; DPOAE, Distortion product Oto-

Acoustic Emission; EP, Endocochlear Potential; ER, Endoplasmic Reticulum; FDA, Food and Drug Administration; Gpx4, Glutathione Peroxidase 4; GR, Glutathione Reductase; GSH.Px, Glutathione Peroxidase; HDAC, Histone Deacetylase; HDACi, Histone Deacetylase Inhibitor; HEI-OC1, House Ear Institute-Organ of Corti 1; IG, Immunoglobulin; IHCs, Inner Hair Cells; IL, Interleukin; iNOS, Inducible Nitric Oxide Synthase; i.p., Intraperitoneal; i.v., Intravenous; JAK-STAT, Janus Kinase-Signal Transducer and Activator of Transcription; LC3-II, Microtubule-associated Protein 1A/1B-light Chain 3; LRP2, LDL Receptor-related Protein 2; MAPK, Mitogen-Activated Protein Kinase; MET, Mechano-Electrical Transduction; MLKL, Mixed Lineage Kinase Domain-Like Pseudokinase; mtDNA, mitochondrial DNA; mTOR, Mammalian Target of Rapamycin; mtROS, mitochondrial ROS; NAC, N-acetyl-L-cysteine; NCI, National Cancer Institute; NF- κ B, Nuclear Factor Kappa-light-chain-enhancer of activated B cells; NLRX1, Nucleotide-binding Domain and Leucine-rich-repeat-containing Family Member X1; NOX3, NADPH Oxidase 3; Nrf2, Nuclear Factor Erythroid 2-Related Factor 2; O₂⁻, Superoxide; OCT2, Organic Cation Transporter 2; OHCs, Outer Hair Cells; ONOO⁻, Peroxynitrite; RIPK, Receptor-Interacting Protein Kinase; ROS, Reactive Oxygen Species; SGN, Spiral Ganglion Neuron; SIRT, Sirtuin; SOD, Superoxide Dismutase; STAT1, Signal Transducer and Activator of Transcription 1; STS, Sodium Thiosulfate; SV, Stria Vascularis; TNF- α , Tumor Necrosis Factor Alpha; TRPV1, Transient Receptor Potential Vanilloid 1.

Address correspondence to: Xiaodong Tan, Department of Otolaryngology-Head and Neck Surgery, Feinberg School of Medicine, Northwestern University, Chicago, Illinois 60611, USA. Tel: 001-312-503-3013; E-mail: xiaodong.tan@northwestern.edu

References

- [1] NCI. Side effects of cancer treatment - National Cancer Institute. n.d.; <https://www.cancer.gov/about-cancer/treatment/side-effects>.
- [2] NCI. Cancer Drugs - National Cancer Institute. n.d.; <https://www.cancer.gov/about-cancer/treatment/drugs>.
- [3] Chemocare. Chemotherapy Drugs and Drugs often Used During Chemotherapy. n.d.; <https://chemocare.com/chemotherapy/drug-info/default.aspx>.
- [4] Beaumont. Chemotherapy Drugs and Side Effects. 2022; <https://www.beaumont.org/treatments/chemotherapy-drugs-side-effects>.
- [5] ACS. Chemotherapy Side Effects. 2022; <https://www.cancer.org/treatment/treatments-and-side-effects/treatment-types/chemotherapy/chemotherapy-side-effects.html>.
- [6] Tavanai E, Khalili ME, Shahidipour Z, Jalaie S, Ghahraman MA, Rouhbakhsh N and Rahimi V. Hearing handicaps, communication difficulties and depression in the older adults: a comparison of hearing aid users and non-users. *Eur Arch Otorhinolaryngol* 2023; 280: 5229-5240.
- [7] Patil JD, Alrashid MA, Eltabbakh A and Fredericks S. The association between stress, emotional states, and tinnitus: a mini-review. *Front Aging Neurosci* 2023; 15: 1131979.
- [8] Brewster KK, Deal JA, Lin FR and Rutherford BR. Considering hearing loss as a modifiable risk factor for dementia. *Expert Rev Neurother* 2022; 22: 805-813.
- [9] Minasian LM, Frazier AL, Sung L, O'Mara A, Kelaghan J, Chang KW, Krailo M, Pollock BH, Reaman G and Freyer DR. Prevention of cisplatin-induced hearing loss in children: informing the design of future clinical trials. *Cancer Med* 2018; 7: 2951-2959.
- [10] Dhillon S. Sodium thiosulfate: pediatric first approval. *Paediatr Drugs* 2023; 25: 239-244.
- [11] Saito T, Zhang ZJ, Tokuriki M, Shibamori Y, Yamamoto T, Noda I, Ohtsubo T and Saito H. Doxorubicin ototoxicity is induced in mice by combination treatment with cyclosporin A. *Acta Otolaryngol* 2001; 121: 787-793.
- [12] Huang D, Savage SR, Calinawan AP, Lin C, Zhang B, Wang P, Starr TK, Birrer MJ and Paulovich AG. A highly annotated database of genes associated with platinum resistance in cancer. *Oncogene* 2021; 40: 6395-6405.
- [13] Rybak LP, Mukherjea D, Jajoo S and Ramkumar V. Cisplatin ototoxicity and protection: clinical and experimental studies. *Tohoku J Exp Med* 2009; 219: 177-186.
- [14] Schacht J, Talaska AE and Rybak LP. Cisplatin and aminoglycoside antibiotics: hearing loss and its prevention. *Anat Rec (Hoboken)* 2012; 295: 1837-1850.
- [15] Rybak LP. Mechanisms of cisplatin ototoxicity and progress in otoprotection. *Curr Opin Otolaryngol Head Neck Surg* 2007; 15: 364-369.
- [16] Rybak LP, Whitworth CA, Mukherjea D and Ramkumar V. Mechanisms of cisplatin-induced ototoxicity and prevention. *Hear Res* 2007; 226: 157-167.
- [17] Kelland L. The resurgence of platinum-based cancer chemotherapy. *Nat Rev Cancer* 2007; 7: 573-584.
- [18] Benedetti Panici P, Gregg S, Scambia G, Baiocchi G, Lomonaco M, Conti G and Mancuso S.

Hearing loss during chemotherapy

- Efficacy and toxicity of very high-dose cisplatin in advanced ovarian carcinoma: 4-year survival analysis and neurological follow-up. *Int J Gynecol Cancer* 1993; 3: 44-53.
- [19] Yuce S, Seker MM, Koc S, Uysal IO, Kacan T, Dogan M, Dogan M, Babacan NA and Kilickap S. Oxaliplatin and ototoxicity: is it really safe for hearing? *Turk J Med Sci* 2014; 44: 586-589.
- [20] Gersten BK, Fitzgerald TS, Fernandez KA and Cunningham LL. Ototoxicity and platinum uptake following cyclic administration of platinum-based chemotherapeutic agents. *J Assoc Res Otolaryngol* 2020; 21: 303-321.
- [21] Oh SY, Wasif N, Garcon MC, Rodriguez G and Saif MW. Ototoxicity associated with oxaliplatin in a patient with pancreatic cancer. *JOP* 2013; 14: 676-679.
- [22] Kim SB, Kim SY, Kim KH and Kim TN. Oxaliplatin-induced sudden hearing loss in a patient with pancreatic cancer. *Korean J Gastroenterol* 2020; 76: 261-264.
- [23] Guvenc MG, Dizdar D, Dizdar SK, Okutur SK and Demir G. Sudden hearing loss due to oxaliplatin use in a patient with colon cancer. *J Chemother* 2016; 28: 341-342.
- [24] Moss PE, Hickman S and Harrison BR. Ototoxicity associated with vinblastine. *Ann Pharmacother* 1999; 33: 423-425.
- [25] Nirban RK, Kapoor A, Narayan S, Maharia S, Singhal MK and Kumar HS. Ototoxicity following Vinblastine chemotherapy in a patient of Hodgkin's Lymphoma. *Indian J Otolaryngol* 2015; 21: 229-230.
- [26] Lugassy G and Shapira A. Sensorineural hearing loss associated with vincristine treatment. *Blut* 1990; 61: 320-321.
- [27] Aydogdu I, Ozturan O, Kuku I, Kaya E, Sevinc A and Yildiz R. Bilateral transient hearing loss associated with vincristine therapy: case report. *J Chemother* 2000; 12: 530-532.
- [28] Kalcioğlu MT, Kuku I, Kaya E, Oncel S and Aydogdu I. Bilateral hearing loss during vincristine therapy: a case report. *J Chemother* 2003; 15: 290-292.
- [29] Rosende L, Vereá-Hernando MM, de Andres A, Pineyro-Molina F, Barja J, Castro-Castro S and Fonseca E. Hypoacusia in a patient treated by isotretinoin. *Case Rep Med* 2011; 2011: 789143.
- [30] Lee JH, Lee JH and Lee JM. All-trans retinoic acid-induced ototoxicity during chemotherapy in pediatric acute promyelocytic leukemia. *Children (Basel)* 2021; 8: 27.
- [31] Akdag M, Akkurt ZM, Gul A, Ucmak D, Yilmaz B, Sengul E and Topcu I. The effects of oral isotretinoin (13-Cis retinoic acid) on the inner ear: a prospective clinical study. *Clin Invest Med* 2014; 37: E102-E107.
- [32] Karabulut H, Karadag AS, Acar B, Dagli M, Karabulut I, Ozmen E, Babademez MA and Karasen RM. The effect of oral isotretinoin (13-cis retinoic acid) on hearing systems in patients with acne vulgaris: a prospective study. *Int J Dermatol* 2011; 50: 1139-1143.
- [33] Bragado P, Armesilla A, Silva A and Porras A. Apoptosis by cisplatin requires p53 mediated p38alpha MAPK activation through ROS generation. *Apoptosis* 2007; 12: 1733-1742.
- [34] Choi YM, Kim HK, Shim W, Anwar MA, Kwon JW, Kwon HK, Kim HJ, Jeong H, Kim HM, Hwang D, Kim HS and Choi S. Mechanism of cisplatin-induced cytotoxicity is correlated to impaired metabolism due to mitochondrial ROS generation. *PLoS One* 2015; 10: e0135083.
- [35] Pinto AL and Lippard SJ. Binding of the antitumor drug cis-diamminedichloroplatinum(II) (cisplatin) to DNA. *Biochim Biophys Acta* 1985; 780: 167-180.
- [36] Chvalova K, Brabec V and Kasparkova J. Mechanism of the formation of DNA-protein cross-links by antitumor cisplatin. *Nucleic Acids Res* 2007; 35: 1812-1821.
- [37] Pan B, Akyuz N, Liu XP, Asai Y, Nist-Lund C, Kurima K, Derfler BH, Gyorgy B, Limapichat W, Walujkar S, Wimalasena LN, Sotomayor M, Corey DP and Holt JR. TMC1 forms the pore of mechanosensory transduction channels in vertebrate inner ear hair cells. *Neuron* 2018; 99: 736-753, e736.
- [38] Beurg M, Evans MG, Hackney CM and Fettiplace R. A large-conductance calcium-selective mechanotransducer channel in mammalian cochlear hair cells. *J Neurosci* 2006; 26: 10992-11000.
- [39] Abeytunge S, Gianoli F, Hudspeth AJ and Kozlov AS. Rapid mechanical stimulation of inner-ear hair cells by photonic pressure. *Elife* 2021; 10: e65930.
- [40] Caprara GA and Peng AW. Mechanotransduction in mammalian sensory hair cells. *Mol Cell Neurosci* 2022; 120: 103706.
- [41] Ó Maoiléidigh D and Ricci AJ. A bundle of mechanisms: inner-ear hair-cell mechanotransduction. *Trends Neurosci* 2019; 42: 221-236.
- [42] Wangemann P. K⁺ cycling and the endocochlear potential. *Hear Res* 2002; 165: 1-9.
- [43] Wangemann P, Liu J and Marcus DC. Ion transport mechanisms responsible for K⁺ secretion and the transepithelial voltage across marginal cells of stria vascularis in vitro. *Hear Res* 1995; 84: 19-29.
- [44] Wangemann P. Supporting sensory transduction: cochlear fluid homeostasis and the endocochlear potential. *J Physiol* 2006; 576: 11-21.

Hearing loss during chemotherapy

- [45] Locher H, de Groot JC, van Iperen L, Huisman MA, Frijns JH and Chuva de Sousa Lopes SM. Development of the stria vascularis and potassium regulation in the human fetal cochlea: Insights into hereditary sensorineural hearing loss. *Dev Neurobiol* 2015; 75: 1219-1240.
- [46] Prayuenyong P, Baguley DM, Kros CJ and Steyger PS. Preferential cochleotoxicity of cisplatin. *Front Neurosci* 2021; 15: 695268.
- [47] Tan WJT and Vlajkovic SM. Molecular characteristics of cisplatin-induced ototoxicity and therapeutic interventions. *Int J Mol Sci* 2023; 24: 16545.
- [48] Nyberg S, Abbott NJ, Shi X, Steyger PS and Dabdoub A. Delivery of therapeutics to the inner ear: the challenge of the blood-labyrinth barrier. *Sci Transl Med* 2019; 11: eaao0935.
- [49] Shi X. Pathophysiology of the cochlear intrastrial fluid-blood barrier (review). *Hear Res* 2016; 338: 52-63.
- [50] Fetoni AR, Troiani D, Eramo SL, Rolesi R and Paludetti Troiani G. Efficacy of different routes of administration for Coenzyme Q10 formulation in noise-induced hearing loss: systemic versus transtympanic modality. *Acta Otolaryngol* 2012; 132: 391-399.
- [51] Howell SB and Taetle R. Effect of sodium thiosulfate on cis-dichlorodiammineplatinum(II) toxicity and antitumor activity in L1210 leukemia. *Cancer Treat Rep* 1980; 64: 611-616.
- [52] Paken J, Govender CD, Pillay M and Sewram V. A review of cisplatin-associated ototoxicity. *Semin Hear* 2019; 40: 108-121.
- [53] Dasari S and Tchounwou PB. Cisplatin in cancer therapy: molecular mechanisms of action. *Eur J Pharmacol* 2014; 740: 364-378.
- [54] McAlpine D and Johnstone BM. The ototoxic mechanism of cisplatin. *Hear Res* 1990; 47: 191-203.
- [55] Ravi R, Somani SM and Rybak LP. Mechanism of cisplatin ototoxicity: antioxidant system. *Pharmacol Toxicol* 1995; 76: 386-394.
- [56] Sheth S, Mukherjea D, Rybak LP and Ramkumar V. Mechanisms of cisplatin-induced ototoxicity and otoprotection. *Front Cell Neurosci* 2017; 11: 338.
- [57] Karasawa T and Steyger PS. An integrated view of cisplatin-induced nephrotoxicity and ototoxicity. *Toxicol Lett* 2015; 237: 219-227.
- [58] Lee CH, Lee DH, Lee SM and Kim ASY. Otoprotective effects of Zingerone on cisplatin-induced ototoxicity. *Int J Mol Sci* 2020; 21: 3503.
- [59] Gentilin E, Simoni E, Candito M, Cazzador D and Astolfi L. Cisplatin-induced ototoxicity: updates on molecular targets. *Trends Mol Med* 2019; 25: 1123-1132.
- [60] Anfuso CD, Cosentino A, Agafonova A, Zappala A, Giurdanella G, Trovato Salinaro A, Calabrese V and Lupo G. Pericytes of stria vascularis are targets of cisplatin-induced ototoxicity: new insights into the molecular mechanisms involved in blood-labyrinth barrier breakdown. *Int J Mol Sci* 2022; 23: 15790.
- [61] Sluyter S, Klis SF, de Groot JC and Smoorenburg GF. Alterations in the stria vascularis in relation to cisplatin ototoxicity and recovery. *Hear Res* 2003; 185: 49-56.
- [62] Freitas MR, Silva VC, Brito GA, Carvalho Junior JV, Gomes Junior RM and Ribeiro Rde A. Distortion-product otoacoustic emissions and auditory brainstem responses sensitivity assessment in cisplatin-induced ototoxicity in rats. *Braz J Otorhinolaryngol* 2009; 75: 476-484.
- [63] Chen Y, Bielefeld EC, Mellott JG, Wang W, Mafi AM, Yamoah EN and Bao J. Early physiological and cellular indicators of cisplatin-induced ototoxicity. *J Assoc Res Otolaryngol* 2021; 22: 107-126.
- [64] Borse V, Al Aameri RFH, Sheehan K, Sheth S, Kaur T, Mukherjea D, Tupal S, Lowy M, Ghosh S, Dhukhwa A, Bhatta P, Rybak LP and Ramkumar V. Epigallocatechin-3-gallate, a prototypic chemopreventative agent for protection against cisplatin-based ototoxicity. *Cell Death Dis* 2017; 8: e2921.
- [65] Thomas AJ, Hailey DW, Stawicki TM, Wu P, Coffin AB, Rubel EW, Raible DW, Simon JA and Ou HC. Functional mechanotransduction is required for cisplatin-induced hair cell death in the zebrafish lateral line. *J Neurosci* 2013; 33: 4405-4414.
- [66] More SS, Akil O, Ianculescu AG, Geier EG, Lustig LR and Giacomini KM. Role of the copper transporter, CTR1, in platinum-induced ototoxicity. *J Neurosci* 2010; 30: 9500-9509.
- [67] Ciarimboli G, Deuster D, Knief A, Sperling M, Holtkamp M, Edemir B, Pavenstadt H, Lanvers-Kaminsky C, am Zehnhoff-Dinnesen A, Schinkel AH, Koepsell H, Jurgens H and Schlatter E. Organic cation transporter 2 mediates cisplatin-induced oto- and nephrotoxicity and is a target for protective interventions. *Am J Pathol* 2010; 176: 1169-1180.
- [68] Breglio AM, Rusheen AE, Shide ED, Fernandez KA, Spielbauer KK, McLachlin KM, Hall MD, Amable L and Cunningham LL. Cisplatin is retained in the cochlea indefinitely following chemotherapy. *Nat Commun* 2017; 8: 1654.
- [69] Thomas JP, Lautermann J, Liedert B, Seiler F and Thomale J. High accumulation of platinum-DNA adducts in stria marginal cells of the cochlea is an early event in cisplatin but not carboplatin ototoxicity. *Mol Pharmacol* 2006; 70: 23-29.
- [70] Ishida S, Lee J, Thiele DJ and Herskowitz I. Uptake of the anticancer drug cisplatin mediated by the copper transporter Ctr1 in yeast and mammals. *Proc Natl Acad Sci U S A* 2002; 99: 14298-14302.

Hearing loss during chemotherapy

- [71] Riedemann L, Lanvers C, Deuster D, Peters U, Boos J, Jurgens H and am Zehnhoff-Dinnesen A. Megalin genetic polymorphisms and individual sensitivity to the ototoxic effect of cisplatin. *Pharmacogenomics J* 2008; 8: 23-28.
- [72] Tang Q, Wang X, Jin H, Mi Y, Liu L, Dong M, Chen Y and Zou Z. Cisplatin-induced ototoxicity: updates on molecular mechanisms and otoprotective strategies. *Eur J Pharm Biopharm* 2021; 163: 60-71.
- [73] Kimitsuki T, Nakagawa T, Hisashi K, Komune S and Komiyama S. Cisplatin blocks mechanoelectric transducer current in chick cochlear hair cells. *Hear Res* 1993; 71: 64-68.
- [74] Li J, Liu C, Kaefer S, Youssef M and Zhao B. The mechanotransduction channel and organic cation transporter are critical for cisplatin ototoxicity in murine hair cells. *Front Mol Neurosci* 2022; 15: 835448.
- [75] van Ruijven MW, de Groot JC and Smoorenburg GF. Time sequence of degeneration pattern in the guinea pig cochlea during cisplatin administration. A quantitative histological study. *Hear Res* 2004; 197: 44-54.
- [76] Binks SP and Dobrota M. Kinetics and mechanism of uptake of platinum-based pharmaceuticals by the rat small intestine. *Biochem Pharmacol* 1990; 40: 1329-1336.
- [77] Previati M, Lanzoni I, Astolfi L, Fagioli F, Vecchiati G, Pagnoni A, Martini A and Capitani S. Cisplatin cytotoxicity in organ of Corti-derived immortalized cells. *J Cell Biochem* 2007; 101: 1185-1197.
- [78] Hazlitt RA, Min J and Zuo J. Progress in the development of preventative drugs for cisplatin-induced hearing loss. *J Med Chem* 2018; 61: 5512-5524.
- [79] Yu D, Gu J, Chen Y, Kang W, Wang X and Wu H. Current strategies to combat cisplatin-induced ototoxicity. *Front Pharmacol* 2020; 11: 999.
- [80] Rybak LP, Husain K, Morris C, Whitworth C and Somani S. Effect of protective agents against cisplatin ototoxicity. *Am J Otol* 2000; 21: 513-520.
- [81] Banfi B, Malgrange B, Knisz J, Steger K, Dubois-Dauphin M and Krause KH. NOX3, a superoxide-generating NADPH oxidase of the inner ear. *J Biol Chem* 2004; 279: 46065-46072.
- [82] Mukherjea D, Jajoo S, Sheehan K, Kaur T, Sheth S, Bunch J, Perro C, Rybak LP and Ramkumar V. NOX3 NADPH oxidase couples transient receptor potential vanilloid 1 to signal transducer and activator of transcription 1-mediated inflammation and hearing loss. *Antioxid Redox Signal* 2011; 14: 999-1010.
- [83] Wu J, Ye J, Kong W, Zhang S and Zheng Y. Programmed cell death pathways in hearing loss: a review of apoptosis, autophagy and programmed necrosis. *Cell Prolif* 2020; 53: e12915.
- [84] Mukherjea D, Jajoo S, Whitworth C, Bunch JR, Turner JG, Rybak LP and Ramkumar V. Short interfering RNA against transient receptor potential vanilloid 1 attenuates cisplatin-induced hearing loss in the rat. *J Neurosci* 2008; 28: 13056-13065.
- [85] Kaur T, Mukherjea D, Sheehan K, Jajoo S, Rybak LP and Ramkumar V. Short interfering RNA against STAT1 attenuates cisplatin-induced ototoxicity in the rat by suppressing inflammation. *Cell Death Dis* 2011; 2: e180.
- [86] Rybak LP, Whitworth C and Somani S. Application of antioxidants and other agents to prevent cisplatin ototoxicity. *Laryngoscope* 1999; 109: 1740-1744.
- [87] Lautermann J, Crann SA, McLaren J and Schacht J. Glutathione-dependent antioxidant systems in the mammalian inner ear: effects of aging, ototoxic drugs and noise. *Hear Res* 1997; 114: 75-82.
- [88] Paken J, Govender CD, Pillay M and Sewram V. Cisplatin-associated ototoxicity: a review for the health professional. *J Toxicol* 2016; 2016: 1809394.
- [89] Ramkumar V, Mukherjea D, Dhukhwa A and Rybak LP. Oxidative stress and inflammation caused by cisplatin ototoxicity. *Antioxidants (Basel)* 2021; 10: 1919.
- [90] Garcia-Berrocal JR, Nevado J, Ramirez-Camacho R, Sanz R, Gonzalez-Garcia JA, Sanchez-Rodriguez C, Cantos B, Espana P, Verdaguier JM and Trinidad Cabezas A. The anticancer drug cisplatin induces an intrinsic apoptotic pathway inside the inner ear. *Br J Pharmacol* 2007; 152: 1012-1020.
- [91] Kamogashira T, Fujimoto C and Yamasoba T. Reactive oxygen species, apoptosis, and mitochondrial dysfunction in hearing loss. *Biomed Res Int* 2015; 2015: 617207.
- [92] Wong AC and Ryan AF. Mechanisms of sensorineural cell damage, death and survival in the cochlea. *Front Aging Neurosci* 2015; 7: 58.
- [93] Romano A, Capozza MA, Mastrangelo S, Maurizi P, Triarico S, Rolesi R, Attina G, Fetoni AR and Ruggiero A. Assessment and management of platinum-related ototoxicity in children treated for cancer. *Cancers (Basel)* 2020; 12: 1266.
- [94] Benkafadar N, Menardo J, Bourien J, Nouvian R, Francois F, Decaudin D, Maiorano D, Puel JL and Wang J. Reversible p53 inhibition prevents cisplatin ototoxicity without blocking chemotherapeutic efficacy. *EMBO Mol Med* 2017; 9: 7-26.
- [95] Kim HJ, Lee JH, Kim SJ, Oh GS, Moon HD, Kwon KB, Park C, Park BH, Lee HK, Chung SY, Park R and So HS. Roles of NADPH oxidases in cisplatin-induced reactive oxygen species generation and ototoxicity. *J Neurosci* 2010; 30: 3933-3946.

Hearing loss during chemotherapy

- [96] So HS, Park C, Kim HJ, Lee JH, Park SY, Lee JH, Lee ZW, Kim HM, Kalinec F, Lim DJ and Park R. Protective effect of T-type calcium channel blocker flunarizine on cisplatin-induced death of auditory cells. *Hear Res* 2005; 204: 127-139.
- [97] Monroe JD, Johnston AM and Smith ME. The monofunctional platinum(II) compounds, phenanthriplatin and pyriplatin, modulate apoptosis signaling pathways in HEI-OC1 auditory hybridoma cells. *Neurotoxicology* 2020; 79: 104-109.
- [98] Lee HS, Kim YR, Lee IK, Kim UK, Baek JI and Lee KY. KL1333, a derivative of beta-lapachone, protects against cisplatin-induced ototoxicity in mouse cochlear cultures. *Biomed Pharmacother* 2020; 126: 110068.
- [99] Murata T, Hibasami H, Maekawa S, Tagawa T and Nakashima K. Preferential binding of cisplatin to mitochondrial DNA and suppression of ATP generation in human malignant melanoma cells. *Biochem Int* 1990; 20: 949-955.
- [100] Olivero OA, Semino C, Kassim A, Lopez-Larraz DM and Poirier MC. Preferential binding of cisplatin to mitochondrial DNA of Chinese hamster ovary cells. *Mutat Res* 1995; 346: 221-230.
- [101] Yang Z, Schumaker LM, Egorin MJ, Zuhowski EG, Guo Z and Cullen KJ. Cisplatin preferentially binds mitochondrial DNA and voltage-dependent anion channel protein in the mitochondrial membrane of head and neck squamous cell carcinoma: possible role in apoptosis. *Clin Cancer Res* 2006; 12: 5817-5825.
- [102] Teitz T, Fang J, Goktug AN, Bonga JD, Diao S, Hazlitt RA, Iconaru L, Morfouace M, Currier D, Zhou Y, Umans RA, Taylor MR, Cheng C, Min J, Freeman B, Peng J, Roussel MF, Kriwacki R, Guy RK, Chen T and Zuo J. CDK2 inhibitors as candidate therapeutics for cisplatin- and noise-induced hearing loss. *J Exp Med* 2018; 215: 1187-1203.
- [103] Devarajan P, Savoca M, Castaneda MP, Park MS, Esteban-Cruciani N, Kalinec G and Kalinec F. Cisplatin-induced apoptosis in auditory cells: role of death receptor and mitochondrial pathways. *Hear Res* 2002; 174: 45-54.
- [104] Zong S, Liu T, Wan F, Chen P, Luo P and Xiao H. Endoplasmic reticulum stress is involved in cochlear cell apoptosis in a cisplatin-induced ototoxicity rat model. *Audiol Neurootol* 2017; 22: 160-168.
- [105] Kishino A, Hayashi K, Maeda M, Jike T, Hidai C, Nomura Y and Oshima T. Caspase-8 regulates endoplasmic reticulum stress-induced necroptosis independent of the apoptosis pathway in auditory cells. *Int J Mol Sci* 2019; 20: 5896.
- [106] Hu H, Tian M, Ding C and Yu S. The C/EBP homologous protein (CHOP) transcription factor functions in endoplasmic reticulum stress-induced apoptosis and microbial infection. *Front Immunol* 2018; 9: 3083.
- [107] Chung WH, Boo SH, Chung MK, Lee HS, Cho YS and Hong SH. Proapoptotic effects of NF-kappaB on cisplatin-induced cell death in auditory cell line. *Acta Otolaryngol* 2008; 128: 1063-1070.
- [108] Babolmorad G, Latif A, Domingo IK, Pollock NM, Delyea C, Rieger AM, Allison WT and Bhavsar AP. Toll-like receptor 4 is activated by platinum and contributes to cisplatin-induced ototoxicity. *EMBO Rep* 2021; 22: e51280.
- [109] So H, Kim H, Lee JH, Park C, Kim Y, Kim E, Kim JK, Yun KJ, Lee KM, Lee HY, Moon SK, Lim DJ and Park R. Cisplatin cytotoxicity of auditory cells requires secretions of proinflammatory cytokines via activation of ERK and NF-kappaB. *J Assoc Res Otolaryngol* 2007; 8: 338-355.
- [110] So H, Kim H, Kim Y, Kim E, Pae HO, Chung HT, Kim HJ, Kwon KB, Lee KM, Lee HY, Moon SK and Park R. Evidence that cisplatin-induced auditory damage is attenuated by downregulation of pro-inflammatory cytokines via Nrf2/HO-1. *J Assoc Res Otolaryngol* 2008; 9: 290-306.
- [111] Watanabe K, Inai S, Jinnouchi K, Bada S, Hess A, Michel O and Yagi T. Nuclear-factor kappa B (NF-kappa B)-inducible nitric oxide synthase (iNOS/NOS II) pathway damages the stria vascularis in cisplatin-treated mice. *Anticancer Res* 2002; 22: 4081-4085.
- [112] Beckman JS, Beckman TW, Chen J, Marshall PA and Freeman BA. Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci U S A* 1990; 87: 1620-1624.
- [113] Lee JE, Nakagawa T, Kita T, Kim TS, Iguchi F, Endo T, Shiga A, Lee SH and Ito J. Mechanisms of apoptosis induced by cisplatin in marginal cells in mouse stria vascularis. *ORL J Otorhinolaryngol Relat Spec* 2004; 66: 111-118.
- [114] Wang J, Ladrech S, Pujol R, Brabet P, Van De Water TR and Puel JL. Caspase inhibitors, but not c-Jun NH2-terminal kinase inhibitor treatment, prevent cisplatin-induced hearing loss. *Cancer Res* 2004; 64: 9217-9224.
- [115] Achkar IW, Abdulrahman N, Al-Sulaiti H, Joseph JM, Uddin S and Mraiche F. Cisplatin based therapy: the role of the mitogen activated protein kinase signaling pathway. *J Transl Med* 2018; 16: 96.
- [116] Ruhl D, Du TT, Wagner EL, Choi JH, Li S, Reed R, Kim K, Freeman M, Hashisaki G, Lukens JR and Shin JB. Necroptosis and apoptosis con-

Hearing loss during chemotherapy

- tribute to cisplatin and aminoglycoside ototoxicity. *J Neurosci* 2019; 39: 2951-2964.
- [117] Morris R, Kershaw NJ and Babon JJ. The molecular details of cytokine signaling via the JAK/STAT pathway. *Protein Sci* 2018; 27: 1984-2009.
- [118] Kaur T, Borse V, Sheth S, Sheehan K, Ghosh S, Tupal S, Jajoo S, Mukherjea D, Rybak LP and Ramkumar V. Adenosine A1 receptor protects against cisplatin ototoxicity by suppressing the NOX3/STAT1 inflammatory pathway in the cochlea. *J Neurosci* 2016; 36: 3962-3977.
- [119] Kim HJ, Oh GS, Lee JH, Lyu AR, Ji HM, Lee SH, Song J, Park SJ, You YO, Sul JD, Park C, Chung SY, Moon SK, Lim DJ, So HS and Park R. Cisplatin ototoxicity involves cytokines and STAT6 signaling network. *Cell Res* 2011; 21: 944-956.
- [120] Ikeda K, Sunose H and Takasaka T. Effects of free radicals on the intracellular calcium concentration in the isolated outer hair cell of the guinea pig cochlea. *Acta Otolaryngol* 1993; 113: 137-141.
- [121] Yoshida T, Inoue R, Morii T, Takahashi N, Yamamoto S, Hara Y, Tominaga M, Shimizu S, Sato Y and Mori Y. Nitric oxide activates TRP channels by cysteine S-nitrosylation. *Nat Chem Biol* 2006; 2: 596-607.
- [122] Kurabi A, Keithley EM, Housley GD, Ryan AF and Wong AC. Cellular mechanisms of noise-induced hearing loss. *Hear Res* 2017; 349: 129-137.
- [123] Choi MJ, Kang H, Lee YY, Choo OS, Jang JH, Park SH, Moon JS, Choi SJ and Choung YH. Cisplatin-induced ototoxicity in rats is driven by RIP3-dependent necroptosis. *Cells* 2019; 8: 409.
- [124] Dhuriya YK and Sharma D. Necroptosis: a regulated inflammatory mode of cell death. *J Neuroinflammation* 2018; 15: 199.
- [125] Cai Z, Jitkaew S, Zhao J, Chiang HC, Choksi S, Liu J, Ward Y, Wu LG and Liu ZG. Plasma membrane translocation of trimerized MLKL protein is required for TNF-induced necroptosis. *Nat Cell Biol* 2014; 16: 55-65.
- [126] Yin H, Yang Q, Cao Z, Li H, Yu Z, Zhang G, Sun G, Man R, Wang H and Li J. Activation of NLRX1-mediated autophagy accelerates the ototoxic potential of cisplatin in auditory cells. *Toxicol Appl Pharmacol* 2018; 343: 16-28.
- [127] Yin H, Sun G, Yang Q, Chen C, Qi Q, Wang H and Li J. NLRX1 accelerates cisplatin-induced ototoxicity in HEI-OC1 cells via promoting generation of ROS and activation of JNK signaling pathway. *Sci Rep* 2017; 7: 44311.
- [128] Fang B and Xiao H. Rapamycin alleviates cisplatin-induced ototoxicity in vivo. *Biochem Biophys Res Commun* 2014; 448: 443-447.
- [129] Youn CK, Kim J, Park JH, Do NY and Cho SI. Role of autophagy in cisplatin-induced ototoxicity. *Int J Pediatr Otorhinolaryngol* 2015; 79: 1814-1819.
- [130] Yang Q, Sun G, Yin H, Li H, Cao Z, Wang J, Zhou M, Wang H and Li J. PINK1 protects auditory hair cells and spiral ganglion neurons from cisplatin-induced ototoxicity via inducing autophagy and inhibiting JNK signaling pathway. *Free Radic Biol Med* 2018; 120: 342-355.
- [131] Cho SI, Jo ER and Song H. Mitophagy impairment aggravates cisplatin-induced ototoxicity. *Biomed Res Int* 2021; 2021: 5590973.
- [132] Li Y, Zhang T, Song Q, Gao D, Li Y, Jie H, Huang P, Zheng G, Yang J and He J. Cisplatin ototoxicity mechanism and antagonistic intervention strategy: a scope review. *Front Cell Neurosci* 2023; 17: 1197051.
- [133] Wang X, Zhou Y, Wang D, Wang Y, Zhou Z, Ma X, Liu X and Dong Y. Cisplatin-induced ototoxicity: from signaling network to therapeutic targets. *Biomed Pharmacother* 2023; 157: 114045.
- [134] Yang WS and Stockwell BR. Ferroptosis: death by lipid peroxidation. *Trends Cell Biol* 2016; 26: 165-176.
- [135] Jian B, Pang J, Xiong H, Zhang W, Zhan T, Su Z, Lin H, Zhang H, He W and Zheng Y. Autophagy-dependent ferroptosis contributes to cisplatin-induced hearing loss. *Toxicol Lett* 2021; 350: 249-260.
- [136] Liu Z, Zhang H, Hong G, Bi X, Hu J, Zhang T, An Y, Guo N, Dong F, Xiao Y, Li W, Zhao X, Chu B, Guo S, Zhang X, Chai R and Fu X. Inhibition of Gpx4-mediated ferroptosis alleviates cisplatin-induced hearing loss in C57BL/6 mice. *Mol Ther* 2024; 32: 1387-1406.
- [137] Hu B, Liu Y, Chen X, Zhao J, Han J, Dong H, Zheng Q and Nie G. Ferrostatin-1 protects auditory hair cells from cisplatin-induced ototoxicity in vitro and in vivo. *Biochem Biophys Res Commun* 2020; 533: 1442-1448.
- [138] Mei H, Zhao L, Li W, Zheng Z, Tang D, Lu X and He Y. Inhibition of ferroptosis protects House Ear Institute-Organ of Corti 1 cells and cochlear hair cells from cisplatin-induced ototoxicity. *J Cell Mol Med* 2020; 24: 12065-12081.
- [139] Wang W, Ma P, Gao W, Lu P, Ding X, Chen J, Yuan H and Lu L. Nrf2 knockout affected the ferroptosis signaling pathway against cisplatin-induced hair cell-like HEI-OC1 cell death. *Oxid Med Cell Longev* 2022; 2022: 2210733.
- [140] Cho S, Hong SJ, Kang SH, Park Y and Kim SK. Alpha-lipoic acid attenuates apoptosis and ferroptosis in cisplatin-induced ototoxicity via the reduction of intracellular lipid droplets. *Int J Mol Sci* 2022; 23: 10981.
- [141] Neuwelt EA, Brummett RE, Doolittle ND, Muldoon LL, Kroll RA, Pagel MA, Dojan R, Church V, Remsen LG and Bubalo JS. First evidence of otoprotection against carboplatin-induced

Hearing loss during chemotherapy

- hearing loss with a two-compartment system in patients with central nervous system malignancy using sodium thiosulfate. *J Pharmacol Exp Ther* 1998; 286: 77-84.
- [142] Brock PR, Knight KR, Freyer DR, Campbell KC, Steyger PS, Blakley BW, Rassekh SR, Chang KW, Fligor BJ, Rajput K, Sullivan M and Neuwelt EA. Platinum-induced ototoxicity in children: a consensus review on mechanisms, predisposition, and protection, including a new International Society of Pediatric Oncology Boston ototoxicity scale. *J Clin Oncol* 2012; 30: 2408-2417.
- [143] Videhult P, Laurell G, Wallin I and Ehrsson H. Kinetics of Cisplatin and its monohydrated complex with sulfur-containing compounds designed for local otoprotective administration. *Exp Biol Med (Maywood)* 2006; 231: 1638-1645.
- [144] Sooriyaarachchi M, George GN, Pickering IJ, Narendran A and Gailer J. Tuning the metabolism of the anticancer drug cisplatin with chemoprotective agents to improve its safety and efficacy. *Metallomics* 2016; 8: 1170-1176.
- [145] Herr I, Ucur E, Herzer K, Okouoyo S, Ridder R, Krammer PH, von Knebel Doeberitz M and Debatin KM. Glucocorticoid cotreatment induces apoptosis resistance toward cancer therapy in carcinomas. *Cancer Res* 2003; 63: 3112-3120.
- [146] Poirrier AL, Pincemail J, Van Den Ackerveken P, Lefebvre PP and Malgrange B. Oxidative stress in the cochlea: an update. *Curr Med Chem* 2010; 17: 3591-3604.
- [147] Fouladi M, Chintagumpala M, Ashley D, Kellie S, Gururangan S, Hassall T, Gronewold L, Stewart CF, Wallace D, Broniscer A, Hale GA, Kasow KA, Merchant TE, Morris B, Krasin M, Kun LE, Boyett JM and Gajjar A. Amifostine protects against cisplatin-induced ototoxicity in children with average-risk medulloblastoma. *J Clin Oncol* 2008; 26: 3749-3755.
- [148] Wang J, Lloyd Faulconbridge RV, Fetoni A, Guitton MJ, Pujol R and Puel JL. Local application of sodium thiosulfate prevents cisplatin-induced hearing loss in the guinea pig. *Neuropharmacology* 2003; 45: 380-393.
- [149] Bear ZW and Mikulec AA. Intratympanic steroid therapy for treatment of idiopathic sudden sensorineural hearing loss. *Mo Med* 2014; 111: 352-356.
- [150] Hill GW, Morest DK and Parham K. Cisplatin-induced ototoxicity: effect of intratympanic dexamethasone injections. *Otol Neurotol* 2008; 29: 1005-1011.
- [151] Calli C, Pinar E, Oncel S, Alper Bagriyanik H and Umut Sakarya E. Recovery of hearing in Cisplatin-induced ototoxicity in the Guinea pig with intratympanic dexamethasone. *Indian J Otolaryngol Head Neck Surg* 2012; 64: 46-50.
- [152] Wimmer C, Mees K, Stumpf P, Welsch U, Reichel O and Suckfull M. Round window application of D-methionine, sodium thiosulfate, brain-derived neurotrophic factor, and fibroblast growth factor-2 in cisplatin-induced ototoxicity. *Otol Neurotol* 2004; 25: 33-40.
- [153] Wolters FL, Klis SF, Hamers FP, de Groot JC and Smoorenburg GF. Perilymphatic application of alpha-melanocyte stimulating hormone ameliorates hearing loss caused by systemic administration of cisplatin. *Hear Res* 2004; 189: 31-40.
- [154] Sarafraz Z, Ahmadi A and Daneshi A. Transtympanic injections of n-acetylcysteine and dexamethasone for prevention of cisplatin-induced ototoxicity: double blind randomized clinical trial. *Int Tinnitus J* 2018; 22: 40-45.
- [155] Meyer F, Rolland V, Bairati I, Guitton M, Fortin A and Côté M. Randomized controlled trial to test the efficacy of trans-tympanic injections of a sodium thiosulfate gel to prevent cisplatin-induced ototoxicity. *J Clin Oncol* 2017; 35: e17560.
- [156] Marshak T, Steiner M, Kaminer M, Levy L and Shupak A. Prevention of cisplatin-induced hearing loss by intratympanic dexamethasone: a randomized controlled study. *Otolaryngol Head Neck Surg* 2014; 150: 983-990.
- [157] McCall AA, Swan EE, Borenstein JT, Sewell WF, Kujawa SG and McKenna MJ. Drug delivery for treatment of inner ear disease: current state of knowledge. *Ear Hear* 2010; 31: 156-165.
- [158] Borenstein JT. Intracochlear drug delivery systems. *Expert Opin Drug Deliv* 2011; 8: 1161-1174.
- [159] Rivera T, Sanz L, Camarero G and Varela-Nieto I. Drug delivery to the inner ear: strategies and their therapeutic implications for sensorineural hearing loss. *Curr Drug Deliv* 2012; 9: 231-242.
- [160] Peppi M, Marie A, Belline C and Borenstein JT. Intracochlear drug delivery systems: a novel approach whose time has come. *Expert Opin Drug Deliv* 2018; 15: 319-324.
- [161] Liu SS and Yang R. Inner ear drug delivery for sensorineural hearing loss: current challenges and opportunities. *Front Neurosci* 2022; 16: 867453.
- [162] Fernandez R, Harrop-Jones A, Wang X, Dellamary L, LeBel C and Piu F. The sustained-exposure dexamethasone formulation OTO-104 offers effective protection against cisplatin-induced hearing loss. *Audiol Neurootol* 2016; 21: 22-29.
- [163] Rolland V, Meyer F, Guitton MJ, Bussi eres R, Philippon D, Bairati I, Leclerc M and C ot e M. A randomized controlled trial to test the efficacy of trans-tympanic injections of a sodium thiosulfate gel to prevent cisplatin-induced ototox-

Hearing loss during chemotherapy

- icity in patients with head and neck cancer. *J Otolaryngol Head Neck Surg* 2019; 48: 4.
- [164] Naples JG, Ruckenstein MJ, Singh J, Cox BC and Li D. Intratympanic diltiazem-chitosan hydrogel as an otoprotectant against cisplatin-induced ototoxicity in a mouse model. *Otol Neurotol* 2020; 41: 115-122.
- [165] Martin-Saldana S, Palao-Suay R, Trinidad A, Aguilar MR, Ramirez-Camacho R and San Roman J. Otoprotective properties of 6alpha-methylprednisolone-loaded nanoparticles against cisplatin: in vitro and in vivo correlation. *Nanomedicine* 2016; 12: 965-976.
- [166] Cai H, Liang Z, Huang W, Wen L and Chen G. Engineering PLGA nano-based systems through understanding the influence of nanoparticle properties and cell-penetrating peptides for cochlear drug delivery. *Int J Pharm* 2017; 532: 55-65.
- [167] Plontke SK, Biegner T, Kammerer B, Delabar U and Salt AN. Dexamethasone concentration gradients along scala tympani after application to the round window membrane. *Otol Neurotol* 2008; 29: 401-406.
- [168] Salt AN and Plontke SK. Principles of local drug delivery to the inner ear. *Audiol Neurootol* 2009; 14: 350-360.
- [169] Salt AN and Plontke SK. Pharmacokinetic principles in the inner ear: influence of drug properties on intratympanic applications. *Hear Res* 2018; 368: 28-40.
- [170] Rybak LP and Kelly T. Ototoxicity: bioprotective mechanisms. *Curr Opin Otolaryngol Head Neck Surg* 2003; 11: 328-333.
- [171] Treskes M, Nijtmans LG, Fichtinger-Schepman AM and van der Vijgh WJ. Effects of the modulating agent WR2721 and its main metabolites on the formation and stability of cisplatin-DNA adducts in vitro in comparison to the effects of thiosulphate and diethyldithiocarbamate. *Biochem Pharmacol* 1992; 43: 1013-1019.
- [172] Dickey DT, Wu YJ, Muldoon LL and Neuwelt EA. Protection against cisplatin-induced toxicities by N-acetylcysteine and sodium thiosulfate as assessed at the molecular, cellular, and in vivo levels. *J Pharmacol Exp Ther* 2005; 314: 1052-1058.
- [173] Campbell K, Claussen A, Meech R, Verhulst S, Fox D and Hughes L. D-methionine (D-met) significantly rescues noise-induced hearing loss: timing studies. *Hear Res* 2011; 282: 138-144.
- [174] Campbell KC, Rehemtulla A, Sunkara P, Hamstra D, Buhnerkempe M and Ross B. Oral D-methionine protects against cisplatin-induced hearing loss in humans: phase 2 randomized clinical trial in India. *Int J Audiol* 2022; 61: 621-631.
- [175] Marina N, Chang KW, Malogolowkin M, London WB, Frazier AL, Womer RB, Rescorla F, Billmire DF, Davis MM, Perlman EJ, Giller R, Lauer SJ and Olson TA; Children's Oncology Group. Amifostine does not protect against the ototoxicity of high-dose cisplatin combined with etoposide and bleomycin in pediatric germ-cell tumors: a children's oncology Group study. *Cancer* 2005; 104: 841-847.
- [176] Mukherjea D, Dhukhwa A, Sapra A, Bhandari P, Woolford K, Franke J, Ramkumar V and Rybak L. Strategies to reduce the risk of platinum containing antineoplastic drug-induced ototoxicity. *Expert Opin Drug Metab Toxicol* 2020; 16: 965-982.
- [177] Muldoon LL, Pagel MA, Kroll RA, Brummett RE, Doolittle ND, Zuhowski EG, Egorin MJ and Neuwelt EA. Delayed administration of sodium thiosulfate in animal models reduces platinum ototoxicity without reduction of antitumor activity. *Clin Cancer Res* 2000; 6: 309-315.
- [178] Brock PR, Maibach R, Childs M, Rajput K, Roebuck D, Sullivan MJ, Laithier V, Ronghe M, Dall'Igna P, Hiyama E, Brichard B, Skeen J, Mateos ME, Capra M, Rangaswami AA, Ansari M, Rechnitzer C, Veal GJ, Covezzoli A, Brugieres L, Perilongo G, Czauderna P, Morland B and Neuwelt EA. Sodium thiosulfate for protection from cisplatin-induced hearing loss. *N Engl J Med* 2018; 378: 2376-2385.
- [179] Freyer DR, Chen L, Krailo MD, Knight K, Villaluna D, Bliss B, Pollock BH, Ramdas J, Lange B, Van Hoff D, VanSoelen ML, Wiernikowski J, Neuwelt EA and Sung L. Effects of sodium thiosulfate versus observation on development of cisplatin-induced hearing loss in children with cancer (ACCL0431): a multicentre, randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* 2017; 18: 63-74.
- [180] Harned TM, Kalous O, Neuwelt A, Loera J, Ji L, Iovine P, Sposto R, Neuwelt EA and Reynolds CP. Sodium thiosulfate administered six hours after cisplatin does not compromise antineuroblastoma activity. *Clin Cancer Res* 2008; 14: 533-540.
- [181] Wu YJ, Muldoon LL and Neuwelt EA. The chemoprotective agent N-acetylcysteine blocks cisplatin-induced apoptosis through caspase signaling pathway. *J Pharmacol Exp Ther* 2005; 312: 424-431.
- [182] van den Berg JH, Beijnen JH, Balm AJ and Schellens JH. Future opportunities in preventing cisplatin induced ototoxicity. *Cancer Treat Rev* 2006; 32: 390-397.
- [183] Ezerina D, Takano Y, Hanaoka K, Urano Y and Dick TP. N-acetyl cysteine functions as a fast-acting antioxidant by triggering intracellular H(2)S and sulfane sulfur production. *Cell Chem Biol* 2018; 25: 447-459, e444.
- [184] Yoo J, Hamilton SJ, Angel D, Fung K, Franklin J, Parnes LS, Lewis D, Venkatesan V and Win-

Hearing loss during chemotherapy

- quist E. Cisplatin otoprotection using transtympanic L-N-acetylcysteine: a pilot randomized study in head and neck cancer patients. *Laryngoscope* 2014; 124: E87-94.
- [185] Wang W, Chen E, Ding X, Lu P, Chen J, Ma P and Lu L. N-acetylcysteine protect inner hair cells from cisplatin by alleviated cellular oxidative stress and apoptosis. *Toxicol In Vitro* 2022; 81: 105354.
- [186] Thomas Dickey D, Muldoon LL, Kraemer DF and Neuwelt EA. Protection against cisplatin-induced ototoxicity by N-acetylcysteine in a rat model. *Hear Res* 2004; 193: 25-30.
- [187] Choe WT, Chinosornvatana N and Chang KW. Prevention of cisplatin ototoxicity using transtympanic N-acetylcysteine and lactate. *Otol Neurotol* 2004; 25: 910-915.
- [188] Orgel E, Knight KR, Chi YY, Malvar J, Rushing T, Mena V, Eisenberg LS, Rassekh SR, Ross CJD, Scott EN, Neely M, Neuwelt EA, Muldoon LL and Freyer DR. Intravenous N-acetylcysteine to prevent cisplatin-induced hearing loss in children: a nonrandomized controlled phase I trial. *Clin Cancer Res* 2023; 29: 2410-2418.
- [189] Campbell KC, Rybak LP, Meech RP and Hughes L. D-methionine provides excellent protection from cisplatin ototoxicity in the rat. *Hear Res* 1996; 102: 90-98.
- [190] Korver KD, Rybak LP, Whitworth C and Campbell KM. Round window application of D-methionine provides complete cisplatin otoprotection. *Otolaryngol Head Neck Surg* 2002; 126: 683-689.
- [191] Rybak LP, Mukherjea D and Ramkumar V. Mechanisms of cisplatin-induced ototoxicity and prevention. *Semin Hear* 2019; 40: 197-204.
- [192] Alberts DS, Speicher LA, Krutzsch M, Wymer J, Capizzi RL, Conlon J, Barrett A and Aickin M. WR-1065, the active metabolite of amifostine (Ethyol), does not inhibit the cytotoxic effects of a broad range of standard anticancer drugs against human ovarian and breast cancer cells. *Eur J Cancer* 1996; 32A Suppl 4: S17-20.
- [193] Koukourakis MI. Amifostine in clinical oncology: current use and future applications. *Anti-cancer Drugs* 2002; 13: 181-209.
- [194] Church MW, Blakley BW, Burgio DL and Gupta AK. WR-2721 (Amifostine) ameliorates cisplatin-induced hearing loss but causes neurotoxicity in hamsters: dose-dependent effects. *J Assoc Res Otolaryngol* 2004; 5: 227-237.
- [195] Hyppolito MA, de Oliveira AA, Lessa RM and Rossato M. Amifostine otoprotection to cisplatin ototoxicity: a guinea pig study using otoacoustic emission distortion products (DPOEA) and scanning electron microscopy. *Braz J Otorhinolaryngol* 2005; 71: 268-273.
- [196] Kaltenbach JA, Church MW, Blakley BW, McCaslin DL and Burgio DL. Comparison of five agents in protecting the cochlea against the ototoxic effects of cisplatin in the hamster. *Otolaryngol Head Neck Surg* 1997; 117: 493-500.
- [197] Fisher MJ, Lange BJ, Needle MN, Janss AJ, Shu HK, Adamson PC and Phillips PC. Amifostine for children with medulloblastoma treated with cisplatin-based chemotherapy. *Pediatr Blood Cancer* 2004; 43: 780-784.
- [198] Gurney JG, Bass JK, Onar-Thomas A, Huang J, Chintagumpala M, Bouffet E, Hassall T, Gururangan S, Heath JA, Kellie S, Cohn R, Fisher MJ, Panandiker AP, Merchant TE, Srinivasan A, Wetmore C, Qaddoumi I, Stewart CF, Armstrong GT, Broniscer A and Gajjar A. Evaluation of amifostine for protection against cisplatin-induced serious hearing loss in children treated for average-risk or high-risk medulloblastoma. *Neuro Oncol* 2014; 16: 848-855.
- [199] van As JW, van den Berg H and van Dalen EC. Medical interventions for the prevention of platinum-induced hearing loss in children with cancer. *Cochrane Database Syst Rev* 2014; CD009219.
- [200] Duval M and Daniel SJ. Meta-analysis of the efficacy of amifostine in the prevention of cisplatin ototoxicity. *J Otolaryngol Head Neck Surg* 2012; 41: 309-315.
- [201] Freyer DR, Sung L and Reaman GH. Prevention of hearing loss in children receiving cisplatin chemotherapy. *J Clin Oncol* 2009; 27: 317-318; author reply 318-319.
- [202] Sifuentes-Franco S, Sanchez-Macias DC, Carrillo-Ibarra S, Rivera-Valdes JJ, Zuniga LY and Sanchez-Lopez VA. Antioxidant and anti-inflammatory effects of coenzyme Q10 supplementation on infectious diseases. *Healthcare (Basel)* 2022; 10: 487.
- [203] Astolfi L, Simoni E, Valente F, Ghiselli S, Hatzopoulos S, Chicca M and Martini A. Coenzyme Q10 plus multivitamin treatment prevents cisplatin ototoxicity in rats. *PLoS One* 2016; 11: e0162106.
- [204] Scasso F, Sprio AE, Canobbio L, Scanarotti C, Manini G, Berta GN and Bassi AM. Dietary supplementation of coenzyme Q10 plus multivitamins to hamper the ROS mediated cisplatin ototoxicity in humans: a pilot study. *Heliyon* 2017; 3: e00251.
- [205] Martin-Saldana S, Palao-Suay R, Aguilar MR, Ramirez-Camacho R and San Roman J. Polymeric nanoparticles loaded with dexamethasone or alpha-tocopheryl succinate to prevent cisplatin-induced ototoxicity. *Acta Biomater* 2017; 53: 199-210.
- [206] Fernandez K, Spielbauer KK, Rusheen A, Wang L, Baker TG, Eyles S and Cunningham LL. Lovastatin protects against cisplatin-induced

Hearing loss during chemotherapy

- hearing loss in mice. *Hear Res* 2020; 389: 107905.
- [207] Fernandez KA, Allen P, Campbell M, Page B, Townes T, Li CM, Cheng H, Garrett J, Mulquin M, Clements A, Mulford D, Ortiz C, Brewer C, Dubno JR, Newlands S, Schmitt NC and Cunningham LL. Atorvastatin is associated with reduced cisplatin-induced hearing loss. *J Clin Invest* 2021; 131: e142616.
- [208] Hazlitt RA, Teitz T, Bonga JD, Fang J, Diao S, Iconaru L, Yang L, Goktug AN, Currier DG, Chen T, Rankovic Z, Min J and Zuo J. Development of second-generation CDK2 inhibitors for the prevention of cisplatin-induced hearing loss. *J Med Chem* 2018; 61: 7700-7709.
- [209] Ingersoll MA, Malloy EA, Caster LE, Holland EM, Xu Z, Zallocchi M, Currier D, Liu H, He DZZ, Min J, Chen T, Zuo J and Teitz T. BRAF inhibition protects against hearing loss in mice. *Sci Adv* 2020; 6: eabd0561.
- [210] Bowers WJ, Chen X, Guo H, Frisina DR, Federoff HJ and Frisina RD. Neurotrophin-3 transduction attenuates cisplatin spiral ganglion neuron ototoxicity in the cochlea. *Mol Ther* 2002; 6: 12-18.
- [211] Chen FQ, Schacht J and Sha SH. Aminoglycoside-induced histone deacetylation and hair cell death in the mouse cochlea. *J Neurochem* 2009; 108: 1226-1236.
- [212] Drottar M, Liberman MC, Ratan RR and Robertson DW. The histone deacetylase inhibitor sodium butyrate protects against cisplatin-induced hearing loss in guinea pigs. *Laryngoscope* 2006; 116: 292-296.
- [213] Layman WS and Zuo J. Epigenetic regulation in the inner ear and its potential roles in development, protection, and regeneration. *Front Cell Neurosci* 2014; 8: 446.
- [214] Layman WS, Williams DM, Dearman JA, Saucedo MA and Zuo J. Histone deacetylase inhibition protects hearing against acute ototoxicity by activating the Nf-kappaB pathway. *Cell Death Discov* 2015; 1: 15012.
- [215] Wang J, Tian KY, Fang Y, Chang HM, Han YN and Chen FQ. Sulforaphane attenuates cisplatin-induced hearing loss by inhibiting histone deacetylase expression. *Int J Immunopathol Pharmacol* 2021; 35: 20587384211034086.
- [216] He Y, Zheng Z, Liu C, Li W, Zhao L, Nie G and Li H. Inhibiting DNA methylation alleviates cisplatin-induced hearing loss by decreasing oxidative stress-induced mitochondria-dependent apoptosis via the LRP1-PI3K/AKT pathway. *Acta Pharm Sin B* 2022; 12: 1305-1321.
- [217] Wang B, Zhu X, Kim Y, Li J, Huang S, Saleem S, Li RC, Xu Y, Dore S and Cao W. Histone deacetylase inhibition activates transcription factor Nrf2 and protects against cerebral ischemic damage. *Free Radic Biol Med* 2012; 52: 928-936.
- [218] Kubo E, Chhunchha B, Singh P, Sasaki H and Singh DP. Sulforaphane reactivates cellular antioxidant defense by inducing Nrf2/ARE/Prdx6 activity during aging and oxidative stress. *Sci Rep* 2017; 7: 14130.
- [219] Zhang W, Xiong H, Pang J, Su Z, Lai L, Lin H, Jian B, He W, Yang H and Zheng Y. Nrf2 activation protects auditory hair cells from cisplatin-induced ototoxicity independent on mitochondrial ROS production. *Toxicol Lett* 2020; 331: 1-10.
- [220] Mu YR, Zou SY, Li M, Ding YY, Huang X, He ZH and Kong WJ. Role and mechanism of FOXG1-related epigenetic modifications in cisplatin-induced hair cell damage. *Front Mol Neurosci* 2023; 16: 1064579.
- [221] Park JS, Kim SW, Park K, Choung YH, Jou I and Park SM. Pravastatin attenuates noise-induced cochlear injury in mice. *Neuroscience* 2012; 208: 123-132.
- [222] Brand Y, Setz C, Levano S, Listyo A, Chavez E, Pak K, Sung M, Radojevic V, Ryan AF and Bodmer D. Simvastatin protects auditory hair cells from gentamicin-induced toxicity and activates Akt signaling in vitro. *BMC Neurosci* 2011; 12: 114.
- [223] Syka J, Ouda L, Nachtigal P, Solichova D and Semecky V. Atorvastatin slows down the deterioration of inner ear function with age in mice. *Neurosci Lett* 2007; 411: 112-116.
- [224] Whitton DS. Statins and hearing. *Hear Res* 2022; 425: 108453.
- [225] Hati S, Zallocchi M, Hazlitt R, Li Y, Vijayakumar S, Min J, Rankovic Z, Lovas S and Zuo J. AZD5438-PROTAC: A selective CDK2 degrader that protects against cisplatin- and noise-induced hearing loss. *Eur J Med Chem* 2021; 226: 113849.
- [226] Osakabe N, Modafferi S, Ontario ML, Rampulla F, Zimbone V, Migliore MR, Fritsch T, Abdelhameed AS, Maiolino L, Lupo G, Anfuso CD, Genovese E, Monzani D, Wenzel U, Calabrese EJ, Vabulas RM and Calabrese V. Polyphenols in inner ear neurobiology, health and disease: from bench to clinics. *Medicina (Kaunas)* 2023; 59: 2045.
- [227] Lee JS, Kang SU, Hwang HS, Pyun JH, Choung YH and Kim CH. Epicatechin protects the auditory organ by attenuating cisplatin-induced ototoxicity through inhibition of ERK. *Toxicol Lett* 2010; 199: 308-316.
- [228] Kim CH, Kang SU, Pyun J, Lee MH, Hwang HS and Lee H. Epicatechin protects auditory cells against cisplatin-induced death. *Apoptosis* 2008; 13: 1184-1194.
- [229] Yumusakhuyulu AC, Yazici M, Sari M, Binnetoglu A, Kosemihal E, Akdas F, Sirvanci S, Yuksel M, Uneri C and Tutkun A. Protective role of resveratrol against cisplatin induced ototoxicity in

Hearing loss during chemotherapy

- guinea pigs. *Int J Pediatr Otorhinolaryngol* 2012; 76: 404-408.
- [230] Olgun Y, Kirkim G, Kolatan E, Kiray M, Bagriyanik A, Olgun A, Kizmazoglu DC, Ellidokuz H, Serbetcioglu B, Altun Z, Aktas S, Yilmaz O and Guneri EA. Friend or foe? Effect of oral resveratrol on cisplatin ototoxicity. *Laryngoscope* 2014; 124: 760-766.
- [231] Lee CH, Kim KW, Lee SM and Kim SY. Dose-dependent effects of resveratrol on cisplatin-induced hearing loss. *Int J Mol Sci* 2020; 22: 13.
- [232] Fetoni AR, Eramo SL, Paciello F, Rolesi R, Podda MV, Troiani D and Paludetti G. Curcuma longa (curcumin) decreases in vivo cisplatin-induced ototoxicity through heme oxygenase-1 induction. *Otol Neurotol* 2014; 35: e169-177.
- [233] Soyalic H, Gevrek F, Koc S, Avcu M, Metin M and Aladag I. Intraperitoneal curcumin and vitamin E combination for the treatment of cisplatin-induced ototoxicity in rats. *Int J Pediatr Otorhinolaryngol* 2016; 89: 173-178.
- [234] Bahadori F and Demiray M. A realistic view on “the essential medicinal chemistry of curcumin”. *ACS Med Chem Lett* 2017; 8: 893-896.
- [235] Nelson KM, Dahlin JL, Bisson J, Graham J, Pauli GF and Walters MA. The essential medicinal chemistry of curcumin. *J Med Chem* 2017; 60: 1620-1637.
- [236] Fried LE and Arbiser JL. Honokiol, a multifunctional antiangiogenic and antitumor agent. *Antioxid Redox Signal* 2009; 11: 1139-1148.
- [237] Liu Y, Chen L, He X, Fan L, Yang G, Chen X, Lin X, Du L, Li Z, Ye H, Mao Y, Zhao X and Wei Y. Enhancement of therapeutic effectiveness by combining liposomal honokiol with cisplatin in ovarian carcinoma. *Int J Gynecol Cancer* 2008; 18: 652-659.
- [238] Cheng N, Xia T, Han Y, He QJ, Zhao R and Ma JR. Synergistic antitumor effects of liposomal honokiol combined with cisplatin in colon cancer models. *Oncol Lett* 2011; 2: 957-962.
- [239] Maruyama Y, Kuribara H, Morita M, Yuzurihara M and Weintraub ST. Identification of magnolol and honokiol as anxiolytic agents in extracts of saiboku-to, an oriental herbal medicine. *J Nat Prod* 1998; 61: 135-138.
- [240] Liou KT, Shen YC, Chen CF, Tsao CM and Tsai SK. Honokiol protects rat brain from focal cerebral ischemia-reperfusion injury by inhibiting neutrophil infiltration and reactive oxygen species production. *Brain Res* 2003; 992: 159-166.
- [241] Lin YR, Chen HH, Ko CH and Chan MH. Differential inhibitory effects of honokiol and magnolol on excitatory amino acid-evoked cation signals and NMDA-induced seizures. *Neuropharmacology* 2005; 49: 542-550.
- [242] Lo YC, Teng CM, Chen CF, Chen CC and Hong CY. Magnolol and honokiol isolated from *Magnolia officinalis* protect rat heart mitochondria against lipid peroxidation. *Biochem Pharmacol* 1994; 47: 549-553.
- [243] Pillai VB, Samant S, Sundaresan NR, Raghuraman H, Kim G, Bonner MY, Arbiser JL, Walker DL, Jones DP, Gius D and Gupta MP. Honokiol blocks and reverses cardiac hypertrophy in mice by activating mitochondrial Sirt3. *Nat Commun* 2015; 6: 6656.
- [244] Tsai SK, Huang SS and Hong CY. Myocardial protective effect of honokiol: an active component in *Magnolia officinalis*. *Planta Med* 1996; 62: 503-506.
- [245] Chiang CK, Sheu ML, Lin YW, Wu CT, Yang CC, Chen MW, Hung KY, Wu KD and Liu SH. Honokiol ameliorates renal fibrosis by inhibiting extracellular matrix and pro-inflammatory factors in vivo and in vitro. *Br J Pharmacol* 2011; 163: 586-597.
- [246] Chiu JH, Ho CT, Wei YH, Lui WY and Hong CY. In vitro and in vivo protective effect of honokiol on rat liver from peroxidative injury. *Life Sci* 1997; 61: 1961-1971.
- [247] Tan X, Zhou Y, Agarwal A, Lim M, Xu Y, Zhu Y, O'Brien J, Tran E, Zheng J, Gius D and Richter CP. Systemic application of honokiol prevents cisplatin ototoxicity without compromising its antitumor effect. *Am J Cancer Res* 2020; 10: 4416-4434.
- [248] Hall JA, Dominy JE, Lee Y and Puigserver P. The sirtuin family's role in aging and age-associated pathologies. *J Clin Invest* 2013; 123: 973-979.
- [249] Vassilopoulos A, Fritz KS, Petersen DR and Gius D. The human sirtuin family: evolutionary divergences and functions. *Hum Genomics* 2011; 5: 485-496.
- [250] Kelly G. A review of the sirtuin system, its clinical implications, and the potential role of dietary activators like resveratrol: part 1. *Altern Med Rev* 2010; 15: 245-263.
- [251] Kelly GS. A review of the sirtuin system, its clinical implications, and the potential role of dietary activators like resveratrol: part 2. *Altern Med Rev* 2010; 15: 313-328.
- [252] Brown KD, Maqsood S, Huang JY, Pan Y, Harkcom W, Li W, Sauve A, Verdin E and Jaffrey SR. Activation of SIRT3 by the NAD(+) precursor nicotinamide riboside protects from noise-induced hearing loss. *Cell Metab* 2014; 20: 1059-1068.
- [253] Patel S, Shah L, Dang N, Tan X, Almudevar A and White PM. SIRT3 promotes auditory function in young adult FVB/nJ mice but is dispensable for hearing recovery after noise exposure. *PLoS One* 2020; 15: e0235491.

Hearing loss during chemotherapy

- [254] Quan Y, Xia L, Shao J, Yin S, Cheng CY, Xia W and Gao WQ. Adjudin protects rodent cochlear hair cells against gentamicin ototoxicity via the SIRT3-ROS pathway. *Sci Rep* 2015; 5: 8181.
- [255] Someya S, Yu W, Hallows WC, Xu J, Vann JM, Leeuwenburgh C, Tanokura M, Denu JM and Prolla TA. Sirt3 mediates reduction of oxidative damage and prevention of age-related hearing loss under caloric restriction. *Cell* 2010; 143: 802-812.
- [256] Someya S and Prolla TA. Mitochondrial oxidative damage and apoptosis in age-related hearing loss. *Mech Ageing Dev* 2010; 131: 480-486.
- [257] Pang J, Xiong H, Ou Y, Yang H, Xu Y, Chen S, Lai L, Ye Y, Su Z, Lin H, Huang Q, Xu X and Zheng Y. SIRT1 protects cochlear hair cell and delays age-related hearing loss via autophagy. *Neurobiol Aging* 2019; 80: 127-137.
- [258] Zendedel E, Butler AE, Atkin SL and Sahebkar A. Impact of curcumin on sirtuins: a review. *J Cell Biochem* 2018; 119: 10291-10300.
- [259] de Boer VC, de Goffau MC, Arts IC, Hollman PC and Keijer J. SIRT1 stimulation by polyphenols is affected by their stability and metabolism. *Mech Ageing Dev* 2006; 127: 618-627.
- [260] Huang KJ, Kuo CH, Chen SH, Lin CY and Lee YR. Honokiol inhibits in vitro and in vivo growth of oral squamous cell carcinoma through induction of apoptosis, cell cycle arrest and autophagy. *J Cell Mol Med* 2018; 22: 1894-1908.
- [261] Jiang QQ, Fan LY, Yang GL, Guo WH, Hou WL, Chen LJ and Wei YQ. Improved therapeutic effectiveness by combining liposomal honokiol with cisplatin in lung cancer model. *BMC Cancer* 2008; 8: 242.
- [262] Lin JW, Chen JT, Hong CY, Lin YL, Wang KT, Yao CJ, Lai GM and Chen RM. Honokiol traverses the blood-brain barrier and induces apoptosis of neuroblastoma cells via an intrinsic bax-mitochondrion-cytochrome c-caspase protease pathway. *Neuro Oncol* 2012; 14: 302-314.
- [263] Wang X, Duan X, Yang G, Zhang X, Deng L, Zheng H, Deng C, Wen J, Wang N, Peng C, Zhao X, Wei Y and Chen L. Honokiol crosses BBB and BCSFB, and inhibits brain tumor growth in rat 9L intracerebral gliosarcoma model and human U251 xenograft glioma model. *PLoS One* 2011; 6: e18490.
- [264] Enaru B, Socaci S, Farcas A, Socaciu C, Danciu C, Stanila A and Diaconeasa Z. Novel delivery systems of polyphenols and their potential health benefits. *Pharmaceuticals (Basel)* 2021; 14: 946.
- [265] Li S, Li L, Chen J, Fan Y, Wang C, Du Y, Guo C, Chen F and Li W. Liposomal honokiol inhibits glioblastoma growth through regulating macrophage polarization. *Ann Transl Med* 2021; 9: 1644.
- [266] Editorial O. Erratum to liposomal honokiol inhibits glioblastoma growth through regulating macrophage polarization. *Ann Transl Med* 2023; 11: 402.
- [267] Mai NNS, Otsuka Y, Kawano Y and Hanawa T. Preparation and characterization of solid dispersions composed of curcumin, hydroxypropyl cellulose and/or sodium dodecyl sulfate by grinding with vibrational ball milling. *Pharmaceuticals (Basel)* 2020; 13: 383.
- [268] Ha ES, Choi DH, Baek IH, Park H and Kim MS. Enhanced oral bioavailability of resveratrol by using neutralized eudragit e solid dispersion prepared via spray drying. *Antioxidants (Basel)* 2021; 10: 90.
- [269] Wu D, Zhou J, Creyer MN, Yim W, Chen Z, Messersmith PB and Jokerst JV. Phenolic-enabled nanotechnology: versatile particle engineering for biomedicine. *Chem Soc Rev* 2021; 50: 4432-4483.