

Erratum

Lycorine hydrochloride inhibits melanoma cell proliferation, migration and invasion via down-regulating p21^{Cip1/WAF1}: Am J Cancer Res. 2021; 11(4): 1391-1409

Shaomin Shi^{1,2,3,4*}, Chongyang Li^{3,4*}, Yanli Zhang¹, Chaowei Deng^{3,4}, Mengqin Tan^{3,4}, Guangzhao Pan^{3,4}, Juan Du¹, Yacong Ji¹, Qian Li¹, Hanghua Liang^{3,4}, Wei Liu¹, Leiyang Guo¹, Gaichao Zhao^{3,4}, Yaling Liu¹, Hongjuan Cui^{3,4}

¹Department of Dermatology, The Third Hospital of Hebei Medical University, Shijiazhuang 050000, Hebei, China; ²Department of Dermatology, The Fifth Hospital of Shijiazhuang, Shijiazhuang 050000, Hebei, China; ³State Key Laboratory of Silkworm Genome Biology, Southwest University, Chongqing 400715, China; ⁴Cancer Center, Medical Research Institute, Southwest University, Chongqing 400715, China. *Equal contributors.

Received September 8, 2024; Accepted September 13, 2024; Epub September 25, 2024; Published September 30, 2024


In this article, we found that some data in **Figure 1** were wrongly presented. We corrected the figure accordingly. This correction does not change the interpretation or conclusions of the article. We apologize for any inconvenience caused.

The correct **Figure 1** is shown below.

Address correspondence to: Yaling Liu, Department of Dermatology, The Third Hospital of Hebei

Medical University, No. 139, Ziqiang Road, Shijiazhuang 050000, Hebei, China. Tel: +86-0311-88602167; E-mail: yzling_liu@126.com; Hongjuan Cui, State Key Laboratory of Silkworm Genome Biology, Southwest University, No. 2, Tiansheng Road, Beibei 400715, Chongqing, China. Tel: +86-023-68251713; E-mail: hongjuan.cui@gmail.com

Anti-melanoma effect of lycorine hydrochloride



Anti-melanoma effect of lycorine hydrochloride




Figure 1. Anti-proliferative effect of LH on melanoma cells. A. A375 and MV3 cells were treated with LH (0, 10, 20 and 40 μM) for 48 h and proliferation of the cells was observed. Cell percentage in 0 μM group is regarded as 100%. Scale bar, 100 μm. B. Under LH (0, 10, 20, and 40 μM) treatment, cell viability was measured by MTT assay on days 1, 3, 5 and 7. C. Percentage of Edu positive cells after LH (20 μM) treatment for 48 h. Scale bar, 50 μm. D. Cell cycle distribution of the cells after LH (20 μM) treatment for 48 h. E, F. The protein levels of CDK1, CDK2 and Cyclin E1 in LH treated cells were measured by western blotting. Mean ± SD; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.