

Erratum

Ambroxol enhances anti-cancer effect of microtubule-stabilizing drug to lung carcinoma through blocking autophagic flux in lysosome-dependent way: Am J Cancer Res. 2017; 7(12): 2406-2421

Xiulei Zhang¹, Qinyue Chen¹, Meiyu Chen², Xiaoqing Ren¹, Xiaofei Wang¹, Jianghui Qian¹, Yali Sun¹, Xianyi Sha¹

¹Key Laboratory of Smart Drug Delivery, School of Pharmacy, Fudan University, Shanghai, China; ²Department of Pharmacology, School of Pharmacy, Fudan University, Shanghai, China

Received June 21, 2025; Accepted August 25, 2025; Epub November 25, 2025; Published November 30, 2025

The authors regret that the DAPI and TUNEL staining images of group PTX 3 mg/kg + Ax 100 mg/kg in **Figure 6A** were provided incorrectly and inadvertently in the previous process. In this work, the result demonstrated Ax potentiated anti-tumor effect of PTX to A549 subcutaneous tumor with no accompanying toxicity, and higher dosage of Ax exhibited better performance. The corrections made in this corrigendum do not affect the original conclusions. We sincerely apologize for any inconvenience or misunderstanding that these errors may have

caused. We thank the editors and readers for understanding.

The corrected **Figure 6** is shown below.

Address correspondence to: Xianyi Sha, Key Laboratory of Smart Drug Delivery, School of Pharmacy, Fudan University, No. 826 Zhangheng Road, Shanghai, China. Tel: +86-021-51980072; Fax: +86-021-51980072; E-mail: shaxy@fudan.edu.cn




Figure 6. TUNEL and LC3 distribution assay of tumor tissues. A. TUNEL staining (green) of the subcutaneous tumor tissues of each group were performed to show apoptosis cells. Cell nucleus were stained with DAPI (blue). Microscopy 200× magnification. B. LC3 distribution in the subcutaneous tumor tissues of each group. Microscopy 400× magnification. The lower-right images were the enlarged ones.