

Erratum

Livin promotes tumor progression through YAP activation in ovarian cancer: Am J Cancer Res. 2020; 10(10): 3179-3193

Jianhua Gao, Wei Han, Yanan He, Jun Zhou, Jintian Miao, Guangmei Zhang

Department of Obstetrics and Gynecology, The First Affiliated Hospital of Harbin Medical University, Harbin 150001, Heilongjiang, China

Received June 4, 2025; Accepted June 6, 2025; Epub August 25, 2025; Published August 30, 2025

In reviewing our submission, we identified errors in the presentation of **Figures 2E** and **8C**. Specifically, the images in **Figure 2E** were misassembled due to an oversight in removing overlay images during figure preparation. Additionally, the image of the animal in the shYAP/TAZ-Vector group was mistakenly misused. Importantly, the descriptions of the results and the figure legends remain unchanged, and we confirm that these corrections do not affect the overall conclusions of the article. We sincerely apologize for any confusion or inconvenience this may have caused. The corrected **Figures 2** and **8** are as follows.

Address correspondence to: Guangmei Zhang, Department of Obstetrics and Gynecology, The First Affiliated Hospital of Harbin Medical University, No. 23 Post Street, Harbin 150001, Heilongjiang, China.
E-mail: guangmeizhang2014@126.com

Livin promotes ovarian cancer progression





Figure 2. Livin overexpression promotes the proliferation and migration of ovarian cancer cells. A. The cell viability of A2780 and SKOV3 cells was evaluated by CCK-8. B. The colony formation of A2780 and SKOV3 cells transfected with vector and Livin. C, D. The invasive ability of A2780 and SKOV3 cells was measured by Transwell assay. E. The migration potential of A2780 and SKOV3 cells was evaluated by the wound healing test. Scale bars: 200 μ m. *P < 0.05, **P < 0.01.






Figure 8. Livin promotes tumorigenesis and metastasis of ovarian cancer through YAP/TAZ. A. Quantitative analysis of the tumor volume at the indicated time and quantitative analysis of the tumor weight. B. HE, Ki-67 and TUNEL immunohistological staining of tumors from different groups. Scale bars: 100 μ m. C. *In vivo* bioluminescent imaging of nude mice at day 28 after ovarian cancer cell implantation and HE staining for lung occupancy of different groups. Scale bars: 250 μ m. **P < 0.01, ***P < 0.001.