Erratum

HMGA2 regulates CD44 expression to promote gastric cancer cell motility and sphere formation: Am J Cancer Res. 2017; 7(2): 260-274


Junying Sun^{1*}, Baocun Sun^{1,2,3*}, Dongwang Zhu⁴, Xiulan Zhao^{1,2}, Yanhui Zhang³, Xueyi Dong¹, Na Che^{1,2}, Jing Li^{1,2}, Fang Liu¹, Nan Zhao^{1,2}, Danfang Zhang^{1,2}, Tieju Liu^{1,2}, Xian Lin^{1,2}

¹Department of Pathology, Tianjin Medical University, Tianjin 300070, China; ²Department of Pathology, Tianjin General Hospital, Tianjin Medical University, Tianjin 300052, China; ³Department of Pathology, Tianjin Cancer Hospital, Tianjin Medical University, Tianjin 300060, China; ⁴Department of Prosthodontics, Affiliated Stomatological Hospital, Tianjin Medical University, Tianjin 300070, China: *Equal contributors.

Received June 26, 2025; Accepted July 1, 2025; Epub August 25, 2025; Published August 30, 2025

In this article, there are two misrepresented images in Figures 4A and 5A. Hence, we would like to publish this erratum to displace the wrong figures and reflect changes. The authors have confirmed that the errors associated with this figure did not have any significant impact on either the results or the conclusions reported in this study. The authors apologize for any inconvenience or misunderstanding that these errors may have caused. The corrected Figures 4 and 5 are shown below.

Address correspondence to: Baocun Sun, Department of Pathology and General Hospital and Cancer Hospital of Tianjin Medical University, No. 22 Qixiangtai Road, Heping District, Tianjin 300070, China. Tel: +86-13602111192; Fax: +86-22-83336813; E-mail: baocunsun@aliyun.com







Figure 4. HMGA2 increases the clonogenicity and proliferation of GC cells. A. A colony formation assay was performed to analyze colony formation ability. The scale bar represents 200 μ m. B. MTT assays of HMGA2-upregulated MKN74 and MKN28 cells and HMGA2-downregulated MGC803 and MKN45 cells. C. Western blot showing that HMGA2 overexpression increased the expression of c-Myc, Sox2 and Oct4, whereas HMGA2 knockdown decreased the expression of c-Myc, Sox2 and Oct4. D. Luciferase reporter assays showed that HMGA2 promotes c-Myc gene expression. All experiments were repeated three times; * indicates P<0.05.






Figure 5. HMGA2 regulates the migratory capacity of GC cells and increases the expression of EMT-related markers in GC cells. A. Wound healing assay (the scale bar represents 200 μm). Quantitative analysis showed a significant difference at 48 h for MGC803 cells and a significant difference at 72 h for MKN45, MKN28 and MKN74 cells. B. Western blot showing that HMGA2 overexpression increased the expression of Snail and β-catenin, while HMGA2 knockdown decreased the expression of Snail and β-catenin. All experiments were repeated three times; * indicates *P*<0.05.

В