

Erratum

Disulfiram combined with chemoimmunotherapy potentiates pancreatic cancer treatment efficacy through the activation of cGAS-STING signaling pathway via suppressing PARP1 expression: Am J Cancer Res. 2023; 13(5): 2055-2065

Shenan Huang^{1*}, Peiyi Xie^{2*}, Xiaomei Huang^{1*}, Zhimeng Chen^{3*}, Jinhuan Yang⁴, Jian Wang⁵, Cong Liu³, Hui Li^{6,7}, Binghai Zhou³

¹Department of Gastroenterology, The Second Affiliated Hospital of Nanchang University, Nanchang 330006, Jiangxi, P. R. China; ²Department of Liver Surgery and Transplantation, Liver Cancer Institute, Zhongshan Hospital, Fudan University, Shanghai 200032, P. R. China; ³Hepato-Biliary-Pancreatic Surgery Division, Department of General Surgery, The Second Affiliated Hospital of Nanchang University, Nanchang 330006, Jiangxi, P. R. China; ⁴Department of Hepatobiliary Surgery, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou 325035, Zhejiang, P. R. China; ⁵Department of Hepato-Biliary-Pancreatic Surgery, Pingxiang People's Hospital, Pingxiang 337099, Jiangxi, P. R. China; ⁶Department of Liver Surgery and Transplantation, Liver Cancer Institute, Zhongshan Hospital, Fudan University, Key Laboratory of Carcinogenesis and Cancer Invasion, Ministry of Education, Shanghai 200032, P. R. China; ⁷Shanghai Medical College and Zhongshan Hospital Immunotherapy Translational Research Center, Shanghai 200031, P. R. China. *Equal contributors.

Received March 10, 2026; Accepted April 20, 2026; Epub August 25, 2026; Published August 30, 2026

In this article, some errors have been found in **Figure 3**. Therefore, we provide the correct version to displace the wrong figure and reflect changes. We sincerely apologize for this oversight and any confusion it may have caused. The corrected **Figure 3** are shown below.

Address correspondence to: Dr. Hui Li, Department of Liver Surgery and Transplantation, Liver Cancer Institute, Zhongshan Hospital, Fudan University, Key

Laboratory of Carcinogenesis and Cancer Invasion, Ministry of Education, Shanghai 200032, P. R. China. E-mail: li.hui1@zs-hospital.sh.cn; Dr. Binghai Zhou, Hepato-Biliary-Pancreatic Surgery Division, Department of General Surgery, The Second Affiliated Hospital of Nanchang University, Nanchang 330006, Jiangxi, P. R. China. E-mail: 18888291342@163.com

Disulfiram combined with chemoimmunotherapy potentiates PDAC via PARP1




Figure 3. DSF activated antitumor immune response via the cGAS-STING pathway. A. The protein levels of γH2AX in PANC-1 cells treated by chemotherapy and/or DSF (10 μM) were detected by western blot. B. Quantification of the staining results of cytosolic dsDNA in PANC-1 cells. C. S172 phosphorylation of TBK1 in PANC-1 cells treated by chemotherapy and/or DSF (10 μM). D. cGAS protein expression was detected by western blot in PANC-1 cells transfected with shNC or shRNA targeting cGAS. E and F. The mRNA level of IFNα and IFNβ in PANC-1 cells transfected with shNC or shcGAS after being treated with DSF plus chemotherapy or STING agonist (diABZI) was determined by qRT-PCR. G. S172 phosphorylation of TBK1 and total TBK1 protein levels were detected in PANC-1 cells with or without chemotherapy plus DSF treatment (**P < 0.01, ***P < 0.001, NS indicates no significance).