

Review Article

The Oct4 protein: more than a magic stemness marker

Dana Zeineddine^{1*}, Aya Abou Hammoud^{1,2,3*}, Mohamad Mortada¹, H          ^{2,3}

¹Lebanese University, Beyrouth, Liban; ²Univ. Bordeaux, CIRID, UMR 5164, F-33 000 Bordeaux, France; ³CNRS, CIRID, UMR 5164, F-33 000 Bordeaux, France. *Equal contributors.

Received July 22, 2014; Accepted August 7, 2014; Epub September 5, 2014; Published September 15, 2014

Abstract: The Oct4 protein, encoded by the *Pou5f1* gene was the very first master gene, discovered 25 years ago, to be absolutely required for the stemness properties of murine and primate embryonic stem cells. This transcription factor, which has also been shown to be essential for somatic cell reprogramming, displays various functions depending upon its level of expression and has been quoted as a "rheostat" gene. Oct4 protein is in complexes with many different partners and its activity depends upon fine post-translational modifications. This review aims at revisiting some properties of this protein, which has not yet delivered all its potentialities.

Keywords: Oct4/ *Pou5f1*, embryonic stem cells, pluripotency, cell lineages, cancer stem cells

Oct-4: a member of the POU family

Oct-4, a transcription factor also known as Oct-3, Oct-3/4, Otf3 or NF-A3, is encoded by the *Pou5f1* gene (located on chromosome 6 in human and 17 in mouse) and belongs to the POU (Pit, Oct, Unc) family of DNA binding-proteins. These proteins regulate the expression of target genes by binding to the octamer motif ATGCAAAT within their promoter or enhancer regions [1, 2]. Oct4, whose expression is associated with pluripotent properties of stem cells, is an essential factor controlling early stages of mammalian embryogenesis [3-5].

Oct4 expression in early embryos

Maternal murine Oct-4 mRNA and protein (352 AA), deposited in the oocyte, are inherited by the zygote and are necessary for development until the 4-cell stage. Proteins are present at low levels at these early stages of murine embryogenesis. Transcription of zygotic *Pou5f1* gene is activated at the 4- to 8-cell stage. Consequently, high level of nuclear Oct-4 protein is detected in all blastomeres until morula stage. Upon blastocyst formation, Oct4 expression remains high in the inner cell mass (ICM) and is not expressed in the trophectoderm (TE). After implantation of the mouse embryo, transient upregulation of Oct4 in a group of cells of

the ICM triggers their differentiation into primitive endoderm (hypoblast) cells. Subsequently, Oct4 expression is down-regulated in these cells [6-8]. During gastrulation, Oct4 is down-regulated and, after day 8 of gestation, it is confined to primordial germ cells [7, 9, 10]. *In vitro*, Oct4 is highly expressed in undifferentiated embryonic stem (ES) cells, embryonic carcinoma (EC) cells and embryonic germ cells. Upon differentiation of these cells induced by Leukemia Inhibitory Factor (LIF) withdrawal or in the presence of retinoic acid, Oct4 expression is down-regulated with different kinetic [11-13].

In humans, Oct-3/4, unlike in mice, encodes 2 isoforms that are generated by alternative splicing of *Pou5f1* mRNA [14, 15]. These isoforms, Oct4-1A and Oct4-1B (360 and 265 amino acids, respectively), of which the 225 amino acids at the C-termini are identical, differ in sequence at their N termini [15]. Critical amounts of human Oct4-1A are required to sustain stem cell self-renewal and it has been shown that Oct4-1B is not related to stemness. In humans, Oct4 mRNA is present throughout all stages from the unfertilized oocyte to the uncompact morula [16-18]. These stages display a variable expression pattern of Oct4 mRNA, between individual blastomeres of the same cell stage with only cytoplasmic localization of Oct4 proteins. No




Figure 1. LIF signaling and post-translational modifications of some stemness transcription factors. The maintenance of self-renewal and pluripotency in mESC is controlled by extrinsic signaling pathway (LIF) and intrinsic self-renewal factors (eg: Oct4, Sox2 and Nanog). LIF binds to its heterodimeric receptor which leads to the activation of receptor associated Janus Kinases (JAKs) which phosphorylate receptor docking sites and Stat3 on tyrosine 705 (Tyr705). Then, dimers of Phospho-Stat3 translocate to the nucleus and activate the transcription of target genes that are essential for ES cell self-renewal (eg: Klf4, Klf5 which induces expression of Mras GTPase). The activated Stat3 leads also to expression of its own repressor like Socs3, which serves as a negative feedback signal. Phosphorylation, Ubiquitination, SUMOylation and Acetylation of Oct4, Sox2 and Nanog control protein activity, expression, and stability, which result in modulation of ESC self-renewal activity. S stands for “stabilization” of the protein. Arrow indicates “activation” while the line indicates “repression”.

Oct4 proteins are found in the nucleus during these stages [11]. During compaction, the expression of Oct4 protein becomes ubiquitous and abundant in the nuclei of all blastomeres of the morula. In blastocysts, Oct4 transcripts and proteins are present in the ICM [18]. As in murine model, Oct4 is present in human embryonic stem (hES) cells [19], human embryonic carcinoma cells [20] as well as in human embryonic germ cells [21].

Oct4 is a gatekeeper for ES cell pluripotency

Pluripotent embryonic stem cell identity is governed by a core of transcription factors involving Oct4 that acts as a key regulator of pluripotent cells across mammalian species. Oct-4

-/- embryos developed to blastocysts only composed of trophoblast cells and they are dead at the time of implantation [4, 22]. In mouse ES cells, Oct4 knockdown results in an up-regulation of genes associated with endoderm differentiation such as *Gata6* and *a-feto-protein* and those associated with trophoblast differentiation such as *Cdx2* [23, 24].

Recent investigation by elegant disruption of maternal or zygotic Oct4 show also the crucial role of this gene for lineage priming *in vivo* [25]. In addition, Oct-4 is required to maintain the pluripotency in hES cells where it acts as a repressor of the gene encoding for hCG (human chorionic gonadotropin), a placental marker in hES cells. Down-regulation of Oct4 in hES cells, was coincident with a significant increase in transcription of genes associated with trophoblast and endoderm lineages [26, 27]. Furthermore, Oct4, Sox2 and Nanog cooperate to maintain the self-renewal and pluripotency of mouse and human ES cells. These three transcription factors, together bound to the promoters of their own genes [28, 29], allow their activation. In addition, they activate the transcription of genes involved in maintaining ES cell self-renewal. *Dppa4*, *Tdgf1*, *Oct4*, *Nanog*, and *Lefty2* are positively regulated by Oct4, Sox2 and Nanog and preferentially expressed in ES cells [28]. On the other hand Oct4, Sox2 and Nanog co-occupy the promoters of genes that have key roles in differentiation and development and promote their inactivation [30].

The intersection between the transcriptional core (Oct4, Sox2 and Nanog) and LIF signaling

In addition to Oct4, Sox2 and Nanog, the LIF/STAT3 signaling pathway is also known to be critical for the maintenance of pluripotency in murine ES (mES) cells while it seems dispensable in hES cells, despite the expression and the functional activation of the LIF/STAT3 in these cells [31]. Since these first observations, it has been demonstrated that hES cells are closer to murine Epiblast stem cells (primed cells) than to naïve mES cells [32, 33]. In addi-

Oct4: a rheostat effector gene




Figure 2. Protein sequence alignment of mammalian OCT4 proteins. Protein sequences of the different mammalian proteins, retrieved from GenBank, have been aligned with the MultiAlign software [81]. The conserved ERK, PKA and SUMOylation sites are indicated.

tion, recent studies have demonstrated a key role of LIF to induce the reversion of primed to naïve states in the human model [12, 34, 35]. An important challenge of this research field is to characterize the molecular mechanisms of ES cell plasticity.

In mES cells, LIF (Leukemia inhibitory factor), via activation of JAK1/ STAT3 pathway, is an essential cytokine that prevents differentiation. Stat3 activates the expression of a significant number of pluripotency-related genes in ES cells [12, 36-39]. Furthermore, some of the Stat3 target genes are also regulated by Oct4. For example, Oct4 and Stat3 directly bind to the promoter region of *Eed* gene (Embryonic ectoderm development) which encodes for a core component of Polycomb repressive complex 2 [40, 41]. In addition, *Klf4* (Krüppel-like transcription factor 4), a direct downstream target of Oct4 and Stat3, is required for ES cell self-renewal and maintenance of pluripotency [42-44]. These data revealed that the cooperation between the intrinsic Oct4 network and extrinsic LIF pathway is crucial to regulate ES cell self-renewal and pluripotency. However Oct4 and LIF's targets are regulated in different ways upon cell differentiation triggered by LIF withdrawal: indeed, Oct4 mRNA and protein remains at high level for few days upon LIF withdrawal. This is in contrast with many of the LIF 's targets (named *Pluri* genes like *Mras*, *Ceacam1* and 2, *Irak3*, *Esrrb*) whose mRNA expression decreased on the first day of LIF withdrawal concomitantly with cell differentiation [12, 13, 39]. It has also recently been shown that dysregulated *Mras* expression, a small GTPase of ras family, lead to an alteration in Oct4 expression, indicating a potential link between this small GTPase and Oct4 regulation [45], see **Figure 1**. More generally, how the *Pluri* gene cluster, which encodes various types of protein regulators, modulates Master gene activity remains to be resolved.

Oct4 and its dosage effect on the cell fate

Oct4 dosage is important in the determination of the mES cell fate. Depending upon Oct4 expression level, mES cells maintain their pluripotency or differentiate towards trophoblast (low or no Oct4 expression) or primitive endodermal and mesodermal (high Oct4) lineages [23, 46, 47]. This rheostat behavior of gene, which has also been shown for Sox2 [48], revealed that these genes exerts a dose-

dependent action. In addition, it has been well documented that neuronal differentiation of ES cells, under the serum-free culture condition, is enhanced because of the sustained overexpression of Oct4 in ES cells [49]. Moreover, the involvement of Oct4 in the mesendoderm differentiation and cardiac commitment of ES cells was also proved. Indeed, transient increase in Oct4 expression upon TGF β induction, in undifferentiated ES cells and in the epiblast of mouse embryos leads to establishment of cardiogenic lineages [46]. Also, by different approaches it was shown that, depending upon its expression level, Oct4 could form different protein complexes with members of the SOX family: the Sox2/ Oct4 complex, which binds to a canonical binding site, induces expression of genes involved in the maintenance of pluripotency. In contrast, an increase in Oct4 or Sox17 expression level leads to a switch of partners, Sox2 being replaced by Sox17. This complex then binds to a different compressed DNA motif at promoter of genes involved in primitive endoderm and mesendoderm differentiation [46, 47, 50, 51]. Whether the Oct4 protein, along with Sox2 or Sox17, displays various functions as a complex, depending upon their post-translational modification, remains unknown.

Regulation of Oct4 stability at the post-translational level

The mechanisms through which Oct4 protein activity is regulated are largely unclear. Potential phosphorylated or sumoylated conserved residues, present among the mammalian Oct4 proteins, are shown in **Figure 2**. Oct4 can be phosphorylated by protein kinase A and/or ERK MAPK at a highly conserved residue, Ser 229 (murine) or Ser 236 (human) within the POU DNA-binding homeodomain, (see **Figure 2**) [52]. Phosphorylation at this Ser residue sterically hinders both DNA binding and homodimer assembly [53, 54]. Furthermore, ubiquitination of Oct4 (shown for both murine and human proteins) is also a post-translational modification that dramatically reduces its transcriptional activity. During ubiquitination process, the E3 ligase interacts directly with target proteins and promotes the Ubiquitin transfer. WWP2, an E3 ubiquitin ligase that specifically interacts with Oct4 through its Tryptophan-based WW domains, has been identified in murine and human embryonic

stem cells. WWP2 promotes the ubiquitination of Oct4 and its degradation both *in vivo* and *in vitro* [55-59]. Additionally, MEK/ERK signaling, activated upon ES cell differentiation, is involved in the negative regulation of human Oct4-IA protein, through its phosphorylation at Ser111 which alters stability and its subcellular localization [60]. Oct4 is also a target for small ubiquitin-related modifier (SUMO)-1 that occurs at a highly conserved K among the mammalian proteins (K118 in mice and K123 in primates), see **Figure 2**. This conserved lysine is located at the end of the amino-terminal transactivation domain and next to the DNA-binding domain. Sumoylation of Oct4, which does not alter its subnuclear localization, enhances its stability, DNA binding and transactivation functions [61, 62]. Whether sumoylation of Oct4 is involved in its increased level, coupled with its function in primitive endoderm differentiation, remains to be established. Interestingly, it has also been shown that SUMOylation of Oct4 and Sox2 regulated Nanog in an opposing manner: SUMOylation of Oct4 enhanced Nanog expression, while SUMOylated Sox2 inhibited its expression. Moreover, SUMOylation of Oct4 by Pias2 or Sox2 by Pias3 impaired the interaction between Oct4 and Sox2 [63]. All these findings reveal that the post-translational modifications of Oct4, act as a regulatory signal to control its activity and stability in a wide variety of cellular processes. This also applies to other stem cell factors, such as Sox-2 and Nanog (**Figure 1**).

Oct-4 and tumorigenicity

It is well documented that overexpression of Oct4, Sox2 and Nanog, together or separately, led to tumorigenicity, tumor metastasis, and even distant recurrence after chemoradiotherapy in different types of cancer [64-66]. High expression of Oct4 was detected in Prostate [67] and Breast cancer stem cells [68] and in the tumor initiating cells in a p53^{-/-} tumor mice model [69]. Oct4 has a critical role in the survival of these tumor cells. Generally, these transcription factors are more frequently overexpressed in poorly differentiated tumors (compared to well differentiated tumors) and expression level of these stemness-involved factors decreases with the differentiation of cells [70-73]. There are probably conserved molecular mechanisms which could explain dedifferentiation of somatic cells, as observed

in cancers, and somatic cell reprogramming [74-76]. So far, reprogramming has not yet been demonstrated to occur, normally, in particular situation in the adult body. However, we could hypothesize that this potential normal process, if it exists, should be constantly under strict control in adult body [for example, co-expression of Oct4, Nanog, Sox2 and myc, one of the *in vitro* reprogramming cocktails, [75, 77-79] should never occur in normal differentiated adult cells]. We could then hypothesize that cancer formation is the result of uncontrolled reprogramming, both involving Oct4 and many other stemness genes [80].

Conclusion

Oct4, first discovered and characterized 25 years ago [1, 5], is not only a key stemness marker but it is also involved in lineages specification and it is a cell ressetor involved in somatic cell reprogramming *in vitro* [33, 74]. Oct4 is also reexpressed in different types of cancer stem cells, which are tumor cell clusters at the origin of chemotherapy tumor resistance and recidive of cancers. Though, the precise understanding of the molecular mechanisms of Oct4 regulation and particularly of its switch ON and OFF in tissues, depending upon microenvironment, is a challenge in fundamental and applied research fields, for regenerative medicine and cancer therapy.

Acknowledgements

This work was funded by CNRS, University of Bordeaux, the Region Aquitaine, the SIRIC BRIO of Bordeaux and the SFR TransBiomed. AAH is funded by a co-tutelle PhD fellowship of the Hariri Foundation, Liban. We thank Claire Saucourt, a previous PhD student of the group, for her contribution to **Figure 1**.

Disclosure of conflict of interest

None.

Address correspondence to: Dr. Hélène Boeuf, UMR-CNRS-5164-CIRID, Université de Bordeaux, Bat. 1B, RdC, BP14, 146 rue Léo Saignat, 33076 Bordeaux, France. E-mail: helene.boeuf@u-bordeaux2.fr

References

- [1] Scholer HR, Balling R, Hatzopoulos AK, Suzuki N, Gruss P. Octamer binding proteins confer transcriptional activity in early mouse embryogenesis. *EMBO J* 1989; 8: 2551-7.

- [2] Yeom YI, Ha HS, Balling R, Scholer HR, Artzt K. Structure, expression and chromosomal location of the Oct-4 gene. *Mech Dev* 1991; 35: 171-9.
- [3] Boiani M, Scholer HR. Regulatory networks in embryo-derived pluripotent stem cells. *Nat Rev Mol Cell Biol* 2005; 6: 872-84.
- [4] Nichols J, Zevnik B, Anastasiadis K, Niwa H, Klewe-Nebenius D, Chambers I, Schöler H, Smith A. Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. *Cell* 1998; 95: 379-91.
- [5] Scholer HR. Octamania: the POU factors in murine development. *Trends Genet* 1991; 7: 323-9.
- [6] Palmieri SL, Peter W, Hess H, Schöler HR. Oct-4 transcription factor is differentially expressed in the mouse embryo during establishment of the first two extraembryonic cell lineages involved in implantation. *Dev Biol* 1994; 166: 259-67.
- [7] Pesce M, Scholer HR. Oct-4: gatekeeper in the beginnings of mammalian development. *Stem Cells* 2001; 19: 271-8.
- [8] Jerabek S, Merino F, Schöler HR, Cojocaru V. OCT4: Dynamic DNA binding pioneers stem cell pluripotency. *Biochim Biophys Acta* 2014; 1839: 138-154.
- [9] Pan GJ, Chang ZY, Scholer HR, Pei D. Stem cell pluripotency and transcription factor Oct4. *Cell Res* 2002; 12: 321-9.
- [10] Kehler J, Tolkunova E, Koschorz B, Pesce M, Gentile L, Boiani M, Lomeli H, Nagy A, McLaughlin KJ, Schöler HR, Tomilin A. Oct4 is required for primordial germ cell survival. *EMBO Rep* 2004; 5: 1078-83.
- [11] Cauffman G, Van de Velde H, Liebaers I, Van Steirteghem A. Oct-4 mRNA and protein expression during human preimplantation development. *Mol Hum Reprod* 2005; 11: 173-81.
- [12] Trouillas M, Saucourt C, Guillotin B, Gauthereau X, Taupin JL, Moreau JF, Boeuf H. The LIF cytokine: towards adulthood. *Eur Cytokine Netw* 2009; 20: 51-62.
- [13] Trouillas M, Saucourt C, Guillotin B, Gauthereau X, Ding L, Buchholz F, Doss MX, Sachinidis A, Hescheler J, Hummel O, Huebner N, Kolde R, Vilo J, Schulz H, Boeuf H. Three LIF-dependent signatures and gene clusters with atypical expression profiles, identified by transcriptome studies in mouse ES cells and early derivatives. *BMC Genomics* 2009; 10: 73.
- [14] Takeda J, Seino S, Bell GI. Human Oct3 gene family: cDNA sequences, alternative splicing, gene organization, chromosomal location, and expression at low levels in adult tissues. *Nucleic Acids Res* 1992; 20: 4613-20.
- [15] Cauffman G, Liebaers I, Van Steirteghem A, Van de Velde H. POU5F1 isoforms show different expression patterns in human embryonic stem cells and preimplantation embryos. *Stem Cells* 2006; 24: 2685-91.
- [16] Verlinsky Y, Morozov G, Verlinsky O, Koukharenko V, Rechitsky S, Goltsman E, Ivakhnenko V, Gindilis V, Strom CM, Kuliev A. Isolation of cDNA libraries from individual human preimplantation embryos. *Mol Hum Reprod* 1998; 4: 571-5.
- [17] Abdel-Rahman B, Fiddler M, Rappolee D, Pergament E. Expression of transcription regulating genes in human preimplantation embryos. *Hum Reprod* 1995; 10: 2787-92.
- [18] Hansis C, Grifo JA, Krey LC. Oct-4 expression in inner cell mass and trophectoderm of human blastocysts. *Mol Hum Reprod* 2000; 6: 999-1004.
- [19] Reubinoff BE, Pera MF, Fong CY, Trounson A, Bongso A. Embryonic stem cell lines from human blastocysts: somatic differentiation in vitro. *Nat Biotechnol* 2000; 18: 399-404.
- [20] Pera MF, Herszfeld D. Differentiation of human pluripotent teratocarcinoma stem cells induced by bone morphogenetic protein-2. *Reprod Fertil Dev* 1998; 10: 551-5.
- [21] Goto T, Adjaye J, Rodeck CH, Monk M. Identification of genes expressed in human primordial germ cells at the time of entry of the female germ line into meiosis. *Mol Hum Reprod* 1999; 5: 851-60.
- [22] Chambers I, Tomlinson SR. The transcriptional foundation of pluripotency. *Development* 2009; 136: 2311-22.
- [23] Niwa H, Miyazaki J, Smith AG. Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nat Genet* 2000; 24: 372-6.
- [24] Niwa H, Toyooka Y, Shimosato D, Strumpf D, Takahashi K, Yagi R, Rossant J. Interaction between Oct3/4 and Cdx2 determines trophectoderm differentiation. *Cell* 2005; 123: 917-29.
- [25] Le Bin GC, Muñoz-Descalzo S, Kurowski A, Leitch H, Lou X, Mansfield W, Etienne-Dumeau C, Grabole N, Mulas C, Niwa H, Hadjantonakis AK, Nichols J. Oct4 is required for lineage priming in the developing inner cell mass of the mouse blastocyst. *Development* 2014; 141: 1001-10.
- [26] Hay DC, Sutherland L, Clark J, Burdon T. Oct-4 knockdown induces similar patterns of endoderm and trophoblast differentiation markers in human and mouse embryonic stem cells. *Stem Cells* 2004; 22: 225-35.
- [27] Zafarana G, Avery SR, Avery K, Moore HD, Andrews PW. Specific knockdown of OCT4 in human embryonic stem cells by inducible short hairpin RNA interference. *Stem Cells* 2009; 27: 776-82.
- [28] Boyer LA, Lee TI, Cole MF, Johnstone SE, Levine SS, Zucker JP, Guenther MG, Kumar RM, Mur-

- ray HL, Jenner RG, Gifford DK, Melton DA, Jaenisch R, Young RA. Core transcriptional regulatory circuitry in human embryonic stem cells. *Cell* 2005; 122: 947-56.
- [29] Niwa H. How is pluripotency determined and maintained? *Development* 2007; 134: 635-46.
- [30] Christophersen NS, Helin K. Epigenetic control of embryonic stem cell fate. *J Exp Med* 2010; 207: 2287-95.
- [31] Dahéron L, Opitz SL, Zaehres H, Lensch MW, Andrews PW, Itskovitz-Eldor J, Daley GQ. LIF/STAT3 signaling fails to maintain self-renewal of human embryonic stem cells. *Stem Cells* 2004; 22: 770-8.
- [32] Pera MF, Trounson AO. Human embryonic stem cells: prospects for development. *Development* 2004; 131: 5515-25.
- [33] Pera MF. Defining pluripotency. *Nat Methods* 2010; 7: 885-7.
- [34] Gafni O, Weinberger L, Mansour AA, Manor YS, Chomsky E, Ben-Yosef D, Kalma Y, Viukov S, Maza I, Zviran A, Rais Y, Shipony Z, Mukamel Z, Krupalnik V, Zerbib M, Geula S, Caspi I, Schneir D, Shwartz T, Gilad S, Amann-Zalcenstein D, Benjamin S, Amit I, Tanay A, Massarwa R, Novershtern N, Hanna JH. Derivation of novel human ground state naive pluripotent stem cells. *Nature* 2013; 504: 282-6.
- [35] Hanna J, Cheng AW, Saha K, Kim J, Lengner CJ, Soldner F, Cassady JP, Muffat J, Carey BW, Jaenisch R. Human embryonic stem cells with biological and epigenetic characteristics similar to those of mouse ESCs. *Proc Natl Acad Sci U S A* 2010; 107: 9222-7.
- [36] Boeuf H, Hauss C, Graeve FD, Baran N, Keding-er C. Leukemia inhibitory factor-dependent transcriptional activation in embryonic stem cells. *J Cell Biol* 1997; 138: 1207-17.
- [37] Bourillot PY, Savatier P. Krüppel-like transcription factors and control of pluripotency. *BMC Biol* 2010; 8: 125.
- [38] Bourillot PY, Aksoy I, Schreiber V, Wianny F, Schulz H, Hummel O, Hubner N, Savatier P. Novel STAT3 target genes exert distinct roles in the inhibition of mesoderm and endoderm differentiation in cooperation with Nanog. *Stem Cells* 2009; 27: 1760-71.
- [39] Mathieu ME, Saucourt C, Mournetas V, Gauthereau X, Thézé N, Praloran V, Thiébaud P, Boeuf H. LIF-Dependent Signaling: New Pieces in the Lego. *Stem Cell Rev* 2012; 8: 1-15.
- [40] Ura H, Usuda M, Kinoshita K, Sun C, Mori K, Akagi T, Matsuda T, Koide H, Yokota T. STAT3 and Oct-3/4 control histone modification through induction of Eed in embryonic stem cells. *J Biol Chem* 2008; 283: 9713-23.
- [41] Ura H, Murakami K, Akagi T, Kinoshita K, Yamaguchi S, Masui S, Niwa H, Koide H, Yokota T. Eed/Sox2 regulatory loop controls ES cell self-renewal through histone methylation and acetylation. *EMBO J* 2011; 30: 2190-204.
- [42] Li Y, McClintick J, Zhong L, Edenberg HJ, Yoder MC, Chan RJ. Murine embryonic stem cell differentiation is promoted by SOCS-3 and inhibited by the zinc finger transcription factor Klf4. *Blood* 2005; 105: 635-7.
- [43] Niwa H, Ogawa K, Shimosato D, Adachi K. A parallel circuit of LIF signalling pathways maintains pluripotency of mouse ES cells. *Nature* 2009; 460: 118-22.
- [44] Papapetrou EP, Tomishima MJ, Chambers SM, Mica Y, Reed E, Menon J, Tabar V, Mo Q, Studer L, Sadelain M. Stoichiometric and temporal requirements of Oct4, Sox2, Klf4, and c-Myc expression for efficient human iPSC induction and differentiation. *Proc Natl Acad Sci U S A* 2009; 106: 12759-64.
- [45] Mathieu ME, Faucheux C, Saucourt C, Soulet F, Gauthereau X, Fédou S, Trouillas M, Thézé N, Thiébaud P, Boeuf H. MRAS GTPase is a novel stemness marker that impacts mouse embryonic stem cell plasticity and Xenopus embryonic cell fate. *Development* 2013; 140: 3311-22.
- [46] Zeineddine D, Papadimou E, Chebli K, Gineste M, Liu J, Grey C, Thurig S, Behfar A, Wallace VA, Skerjanc IS, Pucéat M. Oct-3/4 dose dependently regulates specification of embryonic stem cells toward a cardiac lineage and early heart development. *Dev Cell* 2006; 11: 535-46.
- [47] Stefanovic S, Puceat M. Oct-3/4: not just a gatekeeper of pluripotency for embryonic stem cell, a cell fate instructor through a gene dosage effect. *Cell Cycle* 2007; 6: 8-10.
- [48] Kopp JL, Ormsbee BD, Desler M, Rizzino A. Small increases in the level of Sox2 trigger the differentiation of mouse embryonic stem cells. *Stem Cells* 2008; 26: 903-11.
- [49] Shimosaki K, Nakashima K, Niwa H, Taga T. Involvement of Oct3/4 in the enhancement of neuronal differentiation of ES cells in neurogenesis-inducing cultures. *Development* 2003; 130: 2505-12.
- [50] Aksoy I, Jauch R, Chen J, Dyla M, Divakar U, Bogu GK, Teo R, Leng Ng CK, Herath W, Lili S, Hutchins AP, Robson P, Kolatkar PR, Stanton LW. Oct4 switches partnering from Sox2 to Sox17 to reinterpret the enhancer code and specify endoderm. *EMBO J* 2013; 32: 938-53.
- [51] Stefanovic S, Abboud N, Désilets S, Nury D, Cowan C, Pucéat M. Interplay of Oct4 with Sox2 and Sox17: a molecular switch from stem cell pluripotency to specifying a cardiac fate. *J Cell Biol* 2009; 186: 665-73.
- [52] Brumbaugh J, Hou Z, Russell JD, Howden SE, Yu P, Ledvina AR, Coon JJ, Thomson JA. Phos-

- phorylation regulates human OCT4. *Proc Natl Acad Sci U S A* 2012; 109: 7162-8.
- [53] Saxe JP, Tomilin A, Schöler HR, Plath K, Huang J. Post-translational regulation of Oct4 transcriptional activity. *PLoS One* 2009; 4: e4467.
- [54] Swaney DL, Wenger CD, Thomson JA, Coon JJ. Human embryonic stem cell phosphoproteome revealed by electron transfer dissociation tandem mass spectrometry. *Proc Natl Acad Sci U S A* 2009; 106: 995-1000.
- [55] Cai N, Li M, Qu J, Liu G-H, Izpisua Belmonte JC. Post-translational modulation of pluripotency. *J Mol Cell Biol* 2012; 4: 262-5.
- [56] Liao B, Jin Y. Wwp2 mediates Oct4 ubiquitination and its own auto-ubiquitination in a dosage-dependent manner. *Cell Res* 2010; 20: 332-44.
- [57] Liao B, Zhong X, Xu H, Xiao F, Fang Z, Gu J, Chen Y, Zhao Y, Jin Y. Itch, an E3 ligase of Oct4, is required for embryonic stem cell self-renewal and pluripotency induction. *J Cell Physiol* 2013; 228: 1443-51.
- [58] Xu H, Wang W, Li C, Yu H, Yang A, Wang B, Jin Y. WWP2 promotes degradation of transcription factor OCT4 in human embryonic stem cells. *Cell Res* 2009; 19: 561-73.
- [59] Xu HM, Liao B, Zhang QJ, Wang BB, Li H, Zhong XM, Sheng HZ, Zhao YX, Zhao YM, Jin Y. Wwp2, an E3 ubiquitin ligase that targets transcription factor Oct-4 for ubiquitination. *J Biol Chem* 2004; 279: 23495-503.
- [60] Spelat R, Ferro F, Curcio F. Serine 111 phosphorylation regulates OCT4A protein subcellular distribution and degradation. *J Biol Chem* 2012; 287: 38279-88.
- [61] Gill G. SUMO and ubiquitin in the nucleus: different functions, similar mechanisms? *Genes Dev* 2004; 18: 2046-59.
- [62] Wei F, Schöler HR, Atchison ML. Sumoylation of Oct4 enhances its stability, DNA binding, and transactivation. *J Biol Chem* 2007; 282: 21551-60.
- [63] Wu Y, Guo Z, Wu H, Wang X, Yang L, Shi X, Du J, Tang B, Li W, Yang L, Zhang Y. SUMOylation represses Nanog expression via modulating transcription factors Oct4 and Sox2. *PLoS One* 2012; 7: e39606.
- [64] Koo BS, Lee SH, Kim JM, Huang S, Kim SH, Rho YS, Bae WJ, Kang HJ, Kim YS, Moon JH, Lim YC. Oct4 is a critical regulator of stemness in head and neck squamous carcinoma cells. *Oncogene* 2014; [Epub ahead of print].
- [65] Boumahdi S, Driessens G, Lapouge G, Rorive S, Nassar D, Le Mercier M, Delatte B, Caauwe A, Lenglez S, Nkusi E, Brohé S, Salmon I, Dubois C, del Marmol V, Fuks F, Beck B, Blanpain C. SOX2 controls tumour initiation and cancer stem-cell functions in squamous-cell carcinoma. *Nature* 2014; 511: 246-50.
- [66] Abubaker K, Luwor RB, Zhu H, McNally O, Quinn MA, Burns CJ, Thompson EW, Findlay JK, Ahmed N. Inhibition of the JAK2/STAT3 pathway in ovarian cancer results in the loss of cancer stem cell-like characteristics and a reduced tumor burden. *BMC Cancer* 2014; 14: 317.
- [67] Ben-Porath I, Thomson MW, Carey VJ, Ge R, Bell GW, Regev A, Weinberg RA. An embryonic stem cell-like gene expression signature in poorly differentiated aggressive human tumors. *Nat Genet* 2008; 40: 499-507.
- [68] Hu T, Liu S, Breiter DR, Wang F, Tang Y, Sun S. Octamer 4 small interfering RNA results in cancer stem cell-like cell apoptosis. *Cancer Res* 2008; 68: 6533-40.
- [69] Darini CY, Pisani DF, Hofman P, Pedeutour F, Sudaka I, Chomienne C, Dani C, Ladoux A. Self-renewal gene tracking to identify tumour-initiating cells associated with metastatic potential. *Oncogene* 2012; 31: 2438-49.
- [70] Patru C, Romao L, Varlet P, Coulombel L, Raponi E, Cadusseau J, Renault-Mihara F, Thirant C, Leonard N, Berhneim A, Mihaescu-Maingot M, Haiech J, Bièche I, Moura-Neto V, Daumas-Duport C, Junier MP, Chneiweiss H. CD133, CD15/SSEA-1, CD34 or side populations do not resume tumor-initiating properties of long-term cultured cancer stem cells from human malignant glioma-neuronal tumors. *BMC Cancer* 2010; 10: 66.
- [71] Thirant C, Bessette B, Varlet P, Puget S, Cadusseau J, Tavares Sdos R, Studler JM, Silvestre DC, Susini A, Villa C, Miquel C, Bogeas A, Surena AL, Dias-Morais A, Léonard N, Pflumio F, Bièche I, Boussin FD, Sainte-Rose C, Grill J, Daumas-Duport C, Chneiweiss H, Junier MP. Clinical relevance of tumor cells with stem-like properties in pediatric brain tumors. *PLoS One* 2011; 6: e16375.
- [72] Thirant C, Galan-Moya EM, Dubois LG, Pinte S, Chafey P, Broussard C, Varlet P, Devaux B, Soncin F, Gavard J, Junier MP, Chneiweiss H. Differential Proteomic Analysis of Human Glioblastoma and Neural Stem Cells Reveals HDGF as a Novel Angiogenic Secreted Factor. *Stem Cells* 2012; 30: 845-853.
- [73] Thirant C, Gavard J, Junier MP, Chneiweiss H. Critical multiple angiogenic factors secreted by glioblastoma stem-like cells underline the need for combinatorial anti-angiogenic therapeutic strategies. *Proteomics Clin Appl* 2013; 7: 79-90.
- [74] Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 2006; 126: 663-76.
- [75] Carey BW, Markoulaki S, Hanna JH, Faddah DA, Buganim Y, Kim J, Ganz K, Steine EJ, Casady JP, Creighton MP, Welstead GG, Gao Q,

Oct4: a rheostat effector gene

- Jaenisch R. Reprogramming factor stoichiometry influences the epigenetic state and biological properties of induced pluripotent stem cells. *Cell Stem Cell* 2011; 9: 588-98.
- [76] Buganim Y, Faddah DA, Jaenisch R. Mechanisms and models of somatic cell reprogramming. *Nat Rev Genet* 2013; 14: 427-39.
- [77] Okita K, Matsumura Y, Sato Y, Okada A, Morizane A, Okamoto S, Hong H, Nakagawa M, Tanabe K, Tezuka K, Shibata T, Kunisada T, Takahashi M, Takahashi J, Saji H, Yamanaka S. A more efficient method to generate integration-free human iPS cells. *Nat Methods* 2011; 8: 409-12.
- [78] Shi Y, Desponts C, Do JT, Hahm HS, Scholer HR, Ding S. Induction of pluripotent stem cells from mouse embryonic fibroblasts by Oct4 and Klf4 with small-molecule compounds. *Cell Stem Cell* 2008; 3: 568-74.
- [79] Inoue H, Nagata N, Kurokawa H, Yamanaka S. iPS cells: a game changer for future medicine. *EMBO J* 2014; 33: 409-17.
- [80] Ohnishi K, Semi K, Yamamoto T, Shimizu M, Tanaka A, Mitsunaga K, Okita K, Osafune K, Arioka Y, Maeda T, Soejima H, Moriwaki H, Yamanaka S, Woltjen K, Yamada Y. Premature termination of reprogramming in vivo leads to cancer development through altered epigenetic regulation. *Cell* 2014; 156: 663-77.
- [81] Corpet F. Multiple sequence alignment with hierarchical clustering. *Nucleic Acids Res* 1988; 16: 10881-90.