Review Article Integrin signaling aberrations in prostate cancer

Hira Lal Goel, Naved Alam, Isaac N.S. Johnson & Lucia R. Languino

Department of Cancer Biology, Prostate Cancer Discovery and Development Program, Cancer Center, University of Massachusetts Medical School, Worcester, MA 01605, USA.

Received April 14, 2009; accepted April 17, 2009; available online April 20, 2009

Abstract: Integrins are cell surface receptors for extracellular matrix proteins and play a key role in cell survival, proliferation, migration and gene expression. Integrin signaling has been shown to be deregulated in several types of cancer, including prostate cancer. This review is focused on integrin signaling pathways known to be deregulated in prostate cancer and known to promote prostate cancer progression.

Key words: Focal adhesion kinase, PTEN, PI 3-kinase/AKT, Ras/Raf/MAPK, cdc2, survivin, Bcl-2

Introduction

Prostate cancer is a significant burden in western countries and has been predicted to account for more than 28,660 deaths and 186,320 new cases in 2008 [1]. Prostate cancer development proceeds through a series of defined states. These include prostatic intraepithelial neoplasia (PIN); high-grade PIN lesions, which usually develop prior to invasive cancer: androgen-sensitive invasive cancer and an androgen-independent castrationresistant state [2, 3]. The current therapies for prostate cancer involve surgery, androgen ablation, or the blockade of the androgen receptor; however, a significantly high percentage of treated prostate cancers eventually grows, despite either castration levels of androgen or the presence of anti-androgens. For these patients, radiation therapy is the only treatment available. Still, a large number of patients relapse.

Integrins are cell surface receptors for extracellular matrix proteins and play a key role in cell survival, proliferation, migration and gene expression. Integrin signaling has been shown to be deregulated in several types of cancer, including prostate cancer. In prostate cancer, tumor cells have a different surrounding matrix than normal cells; thus changes in the integrin profile may be functionally relevant and contribute to aberrant intracellular signaling [4-8]. Several studies have associated deregulation of integrin expression with the progression of prostate cancer to an advanced stage (**Table 1**) [4, 8-11]. This article reviews the literature on the major signaling pathways activated by integrins and their deregulation in prostate cancer.

Integrin deregulation in prostate cancer

Integrins are heterodimers consisting of α and β subunits. At this time, 24 heterodimers of the integrin family, consisting of 18 α and 8 β subunits, have been described [12, 13], and their ability to activate specific signaling pathways has been investigated [13]. Integrin signaling plays a key role in the alteration of cellular growth and tumor progression through the regulation of gene expression, apoptosis, cell adhesion, proliferation, migration and angiogenesis [14, 15], as well as proteinase expression [16]. Most α and β subunits have been shown to be downregulated in prostate cancer, whereas only α 6, β 1, β 3 and β 6 are upregulated [6]. Among the α subunits, several reports show that α 3, α 4, α 5 and α 7 are downregulated [17, 18]; α 2 and α 6 are aberrantly expressed, whereas there are no reports on the remaining subunits [6]. A unique

Table 1. Deregulated expression of integrin subunits in human prostate cancer and metastasis

Subunit	Adenocarcinoma	Metastasis	References	
α6	unknown	↑	Knox et al., 1994 [11]; Bonkhoff et al., 1993 [19]; Nagle et al., 1995 [20]	
α _{llb} (trun- cated)	↑	unknown	Trikha et al, 1998 [33]	
β1	1	unknown	Murant et al., 1997 [10]; Knox et al., 1994 [11]; Goel et al., 2007 [22]	
βз	1	1	Zheng et al., 1999 [30]	
β6	1	↑	Li and Languino, 2007 [31]	
Down-Regulated				
a3, a4 a5	\downarrow	unknown	Nagle et al., 1994 [18]	
α7	Ļ	unknown	Ren et al., 2007 [17]	
βıc	Ļ	unknown	Fornaro et al., 1996, 1998, 1999 [25-27]; Perli- no et al., 2000 [28]	
β4	\downarrow	unknown	Nagle et al., 1995 [20]; Davis et al., 2001 [23]; Allen et al., 1998 [24]	
Other				
α2	Ļ	↑	Nagle et al., 1994 [18]; Bonkhoff et al., 1993 [19]	

Up-Regulated

This table shows the expression of integrin subunits found to be deregulated in human primary and metastatic prostate cancer.

expression pattern has been shown for $\alpha 2$, which is downregulated in prostate cancer, but upregulated in lymph node metastases as compared to primary lesions [18, 19]. An extensive analysis of $\alpha 6$ expression in prostate cancer shows that $\alpha 6$ expression is either maintained or overexpressed in prostate cancer, and increases in lymph node metastases [11, 19-21].

Among the β subunits, β 1, β 3, and β 6 are upregulated, while β 1C and β 4 are downregulated in human prostate cancer [6, 20, 22-24]. No reports are available for β 5, β 7, and β 8. Five β 1 variant subunits, β 1A, β 1B, β 1C, β 1C-2, and β 1D, generated by alternative splicing, have been described. Two variants, β 1C and β 1A, are shown to be expressed in benign prostatic epithelium. β 1C is expressed at both protein and mRNA levels in benign prostatic epithelial cells, but is markedly downregulated in adenocarcinoma [25-28]. Fornaro et al. show that the expression of β 1C integrin increases p^{27kip1} levels, a cell cycle inhibitor, as well as p^{27kip1} association with cyclin A [26]. In contrast, the findings that the expression of the β 1A integrin variant is upregulated and is necessary for the cell's ability to grow in an



Figure 1. Integrin-dependent signaling pathways. Schematic drawing showing the signal transduction pathways regulated by integrins that control prostate cancer cell survival, proliferation, adhesion, migration, and cytoskeletal organization. For a detailed description of integrin downstream effectors like Rac, cdc42, Src, Cas, Rho and Crk, or cytoskeletal proteins like AFAP-110, talin and paxillin, readers should refer to previous articles [38, 86].

anchorage-independent manner [29], point to the important role that the β 1A integrin may have during prostate cancer progression and will be helpful in formulating new therapeutic strategies.

Upregulation of $\alpha\nu\beta3$, $\alpha\nu\beta6$ and the truncated α llb integrin variant has been described [6]. Zheng et al., using human prostate cancer cells isolated from 16 surgical specimens, show that these cells express $\alpha\nu\beta3$, whereas normal prostate epithelial cells do not [30]. Similarly, $\alpha\nu\beta6$ [31, 32] and the truncated α llb integrin variant [33] are found to be expressed in adenocarcinoma.

The β 1 and β 3 integrin subunits are known to localize in focal contacts and to mediate spreading and cytoskeletal rearrangement in normal cells [12, 13]. However, when we either downregulated or upregulated these subunits by siRNA or ectopic expression analysis, we show that cancer cell spreading is not af-

fected [29, 34]. These results demonstrate that the ability of the β 1 and β 3 subunits to promote cancer progression is independent of cell spreading.

Overall, these findings indicate that the expression of selective integrin subunits is deregulated during prostate cancer progression, and that these subunits are potential diagnostic markers in prostate cancer.

Activation of unique signaling pathways by integrins

The expression of the β 1 and β 3 subunits activates specific signaling pathways and supports distinct cancer cell functions [34, 35]. Analysis of the mechanism by which β 1 may promote tumor growth in vivo, shows that β 1 is uniquely required in cancer cells for the localization, expression and function of insulin-like growth factor type 1 receptor (IGF-IR), which is known to support cancer cell proliferation and surviv-

al [29, 35]. The mechanism proposed for β 1 integrins' control of IGF-IR activity involves β 1 recruiting specific adaptors to the plasma membrane, thus increasing the concentration of specific adaptors proximal to the growth factor receptor [35]. This study provides evidence that the β 1 cytodomain plays an important role in mediating β 1 integrin association with either insulin receptor substrate-1 (IRS-1) or Grb2-associated binder1 (Gab1)/SH2containing protein-tyrosine phosphate 2 (Shp2), downstream effectors of IGF-IR. Specifically, β 1A associates with IRS-1 and β 1C with Gab1/Shp2 [29, 35, 36].

In parallel studies, we have discovered that β3 is uniquely required in cancer cells for increasing cdc2 levels, as well as cdc2 kinase activity. While *β*1 integrin expression does not increase cancer cell motility or cdc2 levels, and appears to predominantly modulate cell proliferation and survival, these effects are specific for β 3. Higher levels of cdc2 result in increased cell migration mediated by the specific association of cdc2 with cyclin B2 and the phosphorylation of caldesmon, a substrate of cdc2. These results show that cdc2 acts as a downstream mediator of the avß3 integrin and promotes cancer cell migration [34]. In conclusion, the β1 and β3 integrins promote activation of selective signaling pathways that support prostate cancer progression.

Integrin downstream effectors

Since integrins lack catalytic activity, they depend on intracellular effector proteins to transduce signals [37, 38]. In this section, we discuss the major signaling effectors that are likely to contribute to prostate cancer progression (**Figure 1** and **Table 2**).

Focal adhesion kinase (FAK)

FAK is a non-receptor tyrosine kinase, which becomes activated upon integrin-extracellular matrix (ECM) interactions and integrin clustering [39, 40]. Upon phosphorylation, FAK interacts with several signaling proteins, including Src kinases, Cas, paxillin and Phosphoinositide 3-Kinase (PI 3-Kinase) [39, 40]. FAK signaling is altered in prostate cancer. In normal prostate, FAK expression is absent or weak in secretory epithelium and is expressed predominantly in the basal layers. Prostate carcinoma shows a greater expression of FAK compared to the secretory layer of normal prostate. FAK expression is further increased in invasive prostate cancer [41, 42].

A well established role for FAK is its ability to regulate cancer cell motility [43]. The expression of dominant negative FAK inhibits the migration of prostate carcinoma cells [44]. In our previous study, we show that the β 3 integrin induces cell migration on vitronectin, which is mediated by FAK [30]. Recently, the role of FAK in cell migration has been confirmed by using an inhibitor of FAK phosphorylation, PF-573,228. This inhibitor fails to inhibit cell growth or to induce apoptosis. In contrast, treatment with PF-573,228 inhibits both chemotactic and haptotactic migration concomitant with the inhibition of focal adhesion turnover [45]. In addition, Dasatinib, an inhibitor of Src family kinases/Abl, blocks FAK and Cas signaling in human prostate cancer cells, resulting in the suppression of invasion, migration and adhesion of prostate cancer cells [46].

Bombesin is shown to stimulate PC-3 cell migration and tyrosine phosphorylation of FAK. In addition, bombesin also increases the association between FAK and the B1, B3 and B5 integrins [47]. Bombesin induces relocalization of FAK in focal contacts, followed by its tyrosine phosphorylation and the formation of actin lamellipodia. FAK inhibitors cause reduced cell motility upon bombesin treatment [48]. FAK is also required for bombesin stimulated activation of RhoA, a GTPase required for cell migration [49]. Another example of the role that FAK plays in cell migration is provided by Sumitomo et al., who use Neutral endopeptidase 24.11 (NEP) [50]. NEP is an enzyme which cleaves neuropeptides such as bombesin and endothelin-1. NEP treatment blocks bombesin and endothelin-stimulated cell migration and FAK phosphorylation. This study suggests that NEP expression results in the formation of a complex containing NEP, Lyn and PI 3-Kinase and this complex competitively blocks FAK/PI 3-Kinase interactions [50]. The FAK/PI 3-Kinase interactions are also shown to promote prostate cancer cell invasion: $\alpha 5\beta 1$ interacts with the PHSRN sequence of fibronectin (FN), which induces FAK phosphorylation and FAK association with PI 3-Kinase, resulting in prostate cancer cell invasion [51]. FAK siRNA, or specific PI 3-Kinase inhibitors, block PHSRN-mediated invasion

Downstream	Expression/Activity	Prostate cancer stage	References
Effectors			
FAK	upregulated expression	invasive cancer and me- tastasis	Rovin et al, 2002 [41]; Tremblay et al, 1996 [42]
MAP kinase	increased kinase activity	androgen-independent state	Bakin et al, 2003 [53]
PTEN	downregulated expression	cancer and metastasis	Schmitz et al, 2007 [61]
AKT	increased kinase activity	cancer with high Gleason score	Sun et al, 2001 [67]; Malik et al, 2002 [68]
Survivin	upregulated expression	PIN, primary tumors and metastasis	Shariat et al, 2004 [71]; Krajewska et al, 2003 [72]; Kishi et al, 2004 [73]
Bcl-2	upregulated expression	PIN, primary tumors and metastasis from recurrent cancer	Colombel et al, 1993 [76]; Zellweger et al, 2005 [77]; Krajewska et al, 1996 [78]

 Table 2. Aberrant Integrin-Dependent Pathways in Prostate Cancer

Signaling proteins and inhibitors of apoptosis known to be regulated by integrins and to affect prostate cancer progression are shown. FAK, Focal adhesion kinase; PTEN, phosphatase and tensin homolog; MAP kinase, mitogen-activated protein kinase.

[51]. Overall, these studies highlight a crucial role for the FAK in prostate cancer cell invasion mediated by integrins.

Ras/Raf/MAP kinase

Mitogen-activated protein (MAP) kinases, the principal effectors of Ras and known downstream effectors of integrins, are major regulators of cell proliferation and cell differentiation [52]. Although Ras and Raf mutations are not common in prostate cancer, it is known that the activation of the Ras/MAP kinase pathway might be sufficient for progression towards the androgen-independent state [53, 54]. A high ERK/p38 activity ratio favors prostate tumor growth and activation of $\alpha 5\beta 1$ integrin is proposed as a determinant of the in vivo growth promoting activity of a high ERK/p38 ratio [55]. Furthermore, inhibition of MAP kinase, using U0126, decreases α6 integrin mRNA levels in androgen-independent prostate cancer cells [56]. Thus, blocking MAP kinase activation provides an important tool to regulate integrin signaling during prostate cancer progression.

PTEN

PTEN, a dual specificity phosphatase, has the ability to dephosphorylate inositol phospholipids such as phosphatidylinositol-3,4,5triphosphate (PIP3) and, as a consequence, negatively regulates AKT activation. By virtue of its ability to inhibit the AKT pathway, PTEN

acts as a tumor suppressor [57]. The Pten gene is frequently deleted or mutated in human cancers and is shown to be involved in the regulation of cell migration on integrin substrates [58]. In 1997, PTEN was cloned from the 10q23 region, a region frequently targeted by loss of heterozygosity in advanced cancer [59, 60]. PTEN alterations are common in prostate cancer. Recently, Schmitz et al. have shown that 23% of patients with first time diagnoses lost PTEN expression, and 59% of patients with lymph node metastasis no longer express PTEN. These findings suggest that loss of PTEN expression is a possible early prognostic marker for prostate cancer metastasis [61].

Overexpression of PTEN inhibits cell migration, whereas antisense to PTEN enhances cell migration. These effects are suggested to be mediated by FAK regulation, since overexpression of FAK partially antagonizes the effects of PTEN. Thus, PTEN phosphatase may function as a tumor suppressor by negatively regulating cell interactions with the ECM, mediated by integrins [58]. PTEN is shown to regulate the adhesion and proliferation of LNCaP-C4-2 prostate cancer cells stimulated by vascular endothelial growth factor [62]. PTEN expression inhibits LNCaP-C4-2 cell migration toward calvaria-conditioned medium, but has no effect on migration toward lung-conditioned medium, and this inhibitory effect is dependent on PTEN lipid phosphatase activity [63]. All these studies suggest that PTEN downregulation contributes to integrin activation of signaling pathways that mediate cancer progression, although the mechanisms underlying this cross-talk remain to be investigated.

PI 3-Kinase/AKT pathway

PI 3-Kinase is a major downstream component of the integrin and growth factor signaling pathways [64, 65]. PI 3-Kinase catalyzes the production of the lipid secondary messenger PIP3 at the cell membrane. PIP3, in turn, contributes to the recruitment and activation of a wide range of downstream targets, including the serine-threonine protein kinase AKT [64]. Several studies show that integrin-mediated activation of PI 3-Kinase plays a crucial role in cancer cell survival, preventing anoikis and promoting cell migration (for review, [37, 66]). AKT1 kinase activity is significantly increased in primary carcinomas of the prostate [67]. AKT activation, assessed by immunohistochemical staining of human prostate cancer biopsies, shows greater intensity in high Gleason grade compared to PIN and all other grades of prostate cancer [68]. Similarly, using protein microarrays, it is shown that prostate cancer progression is associated with increased phosphorylation of AKT [69]. Although AKT promotes several integrin-mediated functions, our studies indicate a predominant role for the PI 3-Kinase/AKT pathway in prostate cancer cell migration [70].

Survivin/Bcl-2

Survivin, an important member of the inhibitor of apoptosis family, is a dual regulator of cell proliferation and cell viability. Survivin is expressed in embryonic and fetal organs, but is undetectable in most differentiated tissues. Survivin is shown to be upregulated in prostate cancer, especially in aggressive forms, such as high grade carcinoma and metastasis [71-73]. We demonstrate that β 1 integrin engagement by FN upregulates the expression of survivin. and increases protection from apoptosis induced by the TNF- α in aggressive prostate cancer cells. The expression of dominant negative survivin counteracts the ability of FN to protect cells from undergoing apoptosis. We also show that the regulation of survivin levels by integrins is mediated by the AKT pathway [74]. It should be noted that in addition to integrin-ECM interactions, IGF/mTOR signaling and anti-androgen therapy are associated with the modulation of survivin levels in prostate cancer [75].

Bcl-2 is another important regulator of cell survival. Bcl-2 expression is restricted to the basal cells in normal and hypertrophic prostate glands, but all epithelial cells in areas of PIN express Bcl-2 [76]. All primary prostatic carcinomas and metastases obtained from hormone-refractory tumors are shown to express Bcl-2 [76-78]. This suggests that they may protect tumor cells from apoptosis induced in response to radiotherapy or chemotherapy. Integrin ligation, specifically by $\alpha 5\beta 1$ and $\alpha v\beta 3$, but not $\alpha v\beta 1$, stimulates Bcl-2 expression via the FAK and PI 3-Kinase pathways [79, 80]. This integrin-mediated regulation of Bcl-2 is also controlled by the activation of Ca2+/calmodulin-dependent protein kinase IV, NF-kappaB and CREB transcription factors [79, 80]. Bcl-2 is also known to suppress anoikis induced by guinazoline based α 1adrenoceptor antagonists in prostate cancer cells [81].

All these recent studies highlight a crucial role for survivin and Bcl-2 in prostate cancer cell survival mediated by integrins.

Conclusions and future studies

The studies reviewed here indicate that designing new diagnostic and therapeutic approaches for prostate cancer, based on inhibitors of integrin functions or of integrin downstream signaling, will prove to be a successful strategy. However, the molecular pathways by which integrins contribute to prostate cancer progression, and in general, the molecular mechanisms that promote this disease remain to be fully investigated. Several areas of research appear under-investigated. Among others, a major effort is needed to study the mechanisms by which integrins are deregulated in prostate cancer and to characterize integrinmediated pathways which support survival of prostate cancer stem cells. Furthermore, new preclinical studies to test the efficacy of integrin inhibitors in prostate cancer are necessary. For this purpose, prostate cancer mouse models, such as the TRAMP mouse or the mouse which carries a conditional Pten deletion in the prostate are useful tools. Future studies will also take advantage of the use of recently developed novel small animal molecular imaging approaches, such as bioluminescence imaging (BLI) [82, 83]. A very innovative study, using BLI in mice that ubiquitously express luciferase (FLASH, firefly luciferase activated systemically in homozygotes), proves that we can increase our ability to detect tumor response to therapeutic agents like siR-NAs [84, 85].

In conclusion, studies aimed at elucidating the mechanisms by which deregulation of integrinmediated signaling pathways occurs in prostate cancer will provide novel therapeutic approaches for this disease.

Acknowledgements

This study was supported by NIH grant RO1 CA-89720 and RO1 CA-109874); DOD (PCRP DAMD PC040221) to LRL and from American Cancer Society (Institutional Research Grant-93-033) to HLG. We would like to thank Heather Ehlers for help during the preparation of this article.

Please address correspondences to: Lucia R. Languino, PhD, Department of Cancer Biology, Prostate Cancer Discovery and Development Program, University of Massachusetts Medical School, Worcester, MA 01605, USA. E-mail: <u>lucia.languino@umassmed.edu</u>

References

- Jemal A, Siegel R, Ward E, Hao Y, Xu J, Murray T, Thun MJ. Cancer statistics, 2008. CA Cancer J Clin 2008; 58:71-96.
- [2] Scher HI, Heller G. Clinical states in prostate cancer: toward a dynamic model of disease progression. Urology 2000; 55:323-327.
- [3] De Marzo AM, Meeker AK, Zha S, Luo J, Nakayama M, Platz EA, Isaacs WB, Nelson WG. Human prostate cancer precursors and pathobiology. Urology 2003; 62:55-62.
- [4] Fornaro M, Manes T, Languino LR. Integrins and prostate cancer metastases. Cancer Metastasis Rev 2001; 20:321-331.
- [5] Demetriou MC, Cress AE. Integrin clipping: a novel adhesion switch? J Cell Biochem 2004; 91:26-35.
- [6] Goel HL, Li J, Kogan S, Languino LR. Integrins in prostate cancer progression. Endocr Relat Cancer 2008; 3:657-664.
- [7] Boudreau N, Bissell MJ. Extracellular matrix signaling: integration of form and function in normal and malignant cells. Curr Opin Cell Biol 1998; 10:640-646.
- [8] Knudsen BS, Miranti CK. The impact of cell adhesion changes on proliferation and survival during prostate cancer development and progression. J Cell Biochem 2006; 99:345-361.

- [9] Edlund M, Sung SY, Chung LW. Modulation of prostate cancer growth in bone microenvironments. J Cell Biochem 2004; 91:686-705.
- [10] Murant SJ, Handley J, Stower M, Reid N, Cussenot O, Maitland NJ. Co-ordinated changes in expression of cell adhesion molecules in prostate cancer. Eur J Cancer 1997; 33:263-271.
- [11] Knox JD, Cress AE, Clark V, Manriquez L, Affinito KS, Dalkin BL, Nagle RB. Differential expression of extracellular matrix molecules and the α 6-integrins in the normal and neoplastic prostate. Am J Pathol 1994; 145:167-174.
- [12] Hynes RO. Integrins: bidirectional, allosteric signaling machines. Cell 2002; 110:673-687.
- [13] Alam N, Goel HL, Zarif MJ, Butterfield JE, Perkins HM, Sansoucy BG, Sawyer TK, Languino LR. The integrin-growth factor receptor duet. J Cell Physiol 2007; 213:649-653.
- [14] Akalu A, Cretu A, Brooks PC. Targeting integrins for the control of tumour angiogenesis. Expert Opin Investig Drugs 2005; 14:1475-1486.
- [15] Felding-Habermann B. Integrin adhesion receptors in tumor metastasis. Clin Exp Metastasis 2003; 20:203-213.
- [16] Munshi HG, Stack MS. Reciprocal interactions between adhesion receptor signaling and MMP regulation. Cancer Metastasis Rev 2006; 25:45-56.
- [17] Ren B, Yu YP, Tseng GC, Wu C, Chen K, Rao UN, Nelson J, Michalopoulos GK, Luo JH. Analysis of integrin α7 mutations in prostate cancer, liver cancer, glioblastoma multiforme, and leiomyosarcoma. J Natl Cancer Inst 2007; 99:868-880.
- [18] Nagle RB, Knox JD, Wolf C, Bowden GT, Cress AE. Adhesion molecules, extracellular matrix, and proteases in prostate carcinoma. J Cell Biochem Suppl 1994; 19:232-237.
- [19] Bonkhoff H, Stein U, Remberger K. Differential expression of α 6 and α 2 very late antigen integrins in the normal, hyperplastic, and neoplastic prostate: simultaneous demonstration of cell surface receptors and their extracellular ligands. Hum Pathol 1993; 24:243-248.
- [20] Nagle RB, Hao J, Knox JD, Dalkin BL, Clark V, Cress AE. Expression of hemidesmosomal and extracellular matrix proteins by normal and malignant human prostate tissue. Am J Pathol 1995; 146:1498-1507.
- [21] Pontes-Junior J, Reis ST, Dall'oglio M, Neves de Oliveira LC, Cury J, Carvalho PA, Ribeiro-Filho LA, Moreira Leite KR, Srougi M. Evaluation of the expression of integrins and cell adhesion molecules through tissue microarray in lymph node metastases of prostate cancer. J Carcinog 2009; 8:3.
- [22] Goel HL, Zarif MJ, Saluja V, Breen M, Garlick DS, Jiang Z, Wu CL, Davis RJ, FitzGerald TJ, Languino LR. Down-regulation of β1 integrin in vivo delays prostate cancer progression and increases radiosensitivity. IMPaCT Meeting, Atlanta, GA 2007; September 5-8, 2007.
- [23] Davis TL, Cress AE, Dalkin BL, Nagle RB.

Unique expression pattern of the α 6 β 4 integrin and laminin-5 in human prostate carcinoma. Prostate 2001; 46:240-248.

- [24] Allen MV, Smith GJ, Juliano R, Maygarden SJ, Mohler JL. Downregulation of the β4 integrin subunit in prostatic carcinoma and prostatic intraepithelial neoplasia. Hum Pathol 1998; 29:311-318.
- [25] Fornaro M, Tallini G, Bofetiado CJ, Bosari S, Languino LR. Down-regulation of β1C integrin, an inhibitor of cell proliferation, in prostate carcinoma. Am J Pathol 1996; 149:765-773.
- [26] Fornaro M, Tallini G, Zheng DQ, Flanagan WM, Manzotti M, Languino LR. p27(kip1) acts as a downstream effector of and is coexpressed with the β1C integrin in prostatic adenocarcinoma. J Clin Invest 1999; 103:321-329.
- [27] Fornaro M, Manzotti M, Tallini G, Slear AE, Bosari S, Ruoslahti E, Languino LR. β1C integrin in epithelial cells correlates with a nonproliferative phenotype: forced expression of β1C inhibits prostate epithelial cell proliferation. Am J Pathol 1998; 153:1079-1087.
- [28] Perlino E, Lovecchio M, Vacca RA, Fornaro M, Moro L, Ditonno P, Battaglia M, Selvaggi FP, Mastropasqua MG, Bufo P, Languino LR. Regulation of mRNA and protein levels of β1 integrin variants in human prostate carcinoma. Am J Pathol 2000; 157:1727-1734.
- [29] Goel HL, Breen M, Zhang J, Das I, Aznavoorian-Cheshire S, Greenberg NM, Elgavish A, Languino LR. β1A integrin expression is required for type 1 insulin-like growth factor receptor mitogenic and transforming activities and localization to focal contacts. Cancer Res 2005; 65:6692-6700.
- [30] Zheng DQ, Woodard AS, Fornaro M, Tallini G, Languino LR. Prostatic carcinoma cell migration via $\alpha\nu\beta$ 3 integrin is modulated by a focal adhesion kinase pathway. Cancer Res 1999; 59:1655-1664.
- [31] Li J, Wang T, Goel HL, Jiang Z, Cai Y, Crockett KA, Zhang JZ, Jain D, Coonradt M, Manes T, Violette MS, Weinreb H, Dolinski B, FitzGerald TJ, Languino LR. A Novel Mechanism of Prostate Cancer Growth Mediated by $\alpha\nu\beta6$ Integrin and Androgen Receptor. IMPaCT Meeting, Atlanta, GA 2007; September 5-8, 2007.
- [32] Azare J, Leslie K, Al-Ahmadie H, Gerald W, Weinreb PH, Violette SM, Bromberg J. Constitutively activated Stat3 induces tumorigenesis and enhances cell motility of prostate epithelial cells through integrin β 6. Mol Cell Biol 2007; 27:4444-4453.
- [33] Trikha M, Cai Y, Grignon D, Honn KV. Identification of a novel truncated αllb integrin. Cancer Res 1998; 58:4771-4775.
- [34] Manes T, Zheng DQ, Tognin S, Woodard AS, Marchisio PC, Languino LR. $\alpha\nu\beta3$ integrin expression up-regulates cdc2, which modulates cell migration. J Cell Biol 2003; 161:817-826.
- [35] Goel HL, Fornaro M, Moro L, Teider N, Rhim JS, King M, Languino LR. Selective modulation of

type 1 insulin-like growth factor receptor signaling and functions by β 1 integrins. J Cell Biol 2004; 166:407-418.

- [36] Goel HL, Moro L, King M, Teider N, Centrella M, McCarthy TL, Holgado-Madruga M, Wong AJ, Marra E, Languino LR. β1 integrins modulate cell adhesion by regulating insulin-like growth factor-II levels in the microenvironment. Cancer Res 2006; 66:331-342.
- [37] Reddig PJ, Juliano RL. Clinging to life: cell to matrix adhesion and cell survival. Cancer Metastasis Rev 2005; 24:425-439.
- [38] Goel HL, Languino LR. Integrin signaling in cancer. In: Kumar R, editors. Molecular targeting and signal transduction. Norwell: Kluwer Academic Publishers; 2004. p. 15-31.
- [39] Tilghman RW, Parsons JT. Focal adhesion kinase as a regulator of cell tension in the progression of cancer. Semin Cancer Biol 2007; 18:45-52.
- [40] Mitra SK, Schlaepfer DD. Integrin-regulated FAK-Src signaling in normal and cancer cells. Curr Opin Cell Biol 2006; 18:516-523.
- [41] Rovin JD, Frierson HF, Jr., Ledinh W, Parsons JT, Adams RB. Expression of focal adhesion kinase in normal and pathologic human prostate tissues. Prostate 2002; 53:124-132.
- [42] Tremblay L, Hauck W, Aprikian AG, Begin LR, Chapdelaine A, Chevalier S. Focal adhesion kinase pp125FAK expression, activation and association with paxillin and p50CSK in human metastatic prostate carcinoma. Int J Cancer 1996; 68:164-171.
- [43] Ilic D, Furuta Y, Kanazawa S, Takeda N, Sobue K, Nakatsuji N, Nomura S, Fujimoto J, Okada M, Yamamoto T, Alzawa S. Reduced cell motility and enhanced focal adhesion contact formation in cells from FAK-deficient mice. Nature 1995; 377:539-543.
- [44] Slack JK, Adams RB, Rovin JD, Bissonette EA, Stoker CE, Parsons JT. Alterations in the focal adhesion kinase/Src signal transduction pathway correlate with increased migratory capacity of prostate carcinoma cells. Oncogene 2001; 20:1152-1163.
- [45] Slack-Davis JK, Martin KH, Tilghman RW, Iwanicki M, Ung EJ, Autry C, Luzzio MJ, Cooper B, Kath JC, Roberts WG, Parsons JT. Cellular characterization of a novel focal adhesion kinase inhibitor. J Biol Chem 2007; 282:14845-14852.
- [46] Nam S, Kim D, Cheng JQ, Zhang S, Lee JH, Buettner R, Mirosevich J, Lee FY, Jove R. Action of the Src family kinase inhibitor, dasatinib (BMS-354825), on human prostate cancer cells. Cancer Res 2005; 65:9185-9189.
- [47] Aprikian AG, Tremblay L, Han K, Chevalier S. Bombesin stimulates the motility of human prostate-carcinoma cells through tyrosine phosphorylation of focal adhesion kinase and of integrin-associated proteins. Int J Cancer 1997; 72:498-504.
- [48] Lacoste J, Aprikian AG, Chevalier S. Focal ad-

hesion kinase is required for bombesininduced prostate cancer cell motility. Mol Cell Endocrinol 2005; 235:51-61.

- [49] Zheng R, Iwase A, Shen R, Goodman OB, Jr., Sugimoto N, Takuwa Y, Lerner DJ, Nanus DM. Neuropeptide-stimulated cell migration in prostate cancer cells is mediated by RhoA kinase signaling and inhibited by neutral endopeptidase. Oncogene 2006; 25:5942-5952.
- [50] Sumitomo M, Shen R, Walburg M, Dai J, Geng Y, Navarro D, Boileau G, Papandreou CN, Giancotti FG, Knudsen B, Nanus DM. Neutral endopeptidase inhibits prostate cancer cell migration by blocking focal adhesion kinase signaling. J Clin Invest 2000; 106:1399-1407.
- [51] Zeng ZZ, Jia Y, Hahn NJ, Markwart SM, Rockwood KF, Livant DL. Role of focal adhesion kinase and phosphatidylinositol 3'-kinase in integrin fibronectin receptor-mediated, matrix metalloproteinase-1-dependent invasion by metastatic prostate cancer cells. Cancer Res 2006; 66:8091-8099.
- [52] Pearson G, Robinson F, Beers Gibson T, Xu BE, Karandikar M, Berman K, Cobb MH. Mitogenactivated protein (MAP) kinase pathways: regulation and physiological functions. Endocr Rev 2001; 22:153-183.
- [53] Bakin RE, Gioeli D, Bissonette EA, Weber MJ. Attenuation of Ras signaling restores androgen sensitivity to hormone-refractory C4-2 prostate cancer cells. Cancer Res 2003; 63:1975-1980.
- [54] Weber MJ, Gioeli D. Ras signaling in prostate cancer progression. J Cell Biochem 2004; 91:13-25.
- [55] Aguirre-Ghiso JA, Estrada Y, Liu D, Ossowski L. ERK(MAPK) activity as a determinant of tumor growth and dormancy; regulation by p38(SAPK). Cancer Res 2003; 63:1684-1695.
- [56] Onishi T, Yamakawa K, Franco OE, Kawamura J, Watanabe M, Shiraishi T, Kitazawa S. Mitogen-activated protein kinase pathway is involved in α6 integrin gene expression in androgen-independent prostate cancer cells: role of proximal Sp1 consensus sequence. Biochim Biophys Acta 2001; 1538:218-227.
- [57] Majumder PK, Sellers WR. Akt-regulated pathways in prostate cancer. Oncogene 2005; 24:7465-7474.
- [58] Tamura M, Gu J, Matsumoto K, Aota S, Parsons R, Yamada KM. Inhibition of cell migration, spreading and focal adhesions by tumor suppressor PTEN. Science 1998; 280:1614-1617.
- [59] Steck PA, Pershouse MA, Jasser SA, Yung WK, Lin H, Ligon AH, Langford LA, Baumgard ML, Hattier T, Davis T, Frye C, Hu R, Swedlund B, Teng DH, Tavtigian SV. Identification of a candidate tumour suppressor gene, MMAC1, at chromosome 10q23.3 that is mutated in multiple advanced cancers. Nat Genet 1997; 15:356-362.
- [60] Li J, Yen C, Liaw D, Podsypanina K, Bose S, Wang SI, Puc J, Miliaresis C, Rodgers L,

McCombie R, Bigner SH, Giovanella BC, Ittmann M, Tycko B, Hibshoosh H, Wigler MH, Parsons R. PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. Science 1997; 275:1943-1947.

- [61] Schmitz M, Grignard G, Margue C, Dippel W, Capesius C, Mossong J, Nathan M, Giacchi S, Scheiden R, Kieffer N. Complete loss of PTEN expression as a possible early prognostic marker for prostate cancer metastasis. Int J Cancer 2007; 120:1284-1292.
- [62] Chen J, De S, Brainard J, Byzova TV. Metastatic properties of prostate cancer cells are controlled by VEGF. Cell Commun Adhes 2004; 11:1-11.
- [63] Wu Z, McRoberts KS, Theodorescu D. The role of PTEN in prostate cancer cell tropism to the bone micro-environment. Carcinogenesis 2007; 28:1393-1400.
- [64] Cantley LC. The phosphoinositide 3-kinase pathway. Science 2002; 296:1655-1657.
- [65] Khwaja A, Rodriguez-Viciana P, Wennstrom S, Warne PH, Downward J. Matrix adhesion and Ras transformation both activate a phosphoinositide 3-OH kinase and protein kinase B/Akt cellular survival pathway. EMBO J 1997; 16:2783-2793.
- [66] Lipscomb EA, Mercurio AM. Mobilization and activation of a signaling competent α 6 β 4 integrin underlies its contribution to carcinoma progression. Cancer Metastasis Rev 2005; 24:413-423.
- [67] Sun M, Wang G, Paciga JE, Feldman RI, Yuan ZQ, Ma XL, Shelley SA, Jove R, Tsichlis PN, Nicosia SV, Cheng JQ. AKT1/PKBα kinase is frequently elevated in human cancers and its constitutive activation is required for oncogenic transformation in NIH3T3 cells. Am J Pathol 2001; 159:431-437.
- [68] Malik SN, Brattain M, Ghosh PM, Troyer DA, Prihoda T, Bedolla R, Kreisberg JI. Immunohistochemical demonstration of phospho-Akt in high Gleason grade prostate cancer. Clin Cancer Res 2002; 8:1168-1171.
- [69] Paweletz CP, Charboneau L, Bichsel VE, Simone NL, Chen T, Gillespie JW, Emmert-Buck MR, Roth MJ, Petricoin IE, Liotta LA. Reverse phase protein microarrays which capture disease progression show activation of prosurvival pathways at the cancer invasion front. Oncogene 2001; 20:1981-1989.
- [70] Zheng DQ, Woodard AS, Tallini G, Languino LR. Substrate specificity of $\alpha\nu\beta3$ integrin-mediated cell migration and phosphatidylinositol 3kinase/AKT pathway activation. J Biol Chem 2000; 275:24565-24574.
- [71] Shariat SF, Lotan Y, Saboorian H, Khoddami SM, Roehrborn CG, Slawin KM, Ashfaq R. Survivin expression is associated with features of biologically aggressive prostate carcinoma. Cancer 2004; 100:751-757.
- [72] Krajewska M, Krajewski S, Banares S, Huang X,

Turner B, Bubendorf L, Kallioniemi OP, Shabaik A, Vitiello A, Peehl D, Gao GJ, Reed JC. Elevated expression of inhibitor of apoptosis proteins in prostate cancer. Clin Cancer Res 2003; 9:4914-4925.

- [73] Kishi H, Igawa M, Kikuno N, Yoshino T, Urakami S, Shiina H. Expression of the survivin gene in prostate cancer: correlation with clinicopathological characteristics, proliferative activity and apoptosis. J Urol 2004; 171:1855-1860.
- [74] Fornaro M, Plescia J, Chheang S, Tallini G, Zhu YM, King M, Altieri DC, Languino LR. Fibronectin protects prostate cancer cells from tumor necrosis factor α-induced apoptosis via the AKT/Survivin pathway. J Biol Chem 2003; 278:50402-50411.
- [75] Vaira V, Lee CW, Goel HL, Bosari S, Languino LR, Altieri DC. Regulation of survivin expression by IGF-1/mTOR signaling. Oncogene 2007; 26:2678-2684.
- [76] Colombel M, Symmans F, Gil S, O'Toole KM, Chopin D, Benson M, Olsson CA, Korsmeyer S, Buttyan R. Detection of the apoptosissuppressing oncoprotein bcl-2 in hormone- refractory human prostate cancers. Am J Pathol 1993; 143:390-400.
- [77] Zellweger T, Ninck C, Bloch M, Mirlacher M, Koivisto PA, Helin HJ, Mihatsch MJ, Gasser TC, Bubendorf L. Expression patterns of potential therapeutic targets in prostate cancer. Int J Cancer 2005; 113:619-628.
- [78] Krajewska M, Krajewski S, Epstein JI, Shabaik A, Sauvageot J, Song K, Kitada S, Reed JC. Immunohistochemical analysis of bcl-2, bax, bcl-X, and mcl-1 expression in prostate cancers. Am J Pathol 1996; 148:1567-1576.
- [79] Matter ML, Ruoslahti E. A signaling pathway from the α 5 β 1 and α v β 3 integrins that elevates bcl-2 transcription. J Biol Chem 2001;

276:27757-27763.

- [80] Lee BH, Ruoslahti E. $\alpha 5\beta 1$ integrin stimulates Bcl-2 expression and cell survival through Akt, focal adhesion kinase, and Ca2+/calmodulindependent protein kinase IV. J Cell Biochem 2005; 95:1214-1223.
- [81] Keledjian K, Kyprianou N. Anoikis induction by quinazoline based α1-adrenoceptor antagonists in prostate cancer cells: antagonistic effect of bcl-2. J Urol 2003; 169:150-156.
- [82] Drake JM, Gabriel CL, Henry MD. Assessing tumor growth and distribution in a model of prostate cancer metastasis using bioluminescence imaging. Clin Exp Metastasis 2005; 22:674-684.
- [83] Liao CP, Zhong C, Saribekyan G, Bading J, Park R, Conti PS, Moats R, Berns A, Shi W, Zhou Z, Nikitin AY, Roy-Burman P. Mouse models of prostate adenocarcinoma with the capacity to monitor spontaneous carcinogenesis by bioluminescence or fluorescence. Cancer Res 2007; 67:7525-7533.
- [84] Svensson RU, Barnes JM, Rokhlin OW, Cohen MB, Henry MD. Chemotherapeutic agents upregulate the cytomegalovirus promoter: implications for bioluminescence imaging of tumor response to therapy. Cancer Res 2007; 67:10445-10454.
- [85] Svensson RU, Shey MR, Ballas ZK, Dorkin JR, Goldberg M, Akinc A, Langer R, Anderson DG, Bumcrot D, Henry MD. Assessing siRNA pharmacodynamics in a luciferase-expressing mouse. Mol Ther 2008; 16:1995-2001.
- [86] Zhang J, Park SI, Artime MC, Summy JM, Shah AN, Bomser JA, Dorfleutner A, Flynn DC, Gallick GE. AFAP-110 is overexpressed in prostate cancer and contributes to tumorigenic growth by regulating focal contacts. J Clin Invest 2007; 117:2962-2973.