

Review Article

Roles of circular RNA in breast cancer: present and future

Zehuan Li, Zhanghan Chen, Guohua Hu, Ying Jiang

Department of General Surgery, Zhongshan Hospital, Fudan University, Shanghai 200032, China

Received April 9, 2019; Accepted June 28, 2019; Epub July 15, 2019; Published July 30, 2019

Abstract: Breast cancer is one of the most common cancers with the highest morbidity and mortality among women despite the treatment approaches have advanced including surgery, endocrine therapy and targeted therapy. Novel biomarkers are warranted to be discovered for the early detection, treatment and prognosis for breast cancer. CircRNA is a class of covalently closed single-stranded circular RNA molecules without free 5' or 3' end which makes them well expressed and more stable than their linear counterparts. In this review, we mainly discuss the oncogenic or anti-oncogenic roles of circRNAs can be utilized in the treatment and prognosis of breast cancer. A large number of circRNAs have shown great potential to function in carcinogenesis, metastasis or chemoresistance of breast cancer through transcriptional regulation of RNAs including miRNA and mRNA, in addition to their promise as stable biomarkers that can be used for monitoring breast cancer progression. However, the translation phenomenon of circRNAs in breast cancer and the diagnostic value of circRNAs in breast cancer requires further investigation for which the detection of circRNAs in plasma exosomes could be worthy of a try. Above all, engineered exosomes pre-loaded with engineered anti-oncogenic circRNAs are likely to provide a novel direction in the personal medicine of breast cancer.

Keywords: CircRNA, breast cancer, metastasis, chemoresistance, exosome, fusion-circRNA

Background

Breast cancer (BC) is a globally common cancer which is the most frequently diagnosed cancer in the vast majority of the countries and the leading cause of cancer death among females in over 100 countries [1]. According to the newly data from Global Cancer Statistics 2018, over 2 million females are newly diagnosed with BC (24.2% of the total cases) and more than 60 thousand deaths are caused by BC (15.0%). In addition, incidence rates for BC have been rising in the last decade and have far exceeded those for other cancers in both developed and developing countries. Despite the surgery, endocrine therapy and targeted therapy, the morbidity and mortality of BC remain highest in female patients. Novel biomarkers are warranted to be discovered for the early detection, treatment and prognosis for BC.

Circular RNAs (circRNAs) were discovered in plants and shown to encode subviral agents [2]. It is a class of covalently closed single-

stranded circular RNA molecules formed by back-splicing. CircRNA can escape exonucleic acid degrading enzymes due to lack of a free 5' or 3' end which makes them well expressed and more stable than their linear counterparts. Alternative circularization coupled with alternative splicing can produce a variety of additional circRNAs from one gene, making such molecules have enriched sequences in conserved nucleotides [3]. There are basically four types of circRNAs including exon circRNAs (ecircRNAs), circular intronic RNAs (ciRNAs), exon-intron circRNAs (ElicRNAs), intergenic circRNAs or fusion circRNAs (f-circRNAs), which means they derive often but not always from coding exons [4-7].

A vast number of circRNAs have been discovered in a variety of cancers and they are activated in either inhibiting tumor progression or promoting tumorigenesis. On one hand, circ-PTK2 inhibits TGF- β -induced epithelial mesenchymal transition (EMT) and metastasis by controlling TIF1 γ in non-small cell lung cancer (NSCLC) [8]. Circ-MTO1 inhibits Hepatocellular Carcinoma (HCC) progression [9]. Circ-LARP4

suppress cell proliferation and invasion of gastric cancer by sponging miR-424-5p and regulating LATS1 expression [10]. Circ-FBXW7 in represses glioma tumorigenesis and its expression positively associated with glioblastoma patient overall survival [11]. Circ-ITCH inhibits bladder cancer progression by sponging miR-17/miR-224 and regulating p21, PTEN expression [12]. On the other hand, circRNA PRMT5 promotes metastasis of urothelial carcinoma of the bladder through sponging miR-30 c to induce epithelial - mesenchymal transition [13]. Circ-CCDC66 promotes colon cancer growth and metastasis [14]. In addition to solid tumor, circRNAs are also found in hematologic malignancies like acute myeloid leukemia [15]. Moreover, discovery of Kaposi's sarcoma herpesvirus-encoded circular RNAs and a human antiviral circular RNA further indicate the prominent and extensive potential of circRNAs in malignancies [16].

Main functions of circRNA

CircRNAs can exert various functions according to their parental genes [17], herein we mainly discuss the functions of circRNAs as miRNA sponges, in regulation of parental genes and translation.

miRNA sponges

MicroRNAs (miRNAs) are important post-transcriptional regulators of gene expression that acts by direct base pairing to target sites within untranslated regions of messenger RNAs. In the recent decade, miRNA activity has been proved to be affected by miRNA sponge transcripts, as known as competing endogenous RNA (ceRNA). circRNA contains selectively conserved miRNA target sites that can strongly suppresses miRNA activity, resulting in suppressed effects of miRNAs [18]. A circRNA can serve as a sponge for multiple miRNAs. For instance, circHIPK3 can sponge 9 miRNAs which all have been reported to serve as tumor-suppressive miRNAs and thus regulate cancer cell growth [19]. So far, a few circRNAs have been found in BC to serve as miRNA sponges and the detailed information is listed in **Table 1**.

Regulation of parental genes

CircRNAs can regulate their parental genes directly or indirectly. Indirectly, some circRNAs

regulate parental genes through the ceRNA mechanism. For instance, circGFRA1 might function as miR-34a sponge to regulate parental mRNA GFRA1 expression [20]. Circ-ITCH regulates ITCH by sponging miR-214 and miR-17 [21], has-circ-0001098 regulates BARD1 by sponging miR-3942 [22], hsa_circ_0052112 regulates ZNF83 by sponging miR-125a-5p [23], hsa_circ_0072995 regulates ARHGEF28 by sponging miR-30c-2-3p [24] and thus impact the expression of their parental mRNAs in BC. Intriguingly, some circRNA like circ_IRAK3 regulates IRAK3 through a positive-feedback loop circ_IRAK3/miR-3607/FOXC1/IRAK3 loop in which the target mRNA FOXC1 binds to the promoter of parental gene IRAK3 [25]. Directly, circRNAs can regulate their parental genes in cis. On one hand, circRNAs like circ_CNOT2 which include the start-codon could potentially influence the expression of the linear gene because the linear transcript from which the circRNA was spliced is now forced to use another start codon for its translation, which makes some circRNAs negatively correlated with their parental mRNAs [26]. On the other hand, ElciRNAs, a kind of circRNAs circularized with introns 'retained' between exons, have been found predominantly localized in nucleus and can enhance the expression of their parental genes in cis by interacting with Pol II, U1 snRNP and parental gene promoters [6].

Translation

CircRNAs have been predominantly regarded as a type of non-coding RNA since the advent of high-throughput RNA sequencing (RNA-seq). Most circular RNA like circ-CCDC66 promotes cancer growth and metastasis through noncoding mechanism [14]. However, a subset of circRNAs contains an open reading frame and is translated. Pamudurti et al. verified in *Drosophila*, mouse and rat and showed that a subset of circRNAs like circ-Mb1 generally share the start codon with the hosting RNA, encode proteins with specific protein domains, and are translated in a cap-independent manner [27]. In another study, circ-ZNF609 can also be translated into a protein in a splicing dependent/cap-independent manner. Driven by internal ribosome entry site, circ-FBXW7 encodes a novel protein termed FBXW7-185aa and thus inhibited proliferation and cell cycle acceleration [11]. Moreover, translation of circRNAs is likely to be regulated by multiple mechanisms [28]. For instance, translation of circMb1 may

Circular RNA in breast cancer

Table 1. Functional circRNA-miRNA-mRNA axis in breast cancer

circRNA	Parental gene	Sponged miRNA	Targeted mRNA	Specimen type	Pairs of specimens	Reference
Oncogenic						
hsa_circ_0000479	EPSTI1	miR-4753/6809	BCL11A	Tissue	37	[37]
hsa_circ_0001783	EBLN3	miR-200c-3p	ZEB1/2, ETS1	Tissue	18	[35]
hsa_circ_0001846	UBAP2	miR-661	MTA1	Tissue	78	[38]
hsa_circ_0001982	RNF111	miR-143	N/A	Tissue	29	[33]
hsa_circ_0005230	DNM3OS	miR-618	CBX8	Tissue	76	[43]
hsa_circ_0005239	GFRA1	miR-34a	GFRA1	Tissue	51	[20]
hsa_circ_0005505	IRAK3	miR-3607	FOXC1	Tissue	35	[25]
hsa_circ_0006528	PRELID2	miR-7-5p	Raf1	Tissue	20	[45]
hsa_circ_0007294	ANKS1B	miR-148a-3p miR-152-3p	USF1	Tissue	20	[32]
hsa_circ_0007534	DDX42	miR-593	MUC19	Tissue	40	[40]
hsa_circ_0008039	PRKAR1B	miR-432-5p	E2F3	Tissue	38	[36]
hsa_circ_0008717	ABCB10	miR-1271	N/A	Tissue	32	[34]
hsa_circ_0011946	SCMH1	N/A	RFC3	Tissue	12	[39]
hsa_circ_0052112	ZNF83	miR-125a-5p	ZNF83	N/A	N/A	[23]
hsa_circ_0058514	AGFG1	miR-195-5p	CCNE1	Tissue	40	[44]
hsa_circ_0072995	ARHGEF28	miR-30c-2-3p	ARHGEF28	N/A	N/A	[24]
Anti-oncogenic						
N/A	ITCH	miR-214/miR-17	ITCH	Tissue	91	[21]
hsa_circ_0000911	IFNGR2	miR-449a	Notch1	Tissue	35	[51]
hsa_circ_0001098	BARD1	miR-3942	BARD1	Tissue	15	[22]
hsa_circ_0007874	MTO1	N/A	TRAF4/Eg5	N/A	N/A	[46]
hsa_circ_0087378	NTRK2	miR-1260b	SFRP1	Tissue	23	[52]

be regulated by starvation and FOXO and that of circ-ZNF609 can be modulated by stress conditions [27, 28]. Whether circRNAs founded in breast cancer can be translated remains unknown.

Regulation of circRNAs biogenesis

CircRNAs were commonly recognized as the production of back-splicing of pre-mRNA transcripts and can resistance against RNase R. To the best of our knowledge, biogenesis of circRNA can be regulated in several mechanisms. (1) RNA binding protein (RBP): Simon J. Conn observed that the level of RNA binding protein Quaking (QKI) is regulated in the EMT and the QKI can regulate circRNAs in EMT [29]. During the process, the QKI acts by binding to recognition elements in the introns that near the circRNA-forming splice sites. Besides, the insertion of QKI motifs can also induce circRNA formation; (2) Competitive synthesis mediated by the general splicing factors: The splicing factor

muscleblind (MBL) can promote circRNA biogenesis, which is based on the abundant MBL binding sites in introns flanking sequence and that its protein can directly interact with circ-Mbl and balance its own mRNA and circ-Mbl production [30]. Moreover, it was found that once the canonical pre-mRNA processing event was disturbed it will shift toward circRNA production [31]; (3) Positive circRNA/miRNA/mRNA loop: Circ_ANKS1B (hsa-circ-0007294) abundantly sponges miR-148a-3p and miR-152-3p to increase the expression of transcription factor USF1 while circ-ANKS1B biogenesis in BC was promoted by splicing factor ESRP1 whose expression was also regulated by USF1 [32]. Similarly, circ-IRAK3 expression can be enhanced by its targeting mRNA FOXC1 binding to its parental gene IRAK3 promoter [25]. These regulation mechanisms of circRNAs may help us to better understand and utilize the functions of different circRNAs and we believe that there could be more mechanistic regulation of circRNA which remain to be discovered.

Circular RNA in breast cancer




Figure 1. CircRNAs that are found activated in BC progression including carcinogenesis, metastasis and chemoresistance.

CircRNAs validated in BC

Recently, an increasing number of circRNAs have been discovered in BC through next-generation sequencing and circRNA microarray and have been proven to show potential in diagnosis, treatment and prognosis. These circRNAs are activated in BC progression including carcinogenesis, metastasis and chemoresistance (**Figure 1**).

Carcinogenesis associated circRNAs

Among all the validated aberrantly expressed circRNAs in BC, the upregulated circRNAs are more often found to serve as oncogenic circRNAs or vice versa. For instance, hsa_circ_0001982 is upregulated in BC and promotes BC cell carcinogenesis through decreasing miR-143 (microarray, in vitro) [33]. Similarly, circ_ABCB10 (hsa_circ_0008717) promotes BC proliferation and progression through sponging miR-1271 [34]. More convincingly, hsa_circ_001783 regulates BC progression (in vitro) via sponging miR-200c-3p to regulate ZEB1/2 and ETS1 and correlated with poor clinical outcomes in BC patients [35]. Hsa_circ_0008039 depletion significantly suppressed the proliferation (in vivo), arrested cell-cycle progression and reduced migration mechanistically as ceRNA of miR-432-5p which functionally regulated E2F3 [36]. In TNBC, Circ_EPSTI1 (hsa_circ_000479)-miR-4753/6809-BCL11A axis was proven to affect the proliferation and apoptosis of TNBC (in vivo) also through the mecha-

nism of ceRNA. In addition, circ_EPSTI1 serve as an independent prognostic marker for survival in patients with TNBC [37]. Another in vivo study showed that upregulation of circ_UBAP2 (hsa_circ_0001846) promotes TNBC progression through the miR-661/MTA1 pathway (in vivo) and also predicts poor prognosis [38]. Biomathematical prediction and in vitro experiments manifested downregulation of hsa_circ_0011946 suppresses the migration and invasion of the breast cancer cell line MCF-7 by targeting RFC3 (RNA-seq) [39]. Downregulation of hsa_circ_0007534, having been proven upregulated in CRC and glioma tissue, also suppresses BC cell proliferation and invasion by targeting miR-593/MUC19 axis which is different from that in glioma [40-42]. More circRNA regulation axis have been and are being validated in vitro, including hsa_circ_0052112/miR-125a-5p/ZNF83 [23], hsa_circ_0072995/miR-30c-2-3p/ARHGEF28 (in vitro) [24], hsa_circ_0005230/miR-618/CBX8 [43] and so on.

Metastasis associated circRNAs

Not only do circRNAs participate in the carcinogenesis of BC, but also play pivotal roles in metastasis of BC. Circ_IRAK3 (hsa_circRNA_0005505) was greatly upregulated in metastatic BC cells, including brain metastatic cells, bone metastatic cells and lung metastatic cells, and is correlated with a high risk of recurrence in distant organs and the poor prognosis of BC patients [25]. Consistently, the knockdown of circ_IRAK3 can obviously repress BC

lung metastasis in vivo, and FOXC1 expression was simultaneously decreased in the BC lung metastatic lesions. Circ_ANKS1B (hsa_circ_0007294) abundantly sponges miR-148a-3p and miR-152-3p to increase the expression of transcription factor USF1 which can transcriptionally up-regulate TGF- β 1 expression, subsequently activating TGF- β 1/Smad signaling to promote EMT [32]. Circ_AGFG1 (hsa_circ_0058514) acts as a sponge of miR-195-5p to promote TNBC proliferation, mobility and invasion as well as tumorigenesis and metastasis through regulating CCNE1 expression in vivo [44]. Therefore, circRNA including circ_IRAK3, circ_ANKS1B and circ_AGFG1 might act as potential predictors and therapeutic targets for metastatic BC.

Chemoresistance associated circRNAs

A few studies which analyzed the expression of circRNAs in chemoresistant breast cancer revealed that circRNAs may play a role in either promoting or reversing BC chemoresistance. Hsa_circ_0006528 was identified as upregulated in Adriamycin (ADM) resistant MCF-7 and MDA-MB-231 cells and validated in ADM-resistant tissues. Bioinformatic and correlation analysis showed that circ_0006528/miR-7-5p/Raf1 might be an active axis participating in chemotherapeutic resistance in BC, which requires further verification and functional analysis [45]. Circ_MTO1 (hsa_circ_0007874), downregulated in monastrol resistant cell lines (MCF-7R and MDA-MB-231R), can suppress BC cell viability and promote monastrol-induced cytotoxicity through regulating the TRAF4/Eg5 axis by targeting Eg5 protein and sequestering TRAF4 from binding to Eg5 gene [46]. Still, our knowledge of the role of circRNAs in chemotherapeutic resistance is far limited due to the lack of deep mechanistic investigations and in vivo studies. Whether circRNAs could be a future direction to overcome BC chemoresistance requires further exploration.

Anti-oncogenic circRNAs

In previous studies, most circRNAs proven downregulated in BC play anti-oncogenic roles in BC progression via ceRNA mechanisms. Circ-ITCH is a renowned anti-oncogenic circRNA involved in cancer progression and it is markedly down-regulated in esophageal squamous cell carcinoma [47], colorectal cancer [48], lung

cancer [49], glioma [50] and bladder cancer [12]. Similarly, Circ-ITCH can regulate TNBC progression suppresses TNBC growth and metastasis in vivo by up-regulating the expression of its parental gene ITCH through sponging miR-214 and miR-17, and thereby activating Wnt/ β -catenin pathway [21]. Hsa_circ_000911 exerts its anti-oncogenic function by sponging miR-449a and thereby elevating the expression of Notch1 and promoting the function of the NF- κ B signaling pathway [51]. Hsa_circ_0087378 was also validated to be downregulated in ER-positive BC. Bioinformatic and correlation analysis showed that hsa_circ_0087378/miR-1260b/SFRP1 axis was proposed to be a key regulatory pathway [52]. Zhao et al. found an up-regulated anti-oncogenic circRNA Circ_BARD1 (hsa_circ_0001098) which is induced by TCDD treatment in BC cells. Also, circ_BARD1 can inhibit cell proliferation, block cell cycle, enhance apoptosis and restrain tumor growth in vivo via miR-3942/BARD1 axis [22]. Some circRNAs suppress BC progression through breast cancer stem cells (BCSCs). BCSCs have capacities including self-renewal and pluripotency, which are considered as the cause of treatment failure and are liable for metastatic dissemination. Yan et al. discovered a list of aberrantly expressed circRNAs in BCSCs and matched non-BCSCs and validated that loss of circ_VRK1 can enhance BCSCs' expansion capacities [53]. Together, anti-oncogenic circRNAs may serve as a promising therapeutic target for patients with BC and provide a future direction in the development of a novel treatment strategy for BC.

All these circRNAs involved in BC progression were corroborated to act as miRNA sponge to regulate their target gene. However, rarely has research further investigate the deeper mechanisms in the regulation of these circRNAs. Although some circRNAs are common in several cancers such as circ-ITCH [12, 21, 47, 49], hsa_circ_0007534 [40, 42], hsa_circRNA_0002908 hsa_circRNA_0000950 and so on [54], few of these circRNAs has been proven specific in BC instead of other malignancies. Moreover, a clinical trial on the diagnostic value of circRNA in cardiac disease is ongoing (<https://clinicaltrials.gov/ct2/show/NCT03170830?term=circRNA&rank=1>), which indicates whether circRNA could serve as diagnostic biomarker for BC should also be taken in to consideration. In




Figure 2. Engineered exosomes preloaded with engineered circRNAs may serve as a therapeutic option for BC. The formation and secretion of circRNAs.

addition, circRNAs involving complex background of BC like hypoxic environment are rarely discovered except for HIF1 α -associated circ_DENND4C which promotes proliferation of BC cells [55]. Together, more researches are warranted to find more BC-associated and BC-specific circRNAs and further investigate the mechanisms behind the regulation of circRNA.

Future exploration

Mystery of F-circRNA

It has been well-established that cancer-associated chromosomal translocations give rise to fusion proteins [56, 57]. Similarly, fusion-circRNAs (f-circRNAs) are produced from transcribed exons of distinct genes affected by aberrant chromosomal translocations. Some f-circRNAs, like f-circM9 (MLL/AF9) and f-circEA-2a (EML4/ALK) have been proven to be oncogenic circRNAs that not only contribute to cellular transformation and promote cell viability but also confer resistance to therapy and have tumor-promoting properties in models of leukemia and lung cancer respectively [5, 58]. Apparently, our knowledge of f-circRNAs is lim-

ited. Do f-circRNAs all play oncogenic roles in cancer? How about the expression and functions of f-circRNAs in breast cancer? Whether engineered f-circRNAs with pivotal anti-oncogenic nucleic acid fragments can be synthesized and serve as innovative treatment method? More evidences are warranted to further elucidate the role of f-circRNAs.

Detection of circRNA in BC plasma exosome

Most circRNAs are detected in cell of tissue RNA sample in previous studies on BC, but few is done in plasma (**Table 1**). Detection of circRNA in BC plasma exosome is worthy of trying. Exosomes are 30-100 nm sized microvesicles secreted by most cells, containing a variety of functional proteins and RNAs and have been found in both the blood of healthy individuals and patients with malignancies. Exosomes can be isolated from the blood through various methods, including chemical binding, immunaffinity capture and differential ultracentrifugation and serve as biomarkers for diagnosis and the monitoring of tumour progression [59]. Over 58 330 circRNAs, 15 501 lncRNAs and 18 333 mRNAs have been identified in exosomes

with expression level and possible original tissue [60]. More than 1,000 circRNAs were identified in human serum exosomes, including circ-KLDHC10 identified in serum of CRC patient, which means circRNAs originated from cancer could enter the circulation and be readily measured in the serum [61]. Circ-PDE8A was first identified in liver-metastatic pancreatic cells and further analyzed in exosome. Exosome circRNA secreted from adipocytes like circ-deubiquitination promotes the growth of HCC [62]. Intriguingly, exosomal circRNA derived from tumor like ciRS-133 can promote white adipose browning. However, there are 12,316 lncRNAs and 17,102 mRNAs but no circRNAs yet having been identified in exosomes in BC (www.exorbase.org). Whether exosome circRNA also exist in serum of BC patients? Do adipocytes secrete circRNA and thus regulate the progress of BC? More relevant investigations are warranted to elucidate these questions to further confirm whether the innovation of cancer exosomal circRNA-based diagnostic routines can be applied in the clinic for BC. Since engineered exosomes can be therapeutic options for BC [59], whether engineered exosomes preloaded with engineered anti-oncogenic circRNAs would potentiate a route towards the development of personalized medicine for BC is worthy of exploration (Figure 2).

Conclusions

In summary, a large number of circRNAs, including circ_ANKS1B, circ_AGFG1, circ_IRAK3, circ_GFRA1 and circ_EPSTI1, show great potential to function in carcinogenesis, metastasis or chemoresistance of breast cancer via transcriptional regulation of RNAs including miRNA and mRNA, in addition to their promise as stable biomarkers that can be used for monitoring breast cancer progression. The oncogenic or anti-oncogenic roles of circRNAs can be utilized in the treatment and prognosis of breast cancer. However, the translation phenomenon of circRNAs in breast cancer and the diagnostic value of circRNAs in breast cancer requires further investigation for which the detection of circRNAs in plasma exosomes could be worthy of trying. Moreover, engineered exosomes preloaded with engineered anti-oncogenic circRNAs may be likely to provide a novel direction in the personal medicine of breast cancer.

Acknowledgements

This work was supported by an individualized support project for original scientific research of Fudan University (XM03180313), a Grant (SHDC12014207) from the Shen Kang Hospital Development Center Foundation, and a Grant (14-E-34) from the Funded Project of Baoshan District Science and Technology Commission, Shanghai.

Disclosure of conflict of interest

None.

Abbreviations

BC, breast cancer; BCSC, breast cancer stem cell; CRC, colorectal cancer; ecircRNAs, exon circRNAs; ciRNAs, circular intronic RNAs; ElciRNAs, exon-intron circRNAs; NSCLC, non-small cell lung cancer; TCDD, tetrachlorodibenzo-p-dioxin; TNBC, triple negative breast cancer.

Address correspondence to: Drs. Ying Jiang and Guohua Hu, Department of General Surgery, Zhongshan Hospital, Fudan University, 180 Fenglin Road, Xuhui District, Shanghai 200032, China. Tel: 86-21-64041990; Fax: 86-21-64038472; E-mail: jiang.ying3@zs-hospital.sh.cn (YJ); hu.guohua@zs-hospital.sh.cn (GHH)

References

- [1] Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA and Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018; 68: 394-424.
- [2] Sanger HL, Klotz G, Riesner D, Gross HJ and Kleinschmidt AK. Viroids are single-stranded covalently closed circular RNA molecules existing as highly basepaired rod-like structures. *Proc Natl Acad Sci U S A* 1976; 73: 3852-3856.
- [3] Zhang XO, Wang HB, Zhang Y, Lu X, Chen LL and Yang L. Complementary sequence-mediated exon circularization. *Cell* 2014; 159: 134-147.
- [4] Memczak S, Jens M, Elefsinioti A, Torti F, Krueger J, Rybak A, Maier L, Mackowiak SD, Gregersen LH, Munschauer M, Loewer A, Ziebold U, Landthaler M, Kocks C, Le Noble F and Rajewsky N. Circular RNAs are a large class of animal RNAs with regulatory potency. *Nature* 2013; 495: 333-338.

Circular RNA in breast cancer

- [5] Guarnerio J, Bezzi M, Jeong JC, Paffenholz SV, Berry K, Naldini MM, Lo-Coco F, Tay Y, Beck AH and Pandolfi PP. Oncogenic role of fusion-circRNAs derived from cancer-associated chromosomal translocations. *Cell* 2016; 165: 289-302.
- [6] Li Z, Huang C, Bao C, Chen L, Lin M, Wang X, Zhong G, Yu B, Hu W, Dai L, Zhu P, Chang Z, Wu Q, Zhao Y, Jia Y, Xu P, Liu H and Shan G. Exon-intron circular RNAs regulate transcription in the nucleus. *Nat Struct Mol Biol* 2015; 22: 256-264.
- [7] Han B, Chao J and Yao HH. Circular RNA and its mechanisms in disease: from the bench to the clinic. *Pharmacol Ther* 2018; 187: 31-44.
- [8] Wang L, Tong X, Zhou Z, Wang S, Lei Z, Zhang T, Liu Z, Zeng Y, Li C, Zhao J, Su Z, Zhang C, Liu X, Xu G and Zhang HT. Circular RNA hsa_circ_0008305 (circPTK2) inhibits TGF-beta-induced epithelial-mesenchymal transition and metastasis by controlling TIF1gamma in non-small cell lung cancer. *Mol Cancer* 2018; 17: 140.
- [9] Han D, Li J, Wang H, Su X, Hou J, Gu Y, Qian C, Lin Y, Liu X, Huang M, Li N, Zhou W, Yu Y and Cao X. Circular RNA circMTO1 acts as the sponge of microRNA-9 to suppress hepatocellular carcinoma progression. *Hepatology* 2017; 66: 1151-1164.
- [10] Zhang J, Liu H, Hou L, Wang G, Zhang R, Huang Y, Chen X and Zhu J. Circular RNA_LARP4 inhibits cell proliferation and invasion of gastric cancer by sponging miR-424-5p and regulating LATS1 expression. *Mol Cancer* 2017; 16: 151.
- [11] Yang Y, Gao X, Zhang M, Yan S, Sun C, Xiao F, Huang N, Yang X, Zhao K, Zhou H, Huang S, Xie B and Zhang N. Novel role of FBXW7 circular RNA in repressing glioma tumorigenesis. *J Natl Cancer Inst* 2019; 111: 435.
- [12] Yang C, Yuan W, Yang X, Li P, Wang J, Han J, Tao J, Li P, Yang H, Lv Q and Zhang W. Circular RNA circ-ITCH inhibits bladder cancer progression by sponging miR-17/miR-224 and regulating p21, PTEN expression. *Mol Cancer* 2018; 17: 19.
- [13] Chen X, Chen RX, Wei WS, Li YH, Feng ZH, Tan L, Chen JW, Yuan GJ, Chen SL, Guo SJ, Xiao KH, Liu ZW, Luo JH, Zhou FJ and Xie D. PRMT5 circular RNA promotes metastasis of urothelial carcinoma of the bladder through sponging miR-30c to induce epithelial-mesenchymal transition. *Clin Cancer Res* 2018; 24: 6319-6330.
- [14] Hsiao KY, Lin YC, Gupta SK, Chang N, Yen L, Sun HS and Tsai SJ. Noncoding effects of circular RNA CCDC66 promote colon cancer growth and metastasis. *Cancer Res* 2017; 77: 2339-2350.
- [15] Hirsch S, Blatte TJ, Grasedieck S, Cocciardi S, Rouhi A, Jongen-Lavrencic M, Paschka P, Kronke J, Gaidzik VI, Dohner H, Schlenk RF, Kuchenbauer F, Dohner K, Dolnik A and Bullinger L. Circular RNAs of the nucleophosmin (NPM1) gene in acute myeloid leukemia. *Haematologica* 2017; 102: 2039-2047.
- [16] Tagawa T, Gao S, Koparde VN, Gonzalez M, Spouge JL, Serquina AP, Lurain K, Ramaswami R, Uldrick TS, Yarchoan R and Ziegelbauer JM. Discovery of Kaposi's sarcoma herpesvirus-encoded circular RNAs and a human antiviral circular RNA. *Proc Natl Acad Sci U S A* 2018; 115: 12805-12810.
- [17] Geng Y, Jiang J and Wu C. Function and clinical significance of circRNAs in solid tumors. *J Hematol Oncol* 2018; 11: 98.
- [18] Hansen TB, Jensen TI, Clausen BH, Bramsen JB, Finsen B, Damgaard CK and Kjems J. Natural RNA circles function as efficient microRNA sponges. *Nature* 2013; 495: 384-388.
- [19] Zheng Q, Bao C, Guo W, Li S, Chen J, Chen B, Luo Y, Lyu D, Li Y, Shi G, Liang L, Gu J, He X and Huang S. Circular RNA profiling reveals an abundant circHIPK3 that regulates cell growth by sponging multiple miRNAs. *Nat Commun* 2016; 7: 11215.
- [20] He RF, Liu P, Xie XM, Zhou YJ, Liao QJ, Xiong W, Li XL, Li GY, Zeng ZY and Tang HL. circGFRA1 and GFRA1 act as ceRNAs in triple negative breast cancer by regulating miR-34a. *J Exp Clin Cancer Res* 2017; 36: 145.
- [21] Wang ST, Liu LB, Li XM, Wang YF, Xie PJ, Li Q, Wang R, Wei Q, Kang YH, Meng R and Feng XH. Circ-ITCH regulates triple-negative breast cancer progression through the Wnt/beta-catenin pathway. *Neoplasia* 2019; 66: 232-239.
- [22] Zhao J, Zou H, Han C, Ma J, Zhao J and Tang J. Circular RNA BARD1 (Hsa_circ_0001098) overexpression in breast cancer cells with TCDD treatment could promote cell apoptosis via miR-3942/BARD1 axis. *Cell Cycle* 2018; 17: 2731-2744.
- [23] Zhang HD, Jiang LH, Hou JC, Zhong SL, Zhou SY, Zhu LP, Li J, Wang DD, Sun DW, Ji ZL and Tang JH. Circular RNA hsa_circ_0052112 promotes cell migration and invasion by acting as sponge for miR-125a-5p in breast cancer. *Biomed Pharmacother* 2018; 107: 1342-1353.
- [24] Zhang HD, Jiang LH, Hou JC, Zhou SY, Zhong SL, Zhu LP, Wang DD, Yang SJ, He YJ, Mao CF, Yang Y, Wang JY, Zhang Q, Xu HZ, Yu DD, Zhao JH, Tang JH and Ji ZL. Circular RNA hsa_circ_0072995 promotes breast cancer cell migration and invasion through sponge for miR-30c-2-3p. *Epigenomics* 2018; 10: 1229-1242.
- [25] Wu J, Jiang ZR, Chen C, Hu QS, Fu ZY, Chen JJ, Wang ZD, Wang Q, Li AP, Marks JR, Guo CY,

- Chen Y, Zhou JW, Yang LQ, Lin CR and Wang SY. CircIRAK3 sponges miR-3607 to facilitate breast cancer metastasis. *Cancer Lett* 2018; 430: 179-192.
- [26] Smid M, Wilting S, Uhr K, Rodriguez-Gonzalez G, de Weerd V, Prager-Van der Smissen W, van der Vlugt-Daane M, van Galen A, Nik-Zainal S, Butler A, Martin S, Davies H, Staaf J, van de Vijver M, Richardson A, MacGrogan G, Salgado R, van den Eynden G, Purdie C, Thompson A, Caldas C, Span P, Sweep F, Simpson P, Lakhani S, van Laere S, Desmedt C, Paradiso A, Eyfjord J, Broeks A, Vincent-Solomon A, Futreal A, Knappskog S, King T, Viari A, Borresen-Dale AL, Stunnenberg H, Stratton M, Foekens J, Sieuwerts A and Martens J. The circular RNome of primary breast cancer. *Genome Res* 2019; 29: 356-366.
- [27] Pamudurti NR, Bartok O, Jens M, Ashwal-Fluss R, Stottmeister C, Ruhe L, Hanan M, Wyler E, Perez-Hernandez D, Ramberger E, Shenzen S, Samson M, Dittmar G, Landthaler M, Chekulaeva M, Rajewsky N and Kadener S. Translation of CircRNAs. *Mol Cell* 2017; 66: 9-21, e27.
- [28] Legnini I, Di Timoteo G, Rossi F, Morlando M, Briganti F, Sthandier O, Fatica A, Santini T, Andronache A, Wade M, Laneve P, Rajewsky N and Bozzoni I. Circ-ZNF609 is a circular RNA that can be translated and functions in myogenesis. *Mol Cell* 2017; 66: 22-37, e29.
- [29] Conn SJ, Pillman KA, Toubia J, Conn VM, Salmanidis M, Phillips CA, Roslan S, Schreiber AW, Gregory PA and Goodall GJ. The RNA binding protein quaking regulates formation of circRNAs. *Cell* 2015; 160: 1125-1134.
- [30] Ashwal-Fluss R, Meyer M, Pamudurti NR, Ivanov A, Bartok O, Hanan M, Evantal N, Memczak S, Rajewsky N and Kadener S. circRNA biogenesis competes with pre-mRNA splicing. *Mol Cell* 2014; 56: 55-66.
- [31] Liang D, Tatomer DC, Luo Z, Wu H, Yang L, Chen LL, Cherry S and Wilusz JE. The output of protein-coding genes shifts to circular RNAs When the Pre-mRNA Processing Machinery Is limiting. *Mol Cell* 2017; 68: 940-954, e943.
- [32] Zeng K, He B, Yang BB, Xu T, Chen X, Xu M, Liu X, Sun H, Pan Y and Wang S. The pro-metastasis effect of circANKS1B in breast cancer. *Mol Cancer* 2018; 17: 160.
- [33] Tang YY, Zhao P, Zou TN, Duan JJ, Zhi R, Yang SY, Yang DC and Wang XL. Circular RNA hsa_circ_0001982 promotes breast cancer cell carcinogenesis through decreasing miR-143. *DNA Cell Biol* 2017; 36: 901-908.
- [34] Liang HF, Zhang XZ, Liu BG, Jia GT and Li WL. Circular RNA circ-ABCB10 promotes breast cancer proliferation and progression through sponging miR-1271. *Am J Cancer Res* 2017; 7: 1566-1576.
- [35] Liu Z, Zhou Y, Liang G, Ling Y, Tan W, Tan L, Andrews R, Zhong W, Zhang X, Song E and Gong C. Circular RNA hsa_circ_001783 regulates breast cancer progression via sponging miR-200c-3p. *Cell Death Dis* 2019; 10: 55.
- [36] Liu YH, Lu CT, Zhou YZ, Zhang ZH and Sun L. Circular RNA hsa_circ_0008039 promotes breast cancer cell proliferation and migration by regulating miR-432-5p/E2F3 axis. *Biochemical and Biophysical Research Communications* 2018; 502: 358-363.
- [37] Chen B, Wei W, Huang X, Xie X, Kong Y, Dai D, Yang L, Wang J, Tang H and Xie X. circEPST11 as a prognostic marker and mediator of triple-negative breast cancer progression. *Theranostics* 2018; 8: 4003-4015.
- [38] Wang S, Li Q, Wang Y, Li X, Wang R, Kang Y, Xue X, Meng R, Wei Q and Feng X. Upregulation of circ-UBAP2 predicts poor prognosis and promotes triple-negative breast cancer progression through the miR-661/MTA1 pathway. *Biochem Biophys Res Commun* 2018; 505: 996-1002.
- [39] Zhou J, Zhang WW, Peng F, Sun JY, He ZY and Wu SG. Downregulation of hsa_circ_0011946 suppresses the migration and invasion of the breast cancer cell line MCF-7 by targeting RFC3. *Cancer Manag Res* 2018; 10: 535-544.
- [40] Song LL and Xiao Y. Downregulation of hsa_circ_0007534 suppresses breast cancer cell proliferation and invasion by targeting miR-593/MUC19 signal pathway. *Biochem Biophys Res Commun* 2018; 503: 2603-2610.
- [41] Li GF, Li L, Yao ZQ and Zhuang SJ. Hsa_circ_0007534/miR-761/ZIC5 regulatory loop modulates the proliferation and migration of glioma cells. *Biochem Biophys Res Commun* 2018; 499: 765-771.
- [42] Zhang R, Xu J, Zhao J and Wang X. Silencing of hsa_circ_0007534 suppresses proliferation and induces apoptosis in colorectal cancer cells. *Eur Rev Med Pharmacol Sci* 2018; 22: 118-126.
- [43] Xu Y, Yao Y, Leng K, Ji D, Qu L, Liu Y and Cui Y. Increased expression of circular RNA circ_0005230 indicates dismal prognosis in breast cancer and regulates cell proliferation and invasion via miR-618/CBX8 signal pathway. *Cell Physiol Biochem* 2018; 51: 1710-1722.
- [44] Yang R, Xing L, Zheng X, Sun Y, Wang X and Chen J. The circRNA circAGFG1 acts as a sponge of miR-195-5p to promote triple-negative breast cancer progression through regulating CCNE1 expression. *Mol Cancer* 2019; 18: 4.
- [45] Gao DF, Zhang XF, Liu BB, Meng D, Fang K, Guo ZJ and Li LH. Screening circular RNA related to chemotherapeutic resistance in breast cancer. *Epigenomics* 2017; 9: 1175-1188.

Circular RNA in breast cancer

- [46] Liu YX, Dong YY, Zhao LP, Su LH and Luo J. Circular RNA-MTO1 suppresses breast cancer cell viability and reverses monastrol resistance through regulating the TRAF4/Eg5 axis. *Int J Oncol* 2018; 53: 1752-1762.
- [47] Li F, Zhang L, Li W, Deng J, Zheng J, An M, Lu J and Zhou Y. Circular RNA ITCH has inhibitory effect on ESCC by suppressing the Wnt/beta-catenin pathway. *Oncotarget* 2015; 6: 6001-6013.
- [48] Huang G, Zhu H, Shi Y, Wu W, Cai H and Chen X. cir-ITCH plays an inhibitory role in colorectal cancer by regulating the Wnt/beta-catenin pathway. *PLoS One* 2015; 10: e0131225.
- [49] Wan L, Zhang L, Fan K, Cheng ZX, Sun QC and Wang JJ. Circular RNA-ITCH suppresses lung cancer proliferation via inhibiting the wnt/beta-catenin pathway. *Biomed Res Int* 2016; 2016: 1579490.
- [50] Li F, Ma K, Sun M and Shi S. Identification of the tumor-suppressive function of circular RNA ITCH in glioma cells through sponging miR-214 and promoting linear ITCH expression. *Am J Transl Res* 2018; 10: 1373-1386.
- [51] Wang HL, Xiao Y, Wu L and Ma DC. Comprehensive circular RNA profiling reveals the regulatory role of the circRNA-000911/miR-449a pathway in breast carcinogenesis. *Int J Oncol* 2018; 52: 743-754.
- [52] Yuan C, Zhou L, Zhang L, Yin K, Peng J, Sha R, Zhang S, Xu Y, Sheng X, Wang Y, Lin Y, Xu S, Yin W and Lu J. Identification and integrated analysis of key differentially expressed circular RNAs in ER-positive subtype breast cancer. *Epigenomics* 2019; 11: 297-321.
- [53] Yan NN, Xu HY, Zhang JN, Xu L, Zhang YY, Zhang L, Xu YC and Zhang FC. Circular RNA profile indicates circular RNA VRK1 is negatively related with breast cancer stem cells. *Oncotarget* 2017; 8: 95704-95718.
- [54] Shi PY, Sung J, He BY, Song H, Li ZQ, Kong WM, Wang JP, Wang JM and Xue HC. Profiles of differentially expressed circRNAs in esophageal and breast cancer. *Cancer Manag Res* 2018; 10: 2207-2221.
- [55] Liang G, Liu Z, Tan L, Su AN, Jiang WG and Gong C. HIF1alpha-associated circDENND4C promotes proliferation of breast cancer cells in hypoxic environment. *Anticancer Res* 2017; 37: 4337-4343.
- [56] Takeuchi K, Soda M, Togashi Y, Suzuki R, Sakata S, Hatano S, Asaka R, Hamanaka W, Ninomiya H, Uehara H, Lim Choi Y, Satoh Y, Okumura S, Nakagawa K, Mano H and Ishikawa Y. RET, ROS1 and ALK fusions in lung cancer. *Nat Med* 2012; 18: 378-381.
- [57] Davies KD and Doebele RC. Molecular pathways: ROS1 fusion proteins in cancer. *Clin Cancer Res* 2013; 19: 4040-4045.
- [58] Tan S, Sun D, Pu W, Gou Q, Guo C, Gong Y, Li J, Wei YQ, Liu L, Zhao Y and Peng Y. Circular RNA F-circEA-2a derived from EML4-ALK fusion gene promotes cell migration and invasion in non-small cell lung cancer. *Mol Cancer* 2018; 17: 138.
- [59] Li Z, Qiu Y, Lu W, Jiang Y and Wang J. Immunotherapeutic interventions of Triple Negative Breast Cancer. *J Transl Med* 2018; 16: 147.
- [60] Li S, Li Y, Chen B, Zhao J, Yu S, Tang Y, Zheng Q, Li Y, Wang P, He X and Huang S. exoRBase: a database of circRNA, lncRNA and mRNA in human blood exosomes. *Nucleic Acids Res* 2018; 46: D106-D112.
- [61] Li Y, Zheng QP, Bao CY, Li SY, Guo WJ, Zhao J, Chen D, Gu JR, He XH and Huang SL. Circular RNA is enriched and stable in exosomes: a promising biomarker for cancer diagnosis. *Cell Research* 2015; 25: 981-984.
- [62] Zhang H, Deng T, Ge S, Liu Y, Bai M, Zhu K, Fan Q, Li J, Ning T, Tian F, Li H, Sun W, Ying G and Ba Y. Exosome circRNA secreted from adipocytes promotes the growth of hepatocellular carcinoma by targeting deubiquitination-related USP7. *Oncogene* 2019; 38: 2844-2859.