

Review Article

An overview on therapeutics attenuating amyloid β level in Alzheimer's disease: targeting neurotransmission, inflammation, oxidative stress and enhanced cholesterol levels

Xiaoling Zhou, Yifei Li, Xiaozhe Shi, Chun Ma

The Affiliated Hospital to Changchun University of Chinese Medicine, Changchun, China

Received November 26, 2015; Accepted November 26, 2015; Epub February 15, 2016; Published February 29, 2016

Abstract: Alzheimer's disease (AD) is the most common underlying cause of dementia, and novel drugs for its treatment are needed. Of the different theories explaining the development and progression of AD, "amyloid hypothesis" is the most supported by experimental data. This hypothesis states that the cleavage of amyloid precursor protein (APP) leads to the formation of amyloid beta ($A\beta$) peptides that congregate with formation and deposition of $A\beta$ plaques in the frontal cortex and hippocampus. Risk factors including neurotransmitter modulation, chronic inflammation, metal-induced oxidative stress and elevated cholesterol levels are key contributors to the disease progress. Current therapeutic strategies abating AD progression are primarily based on anti-acetylcholinesterase (AChE) inhibitors as cognitive enhancers. The AChE inhibitor, donepezil, is proven to strengthen cognitive functions and appears effective in treating moderate to severe AD patients. N-Methyl-D-aspartate receptor antagonist, memantine, is also useful, and its combination with donepezil demonstrated a strong stabilizing effect in clinical studies on AD. Nonsteroidal anti-inflammatory drugs delayed the onset and progression of AD and attenuated cognitive dysfunction. Based upon epidemiological evidence and animal studies, antioxidants emerged as potential AD preventive agents; however, clinical trials revealed inconsistencies. Pharmacokinetic and pharmacodynamic profiling demonstrated pleiotropic functions of the hypolipidemic class of drugs, statins, potentially contributing towards the prevention of AD. In addition, targeting the APP processing pathways, stimulating neuroprotective signaling mechanisms, using the amyloid anti-aggregants and $A\beta$ immunotherapy surfaced as well-tested strategies in reducing the AD-like pathology. Overall, this review covers mechanism of inducing the $A\beta$ formation, key risk factors and major therapeutics prevalent in the AD treatment nowadays. It also delineates the need for novel screening approaches towards identifying drugs that may prevent or at least limit the progression of this devastating disease.

Keywords: Donepezil, memantine, antioxidants, statins, screening

Introduction

Dementia is a neurodegenerative condition marked by diminished cognitive and thinking ability, altered personality and loss of reasoning [1]. Alzheimer's Disease (AD) is the best known and the most prevalent cause of dementia, accounting for about 60-80% of all dementia cases [2]. Owing to the increasing average life span, the prevalence of AD is sharply on the rise, with the WHO predicting above 20 million cases by 2020, and Delphi consensus study projecting about 70-80 million AD sufferers by 2050 [3-5]. Though AD is an aging-associated disorder mainly reported in patients aged 65-

85 years, it can affect younger people too, with environmental and genetic factors playing a leading contributory role in such cases [6, 7].

Presence of several pathophysiologic mechanisms in the progression of AD hinders development of a single potential treatment for the disease [8]. Although, a few therapeutic agents have gained prominence in the recent years and have reached the late stages of clinical trials [9], an overall lack of any suitable disease-modifying therapy prevents its effective management. This leads to severe damage and death of the functioning brain cells, which ultimately proves fatal [10]. In this review, we will

Therapeutics attenuating A β in Alzheimer's disease




Figure 1. APP processing pathway stimulating A β plaque deposition and AD pathology.

briefly discuss the current mechanisms, rationales and targets for therapeutic interventions in AD. We will focus on the drugs that are known to suppress AD symptoms, specifically the neurotransmitter modulators, anti-inflammatory compounds, antioxidants and cholesterol-lowering statins. We also highlight the need for new drugs that may slow the disease progression, overcoming the deficiencies of existing therapies. Unlike earlier reviews that generally focus on either causes of AD or its treatment, we concisely provide a comprehensive idea about all the essential factors promoting AD pathogenesis, as well as potential approaches to the disease prevention and therapy.

AD hypotheses and A β Generation

Although the molecular mechanisms of AD pathogenesis are well-investigated, the key reason that governs the pathology is yet contentious. The most accepted "amyloid hypothesis" points to the deposition of amyloid beta (A β) in the neurons and parenchyma as major reason for synaptic, axonal and neural dysfunc-

tions and the consecutive cognitive impairment in patients [11-13] (**Figure 1**). A β formation is largely regulated by a shift in the balance between amyloidogenic and non-amyloidogenic amyloid precursor protein (APP) processing [14]. The APP, predominant in the brain as 695, 751 and 770 amino acid isoforms, undergoes non-amyloidogenic cleavage at the α -secretase site at position 17 of the 40-42-A β amino acid domain. This results in the formation of two fragments, sAPP α and a C-terminal fragment (CTF α). sAPP α , proved to be neuroprotective, is secreted. CTF α undergoes further proteolysis by γ -secretase, yielding p3 peptide and a C-terminal fragment, CTF γ . A down-regulation of this non-amyloidogenic processing or a shift towards the amyloidogenic pathway causes β -secretase to cleave APP before the A β amino acid site, releasing CTF β and sAPP β . CTF β is further cleaved by γ -secretase, generating A β ₄₀ and A β ₄₂. The A β ₄₂ isoform is deemed more toxic, participating more in the plaque formation [15, 16]. Intracellular accumulation of hyperphosphorylated and aggregated tau in the form of neurofibrillary tangles is also closely

Therapeutics attenuating A β in Alzheimer's disease




Figure 2. Major risk factors promoting A β plaque deposition in AD.

Table 1. Important publications on Neurotransmitter Modulators as AD therapeutics

Neurotransmitters	Therapeutics	References
Cholinergic	Tacrine	Nagabukuro et al., 2005; Minarini et al., 2013; Alonso et al., 2005; Guziar et al., 2015
	Donepezil	Cacabelos, 2007
	Galanthamine	Hopkins et al., 2012; Albuquerque et al., 2001
	Rivastigmine	Rodrigues Simoes et al., 2014; Toda et al., 2003
	Xanthostigmine	Belluti et al., 2005
	Pyrrolo-isoxazole derivatives	Anand and Singh, 2012; Razavi et al., 2013;
Glutamatergic	Memantine	Golde, 2006; Farlow et al., 2008; Cummings, 2007; Danysz and Parsons, 2012; Dominguez et al., 2011; Prokselj et al., 2013
GABAergic	Etazolate	Marcade et al., 2008
	Gabapentin	Cooney et al., 2013
Serotonergic	Lecozotan	Schechter et al., 2005
	Tandospirone and Buspirone	Miller et al., 1992; Sumiyoshi et al., 2007
	PRX-3140, PF-04995274 and RQ-00000009 (RQ-9)	Ramirez et al., 2014
	SB-742457	Maher-Edwards et al., 2010
	Lu-AE-58054 (SGS-518)	Ramirez, 2013
	Dimebon	Bezprozvanny, 2010
Histaminergic	SSRI	Nagy et al., 2004
	Carebastine and mepyramine	Mizuguchi et al., 2012
	R-(alpha)-methylhistamine	Motawaj et al., 2010
	Pyrilamine	Kharmate et al., 2007
	Thioperamide and ondansetron	Passani and Blandina, 1998
Adrenergic	Chlorpheniramine	Baronio et al., 2014
	SCH58261	Melani et al., 2003

related to neuronal death in AD [17, 18]. It is believed that A β deposition precedes tau

tangle formation, with the latter triggered by A β -activated calpain and an increased tau pro-

Therapeutics attenuating A β in Alzheimer's disease




Figure 3. Prevalently used AD-drugs targeting neurotransmitters.

teolysis [19, 20]. An association of AD with several vascular and mitochondrial risk factors, e.g. diabetes mellitus, hypertension, atherosclerosis, hypercholesterolemia, metabolic syndrome and obesity, led to a view that vascular pathologies may trigger AD [21]. It is believed that apolipoprotein E (Apo E) genotype is linked with hypercholesterolaemia [22]. Amyloid and vascular mechanisms are closely related and both culminate in A β deposition via disruption and perturbation of A β transporters, especially in the blood-brain barrier (BBB) [23]. Familial AD is governed by the autosomal, dominantly inherited, rare mutations in APP and its processing molecules, such as the γ -secretase components, presenilin (PSEN)1 and PSEN2, and accounts for less than 1% of AD cases overall [24]. The present review will focus specifically on the amyloid-based A β concept of AD, highlighting the major risk factors and therapeutics.

AD therapeutics

Risk factors, such as neurotransmitter modulation, chronic inflammation, metal-induced oxidative stress and elevated cholesterol, primar-

ily contribute to AD progression (**Figure 2**). Altered neurotransmission involving cholinergic dysfunction, increased glutamate release and N-methyl-D-aspartate (NMDA) receptor action, and aberrant gamma-aminobutyric acid (GABA), histamine and serotonin functioning participate in the pathogenesis [25-29]. Elevated levels of brain-specific inflammatory cytokines and a pronounced increase in the metals, iron, copper and zinc, in the amyloid plaques of the AD brain suggest inflammation and metal-induced oxidative stress as key mechanisms in AD pathology [30, 31]. Hypercholesterolemia and cholesterol-linked pathways play an important role in AD [32]. Thus, here we elaborate on these risk factors and their mitigation approach in reducing the A β generation and, thereby, AD pathology.

Targeting neurotransmitters (**Table 1**)

Cholinesterase inhibitors: Cholinergic hypothesis of AD development focuses on the increased hydrolysis of acetylcholine by acetylcholinesterase (AChE). This leads to a reduction in synaptic acetylcholine levels detected in AD brain [33]. The resulting modulation in cholinergic neurotransmission and functions prompts learning-memory impairments and altered intellectual, behavioral and emotional responses. Clinical findings reveal significant cortical and hippocampal atrophy associated with the cholinergic modulations [33]. This cholinergic damage is promoted by butyrylcholinesterase (BuChE) that functions as a co-regulator and enhancer of AChE. However, since the effect of BuChE is more prominent at peripheral tissues, AChE gains predominance as the cholinergic neuromodulator in AD [34]. Of the known clinically-used AChE drugs, tacrine, donepezil, rivastigmine and galantamine (**Figure 3**) are the most commonly used and considered promising for AD treatment. Further, these drugs have been derivatized to improve their efficacy and potency and reduce toxic side effects [35]. Tacrine (**Figure 3**) is the foremost approved of the first-

generation AChE and BuChE inhibitors that has cholinomimetic properties [36]. Owing to tacrin's hepatotoxic effects, ring structure-modified derivatives of tacrine were synthesized [37]. Dual-binding site, homo- and heterodimeric derivatives capable of binding to both active and peripheral AChE sites were generated based on homodimers of two tacrine moieties linked by oligomethylene chains [38]. Of these, linkers with a heptamethylene tether were found to be significantly more potent than tacrine [39]. Incorporation of a protonable amino group in the middle of the tether enhanced AChE selectivity and activity, and an amide group in place of the central methylene group of a heptamethylene linker increased BuChE selectivity [40]. Some derivatives with chloro and iodo moieties were also designed [41]. Heterodimeric derivatives with indanone and phthalimide tagged to tacrine also proved suitable [42]. Tacrine heterodimers were designed by connecting the tacrine moiety with imidazole, piperidine and ferulic acid. A potent tacrine heterodimer, carbacrine, was generated by combining tacrine with the carbazole moiety of carvedilol that had IC₅₀ values of 2.15 nM and 296 nM for AChE and BuChE respectively [43]. Combination of tacrine with the hepatoprotective nitric oxide (NO) donors appeared safe [44]. Interestingly, to reduce Ca²⁺ toxicity, Ca²⁺ channel blockers, such as 1,4-dihydropyridine (DHP) were inserted forming tacripyridines, of which the one with a cycloheximide ring in DHP moiety appeared the most potent as AChE inhibitor (IC₅₀ = 0.37 μ M) [45]. Tacrine-phenyl-benzoheterocyclic derivative demonstrated AChE inhibitory property. Moreover, this benzo derivative functioning was also dependent on the methylene linker chains [46]. Donepezil (IC₅₀ = 5.7 nM) (**Figure 3**) is considered less toxic and physiologically well-accepted and was the second FDA-approved drug for treating AD. The drug enhances cholinergic transmission and attenuates neuronal damage [47]. Interaction of its benzyl piperidine and indanone groups with the indole rings at the *peripheral anionic site* (PAS) proved useful. N-benzylpiperidine derivatives with aroylthiourea, fluoro and a chloro incorporation at indanone system were also designed and demonstrated to have 30-50% of donepezil's IC₅₀ value of [48]. A combination of 3-amino-6-phenylpyridazine with N-benzylpiperidine units yielded a compound that was several times

more potent than donepezil. Combining piperidine, indanone and methylene groups resulted in a compound with the highest potency, with an IC₅₀ of 0.0018 μ M [49]. Galanthamine (IC₅₀ = 800 nM) (**Figure 3**), a tertiary alkaloid drug, manifests AChE activity reduction and modulates nicotinic acetylcholine receptors (nAChR) towards enhancing acetylcholine generation [50]. The drug is a proven allosteric modulator of nAChR [51]. Though less toxic, its reduced potency for acetylcholine release compared to tacrine led to the designing of few derivatives using alkyl linkers, especially eight to ten methylene groups, and a terminal ammonium or phthalimido group with several fold increased efficacy [52]. N-substituted galanthamine derivatives with incorporated benzylpiperidines and alkyl linkers, specifically with six methylene units, appeared to have highest AChE efficacy amongst all derivatives [43]. Rivastigmine (IC₅₀ = 4.15 μ M) (**Figure 3**), with a carbamate moiety, emerged as a new generation of AChE inhibitor which is long-acting and reversible. Benzopyrano[4,3-b]pyrrole carbamate derivatives with further methyl derivatization at carbamoyl nitrogen showed a potent inhibitory property [53]. A combination of donepezil and rivastigmine linked through 5,6-dimethoxy-indan-1-one and dialkyl-benzylamine moieties demonstrated significantly higher AChE rather than BuChE inhibition, indicating selectivity towards the former [54]. For these compounds, variations in the meta- and para-substituted derivatives were evident [54]. A heterodimer of rivastigmine and the serotonin transport inhibitor, fluoxetine, appeared as a potent second-generation dual AChE-SERT inhibitor, emphasizing the importance of incorporating dual functions in drugs [55]. Xanthostigmine derivatives possess the amyloid pro-aggregatory property due to their binding at the AChE peripheral site. Its arylidenebenzocycloalkanone derivative targeted both the active and peripheral sites, and a further incorporation of three or seven methylene units alkoxy spacer chain and arylidene moiety into the arylidene aryl ring moiety enhanced contact with PAS [56]. Of the tested meta- and para-isomers, the para-aminobenzoic acid derivative possessed a K_i value of 53 nM (AChE). Further molecular dynamics and docking studies confirmed their efficacy as AChE inhibitors. Cis-isomers of pyrrolo-isoxazole derivatives with methoxy substitution, especially at the para-position were deemed

highly potent, claiming an anti-amnestic and AChE inhibitory abilities higher than that of donepezil [57]. The polyphenolic compounds, coumarin and its derivatives, such as ensaculin (KA-672 HCl) containing a benzopyran ring and a piperazine substitution [58] and AP2238 having benzylamino group linked to coumarin via phenyl ring, are rising as AChE/BuChE inhibitors with peripheral and catalytic site-binding capacities [59]. Flavonoid derivatives linking flavonoid and benzylpiperidine through oxygen atom or alkoxy group (-OCH₂) spacers proved effective [60]. A replacement of benzyl piperidine moiety with amino alkyl or the conformationally restrained hydrophobic groups, pyrrolidine or piperidine at the meta- or para-positions was more potent, while the latter two demonstrated a greater effect [60]. Carbamate-substituted 5,7-dimethoxyflavanone, having an IC₅₀ of around 10 nM or several folds greater respectively [61].

However, cholinesterase inhibitors are not recommended for patients with advanced AD and are prescribed rather for moderate or mild AD cases. Other side effects of these drugs include unwanted cholinergic stimulation in the intestine, heart, muscle, kidney, and other organs. Thus, AChE inhibitors specifically targeting the cholinergic system of the brain are desired.

Glutamatergic alteration: Modulation in the functioning of glutamatergic neurons is generally viewed as a property of mature AD pathology, mediated strongly by the altered levels of the synaptic glutamate neurotransmitter [62]. NMDA receptor activation inducing excitotoxicity by enhanced synaptic glutamate accumulation damages the glutamatergic neurons and adversely impacts the neuronal proliferation and differentiation, causing learning, memory and cognitive impairments [62]. Further, a direct interaction between the NMDA receptors and APP is also reported, indicating the importance of glutamatergic synaptic transmission in AD [63]. Memantine (1-amino-3,5-dimethyladamantane) (**Figure 3**), an adamantane derivative, is the most used and cost-effective drug targeting the NMDA receptor for AD treatment, especially in the USA and Europe [64, 65]. It is proved through clinical studies that the drug has symptomatic effectiveness and is known to treat moderate to severe AD [66]. As detected through cultured neuronal whole-cell patch clamp recordings, memantine has a modest

affinity of about 1 mM at 70 mV [67]. The proposed mechanism appears uncompetitive, fast, voltage-dependent and with reduced tendency for entrapment in the receptor channel [67]. Memantine is capable of blocking the NMDA receptor channel and in preventing glutamate-mediated excitotoxicity by interacting with Mg²⁺ or binding to NMDA channel close to the magnesium-binding site, and thereby reducing Ca²⁺ entry in the post-synaptic neurons [68]. Other than NMDA receptor-mediated functioning, memantine also reduced A β -induced neuronal apoptosis, marked by an attenuated DNA fragmentation and altered Bcl-2 immunostaining [69]. A non-specific neurotransmitter targeting is also detected with memantine, via its ability to antagonize human α 7nAChR, indicating a concern for studies involving the co-existence of NMDA and nACh receptors [70]. However, for situations that involve aberrant functioning of both nAChR and NMDAR in AD, memantine appears very effective. A modulation of dopaminergic and serotonergic/histaminergic neurotransmission is also reported with memantine [71, 72]. Interestingly, it was claimed that a combination of memantine and cholinesterase inhibitor with different and interconnected activities is preferred: the former takes care of agitation/aggression and delusions, and the latter reduces depression, anxiety and apathy thus indicating complementary activities [73, 74].

GABAergic alteration: It is proven that transgenic mice with early-stage amyloid pathology manifest modulations in the cholinergic neurons, followed by glutamatergic and lastly the inhibitory GABAergic neurons [75]. By inducing GABA_A receptor subunit endocytosis, A β impedes synaptic inhibition [76]. GABA_A undergoes a compensatory increase in a few hippocampal regions and sub-regions where the NMDA and non-NMDA type α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPA)-receptors are reduced at the late AD stages, probably in order to maintain hippocampal functions [77]. The pyrazolopyridine compound etazolate with anxiolytic-like properties at nanomolar to low micromolar pharmacological doses selectively affected the GABA_A receptor and enhanced the propagation of α -secretase pathway towards sAPP α [78]. Another anticonvulsant drug, gabapentin, structurally related to GABA demonstrated prominent positive responses to treatment at low

Therapeutics attenuating A β in Alzheimer's disease

doses in mixed vascular/Alzheimer dementia patients [79]. However, the drug caused sedation at a high dose, and appears to affect other neurotransmitter systems, including serotonin and glutamate [80].

Serotonergic alteration: The hippocampal serotonin 5-hydroxytryptamine (5-HT) receptor, especially 5-HT_{1A} receptor, and the cortical 5-HT₆ serotonergic receptors and their metabolites, particularly 5-HIAA, play key role in the memory and cognition [81]. *Post-mortem* AD brain demonstrated a decrease in 5-HT, and its reduction in the cortex correlated with the neuronal loss at the raphe nuclei [82]. This serotonergic dysfunction is a property of the early-onset AD, reportedly with the mis-regulation of α -secretase processing [83]. Given the location of many of these receptors on terminals of other neurons, it is possible that these changes reflect the loss of cholinergic synapses as well [84, 85]. Thus, preclinical studies on 5-HT_{1A} receptor antagonists reported pro-cognitive effects with the significant abatement of glutamatergic and cholinergic transmission [86]. The 5-HT_{1A} receptor antagonist lecozotan (SRR-333) initially appeared promising, safe and well tolerated in clinical pharmacokinetic studies at a single drug dose [87]. However, adverse side effects prevented its progress into phase II clinical trials. Few partial agonists, such as tandospirone and buspirone, are being considered useful in preventing dementia [88, 89]. Agonists of the 5-HT₄ G-protein-coupled receptor, such as PRX-3140, PF-04995274 and RQ-00000009 (RQ-9), that act by enhancing the acetylcholine release, are in the phase I or II clinical studies [90]. Along with the memory enhancement, RQ-9 was capable of reducing A β deposition in the transgenic Tg2576 mice [90]. 5-HT₆ receptor antagonist, SB-742457, completed phase II clinical studies [91]. SB-742457 went through four phase II trials for mild-to-moderate AD, and was claimed comparable to donepezil [92]. Studies also revealed an additive effect of the 5-HT₆ receptor antagonist, Lu-AE-58054 (SGS-518) with donepezil [85]. An antihistamine drug, dimebon, that binds to the 5-HT₆ receptor, appeared very promising in phase II, but failed in phase III [93]. However, a dimebon derivative, P7C3, seemed to have neuroprotective properties and is being taken up for clinical observations [94]. In addition to antagonists, the 5-HT₆

receptor agonist E-6801 also proved effective in improving cognition synergistically with donepezil [95]. Serotonin reuptake inhibitors (SSRIs), eg. Fluoxetine, sertraline and citalopram, proved useful with AChE inhibitors such as rivastigmine, donepezil, etc. in dementia [96]. It is well accepted that these SSRIs may attenuate A β levels and promote neuronal survival and functioning [97]. Overall, serotonergic therapy, via targeting 5-HT_{1A}, 5-HT₄ and especially 5-HT₆ receptors, is being considered increasingly important, although mostly for mild-to-moderate AD patients on the early stages of AD.

Histaminergic modulation: It is reported that short-term and long-term memory is regulated by histamine, and a degeneration of histaminergic neurons impairs the cognition [98]. Studies on null-mutations of the excitatory receptors, H₁R and H₂R, in frontal cortex, amygdala and hippocampus revealed histidine participation in maintaining the normal synaptic plasticity and learning-memory [99]. For the histaminergic synaptic functioning, an interaction with aminergic and peptidergic systems was also reported [100]. An enhanced activity of the inhibitory H₃R attenuated the cholinergic system as well [101]. Histamine is known to contribute to dendritic cell (DC) functioning and regulates immune responses via increased interleukin IL-6 and IL-10, decreased IL-12, and enhanced secretion of chemokines and matrix metalloproteases-9 and -12, the latter participating in DC migration via the H₂R binding [102]. H₁R and H₄R triggered the T helper 2 (Th₂)-type immune responses and modulated Ca²⁺ influx, involving store-operated calcium entry (SOCE) [103]. A β -dependent alteration in T helper cell memory with altered expression of cytokines and chemokines clearly indicated a link with histidine. An aberrant postreceptor signaling in AD involving phosphoinositide hydrolysis and adenylate cyclase pathways that are constitutively suppressed by the cortical H₃R substantiated association between histidine and AD [102]. Inverse agonists of histidine receptor, carebastine and mepyramine, suppressed H₁R and also histamine generation, and prevented an alteration in Ca²⁺-signaling via alteration of protein kinase A (PKA) and cyclic adenosine monophosphate (cAMP) response element binding protein (CREB) [104]. The H₃R agonist, R-(α)-methylhistamine, was found to distinctly attenuate muscarinic

Therapeutics attenuating A β in Alzheimer's disease

Table 2. Important publications on anti-inflammatory agents as AD therapeutics

Inflammation inhibitor	Therapeutics	References
NSAID	Indomethacin	Hoozemans et al., 2001
	BF389	Blom et al., 1997
	SC-560	Choi et al., 2013
	Nitro-flurbiprofen	Jantzen et al., 2002; Cole et al., 2004
	SD-282	Koistinaho et al., 2002
	Targeting Rho-GTPases	Kubo et al., 2008
	Targeting PPAR	Nenov et al., 2014

acetylcholine receptor-dependent phospholipase C (PLC) activity and calcium-calmodulin- and cAMP-induced protein kinases, thereby predicting an attenuation of the presynaptic excitatory functions of acetylcholine in AD [105]. The H3R antagonists that promote the generation of histamine, acetylcholine, dopamine and norepinephrine, and suppress the adenylyl cyclase-PKA pathways are being promoted as drug targets for AD. The H1R antagonist, pyrilamine, inactivated His-dependent STAT6 hyperphosphorylation, while H2R antagonist, ranitidine, and H3R/H4R antagonist, thioperamide, failed to do so [106]. Rather, H3R antagonists that stimulate the cognitive domains were proposed as new therapeutic agents for AD treatment. Chlorpheniramine, an H1R antagonist, that functioned as a serotonin-norepinephrine reuptake inhibitor altered the cortical and hippocampal cholinergic tone and affected learning and memory in AD [107]. Furthermore, the H3R antagonist, thioperamide, and 5-HT3 antagonist, ondansetron, healed cholinergic deficits, suggesting their probable role in inhibiting the AD pathogenesis [108].

Adenosine receptor: Activation of adenosine receptor of A2A subtype induced the anti-inflammatory IL-10 and prevented A β deposition [109, 110]. An interaction of A β with β_2 -adrenergic receptors induced internalization and degradation of the latter, causing adrenergic and glutamatergic aberrations and reduction in β_2 -adrenergic-stimulated cAMP [111, 112]. It was found that both caffeine and adenosine receptor antagonists averted A β build-up via increased striatal PKA activity and p-CREB levels, and decreased p-JNK and p-ERK [113, 114]. The reduction in A2a adenosine receptors, therefore, stimulated pro-survival anti-

apoptotic cascades and cognitive impairments induced by A β [115]. A selective adenosine A(2A) receptor antagonist SCH58-261 also demonstrated a similar effect [116].

Targeting inflammation (Table 2)

Neuroinflammation is established as intricately associated with AD, involv-

ing the participation of complement, cytokines, chemokines and acute phase proteins [117]. Chronic complement activation leading to membrane blebbing and endocytosis, and neurite opsonization was observed in the vicinity of A β [117]. Activated microglia and astrocyte-generated cytokines, such as the pro-inflammatory IL-1 and tumor necrosis factor alpha (TNF α), regulate cyclooxygenase (COX) activity and extensively participate in APP metabolism [118, 119]. Supportively, cytokine and Transforming Growth factor beta polymorphism is observed in the AD brain [120]. In addition, these cytokines, alongside chemokine IL-8, intracellular adhesion molecule-1, macrophage colony stimulating factor and acute phase proteins such as C-reactive protein, serum amyloid A, and transthyretin coordinate the acute phase mechanisms [121-125]. The inflammatory mediators evidently activate the AChE enzyme activity and thereby aggravate the cholinergic dysfunction in AD [126]. Thus, targeting inflammation to reduce AD pathology appears to be advantageous.

Microglial generation of superoxides and oxidative intermediates is an early feature of AD [127]. Thus, therapeutics that target microglia and prevent microglial activation are being designed. Small molecules directed towards the 13-16 site of A β (HHQK domain) that binds within the microglia [128], and SD-282 that targets microglial P38-MAPK mechanism of inflammation [129] are also being assessed. Drugs aiming at the plaque-linked complement C3 and neurotoxic C5b-9 [130] and the brain opsonins integrated to the inflammatory cascade are hypothesized to suppress the membrane attack complex that mediate the complement cascade-mediated neuronal killing [131].

Therapeutics attenuating A β in Alzheimer's disease

Table 3. Important publications on Antioxidants and Mitochondrial targets as AD therapeutics and preventive agents

Anti-Oxidative stress agents	Therapeutics	References
Antioxidants	Vitamin C	Sano et al., 1997, Zandi et al., 2004
	Carotenoids	Javed et al., 2012; Vijayapadma et al., 2014
	Resveratrol	Joseph et al., 2003; Ho et al., 2009
	Curcumin	Yang et al., 2005, Ringman et al., 2012
	Neu-P11	He et al., 2013
	Green tea and food additives	Kim et al., 2009; Zhao et al., 1989; Goodman et al., 1994; Iuvone et al., 2006
Mitochondrial targeting	Coenzyme Q10 (CoQ10)	Lee et al., 2009
	α -lipoic acid	Siedlak et al., 2009
	MitoQ and plastoquinone	Kapay et al., 2011; McManus et al., 2011
	SS31	Calkins et al., 2011
Nrf2/ARE pathway targets	Tertbutylhydroquinone	Ramsey et al., 2007; Kanninen et al., 2008; Dumont et al., 2012
	Adenovirus-dependent gene delivery	Kanninen et al., 2008; Dumont et al., 2012
Metal Chelation	Desferrioxamine, EDTA and Clioquinol	Mandel et al., 2007; Amit et al., 2008; Hegde et al., 2009

However, the most well-studied compounds are the non-steroidal anti-inflammatory drug (NSAID) that had undergone epidemiological studies. The studies revealed that consistent use of NSAID has positive effect in attenuating the AD risk [132]. Indomethacin inhibited astroglial IL-1-dependent IL-6 secretion involving a reduction in the prostaglandin-2 [133]. A specific cyclooxygenase-2 (COX-2) inhibitor, BF389, followed almost a similar IL mechanism [134]; however, the COX-2 inhibitors failed to prevent cognitive decline. Cognitive restoration and reduction in the AD pathological features in triple transgenic mice by the anti-inflammatory COX-1 inhibitor, SC-560, appeared hopeful [135]. Ibuprofen, rather than adopting the COX inhibition pathway, suppressed γ -secretase activity and reduced A β levels, as found in APP-transgenic mice [136]. A nitro-derivative of ibuprofen, nitro-flurbiprofen, released NO that promoted microglial A β clearance and also prevented the reduction of plasticity-related genes [137, 138]. Very interestingly, the anti-inflammatory mode of action of nitro-ibuprofen was different from SD-282, that is used for suppressing microglial activation in AD [129]. Thus, a suitable NSAID that balances the two opposing properties in terms of microglial activation is essential. Via targeting the BACE activity, the peroxisome proliferator-activated receptor gamma agonists, pioglitazone, prevented cytokine-dependent A β production, while the antagonists, GW0072, prevented NSAID-mediated amyloid formation [139-141]. NSAID regulated Rho-GTPases and restrained the reduction in axonal functioning and astroglial migration and

activation in AD [142]. Another very interesting aspect is that the targeting of the IL-1 responsive element of 5'-untranslated region (5'UTR) of APP proved responsible for driving APP translation [143]. These drugs are also conjectured to target the interaction of AU-rich protein with the 3'-UTR of IL-1 and TNF α , or the thalidomides that block the cytokine translation [143]. Thus, targeting the NSAID-dependent cytokines as well as APP translation appears quite promising. However, long duration treatments with COX inhibitors were found to cause damage to heart, kidney, intestine and other organs [144-146]. Moreover, the belief that NSAIDs function in the ApoE ϵ 4-carrying AD population restricts the use of NSAID in other AD patients [147]. Thus, these limitations associated with the anti-inflammatory targets enlighten the need for newer compounds and drugs for AD.

Targeting metals and oxidative stress (Table 3)

Trapping of Fe, Cu and Zn ions within the amyloid plaques and their interactions with APP and A β are prominent mechanisms accelerating the amyloid pathology [143]. Several experimental techniques, such as proton-induced X-ray emission, epifluorescence microscopy, immersion autometallography [148], synchrotron X-ray fluorescence (SXRF), magnetic resonance imaging (MRI), susceptibility weighted MR (SWI), and laser capture microdissection coupled with X-ray fluorescence microscopy confirmed the metal localization in concentrations of ~15 μ M for Cu²⁺ and about 1 mM for iron and zinc [149-152]. Aluminium is another

Therapeutics attenuating A β in Alzheimer's disease

metal that has emerged as an important participant in AD [152]. Catalytic reactions involving the generation of neurotoxic hydrogen peroxide (H₂O₂) and superoxide ion generation are the major metal-mediated mechanism of A β generation. Thus, superoxide dismutase-1 and co-enzyme Q that possess endogenous antioxidant properties are assessed for their usefulness in animal models exposed to oxidative stress, and are also being tested for efficacy in AD [153]. In the current review section, we will discuss the specific participation of these heavy metals in the AD pathogenesis, followed by the therapeutics targeting the metals and their induced mechanism.

Fe: Cortical and hippocampal over-expression of hemoxygenase (HO-1) promotes heme conversion to Fe²⁺ inducing the mitochondrial insufficiency, enhancing cytochrome C oxidase activity and H₂O₂ generation [154]. Furthermore, an involvement of Fenton reaction (Fe²⁺ + H₂O₂ → Fe³⁺ + OH⁻ + OH[•]) in neurons and astroglia triggers free radical generation and oxidative stress, which then activates neurotoxic mechanisms associated with nuclear factor- κ B, p53, c-Jun transcription factors, DNA damage and apoptosis [155]. These oxidative stress mechanisms culminate in BBB disruption [156], myelin breakdown and eventually cognitive decline in AD [157]. An IRE-IRP binding that regulates APP translation is also altered due to excess iron accumulation via formation of an IRP-1 and [4Fe-4S] cluster that deregulate iron uptake and APP translation [143]. A resemblance of APP 5'-UTR sequence with the ferritin IRE stemloop strongly supports the intense participation of iron in the pathology. Moreover, an iron binding site was also found in the APP 5'-UTR, corroborating the concept of Fe-regulated APP and thereby A β generation [158, 159]. Furthermore, the iron-mediated down-regulation of α -secretase activator furin was also shown to participate in iron homeostasis [160].

Cu: The fact that copper ions bind to histidine, aspartate and tyrosine- residues in A β with a dissociation constant at the attomolar levels, and the Cu-promoted stimulation of Fenton reaction strongly support the participation of Cu in oxidative stress-mediated AD pathogenesis [161]. The released H₂O₂ interacts with copper tyrosinate that cross-links with A β , further enhancing A β generation [162]. The process of Cu-mediated toxicity during AD pathogenesis

was proven to be stimulated by an interaction of copper ions with membrane lipid rafts, altering endocytosis of APP that bears the features of a Cu transporter [163].

Zn: A Micro-PIXE study revealed the Zn(II) levels of around 70-90 μ g/g, amounting to about 1020-1060 μ M concentration in the plaque rim, core and the total plaques [143]. The zinc level as high as 1 mM was also reported in cerebral amyloid plaques, with a pre-synaptic vesicle to postsynaptic neuronal discharge [164]. The APP was found to have a Zn-binding domain located between the amino acids 181 and 200, leading to its binding to the amino acid 6-8 in A β [165]. The alteration in Zn release involves an enhanced expression of zinc transporter proteins (ZNTs, ZnT2-8) that further promotes abnormal Zn deposition in the AD brain [166]. This change in Zn homeostasis alters its normal functions, influencing APP interaction with heparin-like molecules or laminin that participates in the biological process of neurite outgrowth [167]. Zinc deposition in A β promotes γ -secretase activity via enhanced presenilin formation, and also activates transcription factors nuclear factor kappa-B and Specificity protein-1 that bind to APP promoter, thereby promoting amyloidogenic APP synthesis [168].

Al: Studies in APP transgenic Tg2576 mice revealed an increase in insoluble A β deposition upon dietary Al supplementation and consumption of Al-treated drinking water [169]. This was supported by an increased Al levels within A β core plaques peptides of aluminium dust-exposed workers [170]. Apart from inducing oxidative stress, aluminium disrupts neuronal intracellular Ca²⁺ release and stimulates DNA damage during AD pathology [171]. Because of the explicit role of transition metals in AD, metal chelation in modifying the AD progression is being widely accepted. Desferrioxamine, EDTA and clioquinol were found to be effective in attenuating AD pathogenesis in transgenic AD mice, which also proved successful in clinical trials [172-174]. The metals function via an oxidative stress mechanism, and hence antioxidant treatments are deemed as a promising preventive approach in mitigating AD.

Oxidative stress attenuation: antioxidant treatment

Animal studies on APP and PS1 transgenic models, and A β over-expressing animals re-

Therapeutics attenuating A β in Alzheimer's disease

vealed an anti-oxidant-mediated cognitive restoration associated with a reduction in hippocampal and cortical A β deposition. The antioxidants mainly studied include the vitamins (carotenoids, vitamins C and E, lipoic acid, coenzyme Q10, N-acetylcysteine, polyphenols and Ginkgo biloba extract [175]. An attenuation in reactive oxygen species (ROS) generation and A β -induced neurotoxicity in cultured primary neuronal cultures and cell lines, such as PC12 cells, as well as in Tg 2576 AD transgenic mice, was observed using vitamin E [176, 177]. The effect of vitamin E was more pronounced when combined with anti-inflammatory agents, as reported for indomethacin [178]. Patient studies revealed tocopherol-mediated attenuation in AD pathology, and its combination with vitamin C augmented the preventive effect [179, 180]. Combining vitamin E with memantine and donepezil, however, failed to add to the protective effect of vitamin E alone [181, 182]. Mitochondrial targeting aimed at mitigating the ROS generation appeared as a pertinent anti-oxidant mechanism, where coenzyme Q10 (CoQ10, ubiquinone) played a potent role [183]. However, the effect of ubiquinone or its water-soluble cognate, idebenone, though appealing in the pre-clinical studies, failed to prove effective in clinical trials [184, 185]. Treatment with the mitochondrial antioxidant, α -lipoic acid, significantly improved cognition in Tg2576 transgenic AD mice [186]. This was supported by patient data, at a level comparable to AChE drug therapy [187]. A combination of lipoic acid with the anti-oxidant, omega-3-fatty acid, appeared potent in slowing down the functional cognitive decline [188]. *In vitro* effect of lipoic acid in inhibiting the A β oligomerization was also reported [189]. Other mitochondrial antioxidants, such as MitoQ and plastoquinone, prevented oxidative stress, the astroglia-induced neuroinflammation and, ultimately, AD pathology [190, 191]. Plastoquinone also prevented hippocampal modulations of Long Time Potentiation in rats [191], and SS31 prevented alterations in the mitochondrial dynamics of transgenic AD mice neurons [192]. MitoVitE was found to be an advanced and more effective form of MitoQ [193]. Mitochondrial permeability transition pore (mPTP), that regulated mitochondrial functions, was also targeted. Dimebon, which appeared to be promising in this context, failed in clinical trials, probably because of its complex mechanism of action,

which involves AChE reduction and 5-HT₄ stimulation on one hand, and 5-HT₆ and 5-HT₃ activations that impairs cognition [194]. N-acetylcysteine (NAC) is another antioxidant that reduced malondialdehyde, increased glutathione and restored the LTP [195, 196]. Interestingly, a study revealed that a combination of NAC and lipoic acid, along with curcumin, epigallocatechin gallate and the anti-oxidant vitamins is a prominent inducer of normal cognitive functioning in Tg2576 mice [197]. Flavonoids and carotenoids themselves were also widely studied as preventive agents in AD [194]. Natural flavonoid and carotenoid, namely rutin and lutein, prevented dementia [198, 199]. Curcumin proved important in reducing A β and AChE in animal studies; however, the human studies were not so promising [200, 201]. Resveratrol containing blueberry [202] and red grape [203] caused attenuation in A β plaque deposition, with epidemiological surveys proving the red wine-induced reduced memory loss in AD [204]. However, all these natural antioxidants could be claimed effective after completion of clinical trials [194]. Supplementation with the antioxidant melatonin that quenches free radicals reduced fibrillar amyloid burdens and prevented neurodegeneration via reduction of A β -induced neuronal apoptosis, with observed protection in human studies [205]. A melatonin agonist Neu-P11 was found to be effective in rats [206]. Alkaloid and a flavonoid derivative, Silibinin, Ginkgo biloba and a long-terncaffeine intake proved useful against A β -induced damage in transgenic animals [207-210]. Green tea and food additives, such as theanine, rosamarinic acid and nordihydroguaiaretic acid had both antioxidant properties and anti-amyloid features as well [211-214]. An endogenous antioxidant mechanism targeting the nuclear receptor factor 2 (Nrf2)/ antioxidant response element (ARE) pathway also seems to be a potential alternative approach to attenuate the AD pathology [215]. Tertbutylhydroquinone or adenovirus-dependent gene delivery along with the triterpenoid CDDO-methylamide that stimulated Nrf2 expression and translocation protected against AD pathology [216, 217]. Overall, the general prevailing view is that although oxidative stress is a major risk factor for AD, the therapeutic effects of antioxidants alone in clinical trials appear less promising. Probably, the antioxidants bioavailability, water and lipid solubility,

Therapeutics attenuating A β in Alzheimer's disease

Table 4. Important publications on cholesterol-lowering drugs as AD therapeutics

Anti-cholesterol agents	Therapeutics	References
Statin	Simvastatin	Shinohara et al., 2014
	Lovastatin	Friedhoff et al., 2001; Buxbaum et al., 2002
	Atorvastatin, Cerivastatin, Fluvastatin, Pravastatin, Rosuvastatin	Barone et al., 2014

mechanism of action, time and duration of treatment, and combined use are to be taken into consideration for further preclinical and clinical studies.

Targeting hypercholesterolemia (Table 4)

APP is a transmembrane protein, and cholesterol is an integral component of the lipid membrane. Hence, the changes in cholesterol level affect the lipid raft proteins, thereby impacting the APP as well [218]. The binding of cholesterol to CTF β through the GXXXG motif is an important factor that brings APP, γ - and β -secretases close together and enhances the amyloidogenic cleavage [219]. Since the major non-amyloidogenic component, α -secretase, is not a part of the lipid raft, cholesterol contributes greater to the amyloidogenic cleavage pathway of APP [219]. Noticeably, APP is also closely associated with the cholesterol biosynthesis regulatory enzymes - sterol receptor element binding protein (SREBP) and HMG-CoA reductase [220]. Participation of cholesterol in A β aggregation was also reported through its interaction with the ganglioside GM1 found in lipid rafts of CNS [221]. Furthermore, localization of A β degrading enzymes, insulin-degrading enzyme (IDE) and neprilylin (NPE) and plasmin in the lipid rafts points towards the role of cholesterol in influencing the A β degradation [222]. A β clearance is also influenced by ApoE polymorphic alleles, ϵ 4, ϵ 3 and ϵ 2, that participate in the cholesterol transport and impact the brain cholesterol homeostasis [223]. ApoE participates in neuroinflammation by modulating the toll like receptor and the nuclear factor kappaB pathways [224]. In fact, it is believed that the effect of APOE on BBB is inflammation-mediated, since its integral cell component, astrocytes, induce inflammation when activated [225]. Increased levels of oxysterols interact with APP and A β , with several hydroxylated cholesterol forms identified in the AD brain [226]. It is believed that the oxysterols in the form of 24- and 25-hydroxycholesterol (24-OH) that, unlike

cholesterol, can cross the BBB, actually mediate the effects of cholesterol in the AD pathogenesis [226]. A significant overlap of cholesterol and metal-mediated AD pathology was also observed. Firstly, the capacity of both metals and cholesterol to bind monomeric A β and induce its oligomerization emerged as a good explanation [219]. Cholesterol itself undergoes oxidation that may enhance the A β generation [227]. Secondly, the IDE and NEP require metals for their functioning [219]. Generation of oxysterol was also conjectured to be metal-oxidative and stress-dependent [228]. Thus, an increased dietary cholesterol and metal exposure, is hypothesized to promote A β generation and reduce A β degradation in a synergistic way, thereby enhancing plaque formation.

Cholesterol lowering drugs/statins: Because of the well-proven link between cholesterol and AD, statins are believed to be therapeutic for the disease. Simvastatin, via increased PI3K/AKT activity and endothelial NO synthase pathway, contributes towards inhibiting the learning and memory impairment in the Tg2576 mice [229]. However, simvastatin has no effect on the A β level in brain. Rather, administration of simvastatin to A β -immunized mice exacerbated the amyloid angiopathy [230]. The suppression of cholesterol metabolism with the help of HMG-CoA reductase inhibitor lovastatin, or its active metabolite lovastatin acid at 10-60 mg once-daily dose, caused dose-dependent A β reduction in human subjects [231]. Epidemiological studies have demonstrated that hypercholesterolemia was a risk factor for AD, and lovastatin caused a delayed AD onset and attenuated AD development [231, 232]. Atorvastatin was reported to prevent the neuronal degeneration following A β induction. The effects of both atorvastatin and pitavastatin were mediated by attenuation of inflammation and oxidative stress and increase in the glutamatergic transporters [229, 233]. Statins were also found to suppress inflammation and microglial iNOS synthesis and NO generation in

Therapeutics attenuating A β in Alzheimer's disease

Table 5. FDA-approved drugs as AD therapeutics

Drugs	Mode of action	AD symptoms
Donepezil	AChE inhibitor	Mild to severe AD
Memantine	NMDAR antagonist	Moderate to severe AD
Rivastigmine	AChE and BuChE inhibitor	Alzheimer-like pathology
Galantamine	Cholinergic inhibitor	Alzheimer-like pathology

AD via pleiotropic actions involving isoprenyl intermediates [234]. The neuroprotective pleiotropic effects of statin included an increase in SOD activity, activation of PKC, augmentation of endothelial nitric oxide synthase (eNOS) and reduction of CoQ10 levels [235, 236]. The effects observed in clinical trials were analyzed for atorvastatin, cerivastatin, fluvastatin, pravastatin, rosuvastatin and simvastatin and were found to be independent of apoE genotype [237]. There were several controversies in regards to the statins' beneficial role, however. It was later deduced that, despite lower reported AD incidence among the statin-treated users, there is no direct link between statins and the AD risk and development. Moreover, contradicting the phase II study, phase III randomized clinical trials showed a less beneficial role of statins [237]. Thus, it is suggested that clinical trials with a large patient population, different duration and different stages of disease are needed to find out the exact role of the cholesterol-reducing agents in AD. An up-regulation of the heme oxygenase/biliverdin reductase system, probably via inhibition of BACE1, is a likely possible pathway for inhibiting the AD progression. It is hypothesized that the effects of simvastatin, lovastatin, atorvastatin and rosuvastatin are based on this mechanism [238]. Although still debatable, the use of statins may provide a useful strategy for suppressing the AD progression.

Future directions and conclusion

The present review gives an insight into the major pathophysiological risk factors promoting AD. Overall, it is observed that despite elaborate knowledge of the risk factors and mechanism of AD, only a modest choice of therapeutic tools is available for management, prevention, mitigation and treatment of the disease. Presently, symptomatic treatments are the most prevalent, and multiple studies are in progress to identify specific drugs targeting AD. With all the progress made so far, clinical trials have

accepted the AChE inhibitors and a single NMDA receptor agonist as the probable AD therapeutics. AChE inhibitors, donepezil, galantamine and rivastigmine are used in mild to moderate AD. In severe AD or in patients not responding to AChE inhibitors, memantine can be used as an alternative (**Table 5**). A combination of cholinesterase inhibitors, with memantine is also accepted as a treatment option, where the cholinergic drugs appeared more clinically beneficial even at the late stage of the disease [239]. Yet, it remains unclear at which stage of the disease the drugs should be started and which sequence of treatments should be used. A lack of knowledge on the AD-specific pharmacokinetics and bioavailability has led to a reasonably low acceptance of the antioxidants, statins and anti-inflammatory agents as AD therapeutics. An understanding of their usage-dose, latency period, stage at which efficacy is at the peak and the genetic impact may help in proceeding with the agents reducing oxidative stress, inflammation and hyperlipidemia. Targeting BACE and presenilin, the components of the γ -secretase pathway, is envisaged as reasonable approach for inhibiting the amyloidogenic pathway of APP processing. Alternatively, stimulation of signaling pathways triggering the α -secretase-based non-amyloidogenic pathway appears as a reasonable strategy. A very optimistic approach is the targeting of APP 5'-UTR. Worldwide, the drug libraries are being screened for that purpose, with the aim of identifying both the inhibitors of amyloidogenic and the promoters of non-amyloidogenic pathways of A β . Vaccine-based approaches are also designed, bringing the attention to the usefulness of A β immunotherapy. The use of amyloid anti-aggregant strategies is also being investigated. Though clinical trials for the immunotherapies are in progress, detailed clinical studies on patients are yet awaited. However, we look forward to further research and clinical trials towards understanding and identifying novel independent and interdependent strategies that modulate amyloid metabolism in attenuating A β in AD.

Disclosure of conflict of interest

None.

Therapeutics attenuating A β in Alzheimer's disease

Address correspondence to: Dr. Chun Ma, The Affiliated Hospital to Changchun University of Chinese Medicine, Changchun, China. E-mail: silfon@sina.com

References

- [1] Coria F, Rubio I and Bayon C. Alzheimer's disease, beta-amyloidosis, and aging. *Rev Neurosci* 1994; 5: 275-292.
- [2] Mattila J, Soininen H, Koikkalainen J, Rueckert D, Wolz R, Waldemar G and Lotjonen J. Optimizing the diagnosis of early Alzheimer's disease in mild cognitive impairment subjects. *J Alzheimers Dis* 2012; 32: 969-979.
- [3] Ferri CP, Prince M, Brayne C, Brodaty H, Fratiglioni L, Ganguli M, Hall K, Hasegawa K, Hendrie H, Huang Y, Jorm A, Mathers C, Menezes PR, Rimmer E, Sczufca M; Alzheimer's Disease International. Global prevalence of dementia: a Delphi consensus study. *Lancet* 2005; 366: 2112-2117.
- [4] Haan MN and Wallace R. Can dementia be prevented? Brain aging in a population-based context. *Annu Rev Public Health* 2004; 25: 1-24.
- [5] Dufouil C and Alperovitch A. [Epidemiology of Alzheimer's disease]. *Rev Prat* 2005; 55: 1869-1878.
- [6] Selkoe DJ and Podlisny MB. Deciphering the genetic basis of Alzheimer's disease. *Annu Rev Genomics Hum Genet* 2002; 3: 67-99.
- [7] Chin-Chan M, Navarro-Yepes J and Quintanilla-Vega B. Environmental pollutants as risk factors for neurodegenerative disorders: Alzheimer and Parkinson diseases. *Front Cell Neurosci* 2015; 9: 124.
- [8] Castellani RJ and Perry G. The complexities of the pathology-pathogenesis relationship in Alzheimer disease. *Biochem Pharmacol* 2014; 88: 671-676.
- [9] Jia Q, Deng Y and Qing H. Potential therapeutic strategies for Alzheimer's disease targeting or beyond beta-amyloid: insights from clinical trials. *Biomed Res Int* 2014; 2014: 837157.
- [10] Arbogast CE, Welleford EA and Netting FE. State Dementia Plans and the Alzheimer's Disease Movement: Framing Diagnosis, Prognosis, and Motivation. *J Appl Gerontol* 2015; [Epub ahead of print].
- [11] Iadecola C. Cerebrovascular effects of amyloid-beta peptides: mechanisms and implications for Alzheimer's dementia. *Cell Mol Neurobiol* 2003; 23: 681-689.
- [12] Thal DR, Griffin WS and Braak H. Parenchymal and vascular Abeta-deposition and its effects on the degeneration of neurons and cognition in Alzheimer's disease. *J Cell Mol Med* 2008; 12: 1848-1862.
- [13] Calkins MJ and Reddy PH. Amyloid beta impairs mitochondrial anterograde transport and degenerates synapses in Alzheimer's disease neurons. *Biochim Biophys Acta* 2011; 1812: 507-513.
- [14] Canobbio I, Catricala S, Balduino C and Torti M. Calmodulin regulates the non-amyloidogenic metabolism of amyloid precursor protein in platelets. *Biochim Biophys Acta* 2011; 1813: 500-506.
- [15] O'Brien RJ and Wong PC. Amyloid precursor protein processing and Alzheimer's disease. *Annu Rev Neurosci* 2011; 34: 185-204.
- [16] Zhang YW, Thompson R, Zhang H and Xu H. APP processing in Alzheimer's disease. *Mol Brain* 2011; 4: 3.
- [17] Mirra SS, Heyman A, McKeel D, Sumi SM, Crain BJ, Brownlee LM, Vogel FS, Hughes JP, van Belle G and Berg L. The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part II. Standardization of the neuropathologic assessment of Alzheimer's disease. *Neurology* 1991; 41: 479-486.
- [18] Serrano-Pozo A, Frosch MP, Masliah E and Hyman BT. Neuropathological alterations in Alzheimer disease. *Cold Spring Harb Perspect Med* 2011; 1: a006189.
- [19] Paulson JB, Ramsden M, Forster C, Sherman MA, McGowan E and Ashe KH. Amyloid plaque and neurofibrillary tangle pathology in a regulatable mouse model of Alzheimer's disease. *Am J Pathol* 2008; 173: 762-772.
- [20] Ferreira A and Bigio EH. Calpain-mediated tau cleavage: a mechanism leading to neurodegeneration shared by multiple tauopathies. *Mol Med* 2011; 17: 676-685.
- [21] Orsucci D, Mancuso M, Ienco EC, Simoncini C, Siciliano G and Bonuccelli U. Vascular factors and mitochondrial dysfunction: a central role in the pathogenesis of Alzheimer's disease. *Curr Neurovasc Res* 2013; 10: 76-80.
- [22] Huang Y. Apolipoprotein E and Alzheimer disease. *Neurology* 2006; 66: S79-85.
- [23] Erickson MA and Banks WA. Blood-brain barrier dysfunction as a cause and consequence of Alzheimer's disease. *J Cereb Blood Flow Metab* 2013; 33: 1500-1513.
- [24] Zou Z, Liu C, Che C and Huang H. Clinical genetics of Alzheimer's disease. *Biomed Res Int* 2014; 2014: 291862.
- [25] Zadori D, Veres G, Szalardy L, Klivenyi P, Toldi J and Vecsei L. Glutamatergic dysfunctioning in Alzheimer's disease and related therapeutic targets. *J Alzheimers Dis* 2014; 42 Suppl 3: S177-187.
- [26] Verhoeff NP. Acetylcholinergic neurotransmission and the beta-amyloid cascade: implications for Alzheimer's disease. *Expert Rev Neurother* 2005; 5: 277-284.

Therapeutics attenuating A β in Alzheimer's disease

- [27] Luchetti S, Huitinga I and Swaab DF. Neurosteroid and GABA-A receptor alterations in Alzheimer's disease, Parkinson's disease and multiple sclerosis. *Neuroscience* 2011; 191: 6-21.
- [28] Zlomuzica A, Dere D, Binder S, De Souza Silva MA, Huston JP and Dere E. Neuronal histamine and cognitive symptoms in Alzheimer's disease. *Neuropharmacology* 2015; [Epub ahead of print].
- [29] Rodriguez JJ, Noristani HN and Verkhatsky A. The serotonergic system in ageing and Alzheimer's disease. *Prog Neurobiol* 2012; 99: 15-41.
- [30] Vasto S, Candore G, Listi F, Balistreri CR, Colonna-Romano G, Malavolta M, Lio D, Nuzzo D, Mocchegiani E, Di Bona D and Caruso C. Inflammation, genes and zinc in Alzheimer's disease. *Brain Res Rev* 2008; 58: 96-105.
- [31] Bush AI. The metallobiology of Alzheimer's disease. *Trends Neurosci* 2003; 26: 207-214.
- [32] Daneschvar HL, Aronson MD and Smetana GW. Do statins prevent Alzheimer's disease? A narrative review. *Eur J Intern Med* 2015; 26: 666-9.
- [33] Francis PT, Palmer AM, Snape M and Wilcock GK. The cholinergic hypothesis of Alzheimer's disease: a review of progress. *J Neurol Neurosurg Psychiatry* 1999; 66: 137-147.
- [34] Thomsen T, Zendej B, Fischer JP and Kewitz H. In vitro effects of various cholinesterase inhibitors on acetyl- and butyrylcholinesterase of healthy volunteers. *Biochem Pharmacol* 1991; 41: 139-141.
- [35] Anand P and Singh B. A review on cholinesterase inhibitors for Alzheimer's disease. *Arch Pharm Res* 2013; 36: 375-399.
- [36] Nagabukuro H, Hashimoto T, Iwata M and Doi T. Effects of TAK-802, a novel acetylcholinesterase inhibitor, and tamsulosin, an α 1-adrenoceptor antagonist, and their synergistic effects on the urodynamic characteristics in a guinea-pig model of functional bladder outlet obstruction. *BJU Int* 2005; 95: 1071-1076.
- [37] Park SM, Ki SH, Han NR, Cho IJ, Ku SK, Kim SC, Zhao RJ and Kim YW. Tacrine, an oral acetylcholinesterase inhibitor, induced hepatic oxidative damage, which was blocked by liquiritigenin through GSK3-beta inhibition. *Biol Pharm Bull* 2015; 38: 184-192.
- [38] Minarini A, Milelli A, Simoni E, Rosini M, Bolognesi ML, Marchetti C and Tumiatti V. Multifunctional tacrine derivatives in Alzheimer's disease. *Curr Top Med Chem* 2013; 13: 1771-1786.
- [39] Li C, Carlier PR, Ren H, Kan KK, Hui K, Wang H, Li W, Li Z, Xiong K, Clement EC, Xue H, Liu X, Li M, Pang Y and Han Y. Alkylene tether-length dependent gamma-aminobutyric acid type A receptor competitive antagonism by tacrine dimers. *Neuropharmacology* 2007; 52: 436-443.
- [40] Colovic MB, Krstic DZ, Lazarevic-Pasti TD, Bondzic AM and Vasic VM. Acetylcholinesterase inhibitors: pharmacology and toxicology. *Curr Neuropharmacol* 2013; 11: 315-335.
- [41] Romero A, Cacabelos R, Oset-Gasque MJ, Samadi A and Marco-Contelles J. Novel tacrine-related drugs as potential candidates for the treatment of Alzheimer's disease. *Bioorg Med Chem Lett* 2013; 23: 1916-1922.
- [42] Alonso D, Dorransoro I, Rubio L, Munoz P, Garcia-Palomo E, Del Monte M, Bidon-Chanal A, Orozco M, Luque FJ, Castro A, Medina M and Martinez A. Donepezil-tacrine hybrid related derivatives as new dual binding site inhibitors of AChE. *Bioorg Med Chem* 2005; 13: 6588-6597.
- [43] Guzior N, Wieckowska A, Panek D and Malawska B. Recent development of multifunctional agents as potential drug candidates for the treatment of Alzheimer's disease. *Curr Med Chem* 2015; 22: 373-404.
- [44] Lupp A, Appenroth D, Fang L, Decker M, Lehmann J and Fleck C. Tacrine-NO donor and tacrine-ferulic acid hybrid molecules as new anti-Alzheimer agents: hepatotoxicity and influence on the cytochrome P450 system in comparison to tacrine. *Arzneimittelforschung* 2010; 60: 229-237.
- [45] Marco-Contelles J, Leon R, de Los Rios C, Guglietta A, Terencio J, Lopez MG, Garcia AG and Villarroja M. Novel multipotent tacrine-dihydropyridine hybrids with improved acetylcholinesterase inhibitory and neuroprotective activities as potential drugs for the treatment of Alzheimer's disease. *J Med Chem* 2006; 49: 7607-7610.
- [46] Huang L, Su T, Shan W, Luo Z, Sun Y, He F and Li X. Inhibition of cholinesterase activity and amyloid aggregation by berberine-phenyl-benzoheterocyclic and tacrine-phenyl-benzoheterocyclic hybrids. *Bioorg Med Chem* 2012; 20: 3038-3048.
- [47] Cacabelos R. Donepezil in Alzheimer's disease: From conventional trials to pharmacogenetics. *Neuropsychiatr Dis Treat* 2007; 3: 303-333.
- [48] Sugimoto H, Iimura Y, Yamanishi Y and Yamatsu K. Synthesis and structure-activity relationships of acetylcholinesterase inhibitors: 1-benzyl-4-[(5,6-dimethoxy-1-oxindan-2-yl)methyl]piperidine hydrochloride and related compounds. *J Med Chem* 1995; 38: 4821-4829.
- [49] Pang YP and Kozikowski AP. Prediction of the binding site of 1-benzyl-4-[(5,6-dimethoxy-1-oxindan-2-yl)methyl]piperidine in acetylcholinesterase by docking studies with the SYSDOC program. *J Comput Aided Mol Des* 1994; 8: 683-693.

Therapeutics attenuating A β in Alzheimer's disease

- [50] Hopkins TJ, Rupprecht LE, Hayes MR, Blendy JA and Schmidt HD. Galantamine, an acetylcholinesterase inhibitor and positive allosteric modulator of nicotinic acetylcholine receptors, attenuates nicotine taking and seeking in rats. *Neuropsychopharmacology* 2012; 37: 2310-2321.
- [51] Albuquerque EX, Santos MD, Alkondon M, Pereira EF and Maelicke A. Modulation of nicotinic receptor activity in the central nervous system: a novel approach to the treatment of Alzheimer disease. *Alzheimer Dis Assoc Disord* 2001; 15 Suppl 1: S19-25.
- [52] Greenblatt HM, Kryger G, Lewis T, Silman I and Sussman JL. Structure of acetylcholinesterase complexed with (-)-galanthamine at 2.3 Å resolution. *FEBS Lett* 1999; 463: 321-326.
- [53] Bolognesi ML, Bartolini M, Cavalli A, Andrisano V, Rosini M, Minarini A and Melchiorre C. Design, synthesis, and biological evaluation of conformationally restricted rivastigmine analogues. *J Med Chem* 2004; 47: 5945-5952.
- [54] Rodrigues Simoes MC, Dias Viegas FP, Moreira MS, de Freitas Silva M, Riquiel MM, da Rosa PM, Castelli MR, dos Santos MH, Soares MG and Viegas C Jr. Donepezil: an important prototype to the design of new drug candidates for Alzheimer's disease. *Mini Rev Med Chem* 2014; 14: 2-19.
- [55] Toda N, Tago K, Marumoto S, Takami K, Ori M, Yamada N, Koyama K, Naruto S, Abe K, Yamazaki R, Hara T, Aoyagi A, Abe Y, Kaneko T and Kogen H. Design, synthesis and structure-activity relationships of dual inhibitors of acetylcholinesterase and serotonin transporter as potential agents for Alzheimer's disease. *Bioorg Med Chem* 2003; 11: 1935-1955.
- [56] Belluti F, Rampa A, Piazzini L, Bisi A, Gobbi S, Bartolini M, Andrisano V, Cavalli A, Recanatini M and Valenti P. Cholinesterase inhibitors: xanthostigmine derivatives blocking the acetylcholinesterase-induced beta-amyloid aggregation. *J Med Chem* 2005; 48: 4444-4456.
- [57] Anand P and Singh B. Synthesis and evaluation of novel 4-[(3H,3aH,6aH)-3-phenyl]-4,6-dioxo-2-phenyldihydro-2H-pyrrolo[3,4-d]isoxazol-5(3H,6H,6aH-yl)benzoic acid derivatives as potent acetylcholinesterase inhibitors and anti-amnesic agents. *Bioorg Med Chem* 2012; 20: 521-530.
- [58] Razavi SF, Khoobi M, Nadri H, Sakhteman A, Moradi A, Emami S, Foroumadi A and Shafiee A. Synthesis and evaluation of 4-substituted coumarins as novel acetylcholinesterase inhibitors. *Eur J Med Chem* 2013; 64: 252-259.
- [59] Rizzo S, Bartolini M, Ceccarini L, Piazzini L, Gobbi S, Cavalli A, Recanatini M, Andrisano V and Rampa A. Targeting Alzheimer's disease: Novel indanone hybrids bearing a pharmacophoric fragment of AP2238. *Bioorg Med Chem* 2010; 18: 1749-1760.
- [60] Shen Y, Zhang J, Sheng R, Dong X, He Q, Yang B and Hu Y. Synthesis and biological evaluation of novel flavonoid derivatives as dual binding acetylcholinesterase inhibitors. *J Enzyme Inhib Med Chem* 2009; 24: 372-380.
- [61] Sawasdee P, Sabphon C, Sitthiwongwanit D and Kokpol U. Anticholinesterase activity of 7-methoxyflavones isolated from *Kaempferia parviflora*. *Phytother Res* 2009; 23: 1792-1794.
- [62] Walton HS and Dodd PR. Glutamate-glutamine cycling in Alzheimer's disease. *Neurochem Int* 2007; 50: 1052-1066.
- [63] Innocent N, Cousins SL and Stephenson FA. NMDA receptor/amyloid precursor protein interactions: a comparison between wild-type and amyloid precursor protein mutations associated with familial Alzheimer's disease. *Neurosci Lett* 2012; 515: 131-136.
- [64] Golde TE. Disease modifying therapy for AD? *J Neurochem* 2006; 99: 689-707.
- [65] Farlow MR, Graham SM and Alva G. Memantine for the treatment of Alzheimer's disease: tolerability and safety data from clinical trials. *Drug Saf* 2008; 31: 577-585.
- [66] Cummings JL. Treatment of Alzheimer's disease: the role of symptomatic agents in an era of disease-modifying therapies. *Rev Neurol Dis* 2007; 4: 57-62.
- [67] Gilling KE, Jatzke C, Hechenberger M and Parsons CG. Potency, voltage-dependency, agonist concentration-dependency, blocking kinetics and partial untrapping of the uncompetitive N-methyl-D-aspartate (NMDA) channel blocker memantine at human NMDA (GluN1/GluN2A) receptors. *Neuropharmacology* 2009; 56: 866-875.
- [68] Danysz W and Parsons CG. Alzheimer's disease, beta-amyloid, glutamate, NMDA receptors and memantine—searching for the connections. *Br J Pharmacol* 2012; 167: 324-352.
- [69] Miguel-Hidalgo JJ, Paul IA, Wanzo V and Banerjee PK. Memantine prevents cognitive impairment and reduces Bcl-2 and caspase 8 immunoreactivity in rats injected with amyloid beta1-40. *Eur J Pharmacol* 2012; 692: 38-45.
- [70] Pohanka M. Alpha7 nicotinic acetylcholine receptor is a target in pharmacology and toxicology. *Int J Mol Sci* 2012; 13: 2219-2238.
- [71] Dominguez E, Chin TY, Chen CP and Wu TY. Management of moderate to severe Alzheimer's disease: focus on memantine. *Taiwan J Obstet Gynecol* 2011; 50: 415-423.
- [72] Prokselj T, Jerin A and Kogoj A. Memantine may affect pseudobulbar affect in patients with Alzheimer's disease. *Acta Neuropsychiatr* 2013; 25: 361-366.

Therapeutics attenuating A β in Alzheimer's disease

- [73] Atri A, Molinuevo JL, Lemming O, Wirth Y, Pulte I and Wilkinson D. Memantine in patients with Alzheimer's disease receiving donepezil: new analyses of efficacy and safety for combination therapy. *Alzheimers Res Ther* 2013; 5: 6.
- [74] Gauthier S and Molinuevo JL. Benefits of combined cholinesterase inhibitor and memantine treatment in moderate-severe Alzheimer's disease. *Alzheimers Dement* 2013; 9: 326-331.
- [75] Bell KF, de Kort GJ, Steggerda S, Shigemoto R, Ribeiro-da-Silva A and Cuervo AC. Structural involvement of the glutamatergic presynaptic boutons in a transgenic mouse model expressing early onset amyloid pathology. *Neurosci Lett* 2003; 353: 143-147.
- [76] Ulrich D. Amyloid-beta Impairs Synaptic Inhibition via GABA(A) Receptor Endocytosis. *J Neurosci* 2015; 35: 9205-9210.
- [77] Jansen KL, Faull RL, Dragunow M and Synek BL. Alzheimer's disease: changes in hippocampal N-methyl-D-aspartate, quisqualate, neurotensin, adenosine, benzodiazepine, serotonin and opioid receptors—an autoradiographic study. *Neuroscience* 1990; 39: 613-627.
- [78] Marcade M, Bourdin J, Loiseau N, Peillon H, Rayer A, Drouin D, Schweighoffer F and Desire L. Etazolate, a neuroprotective drug linking GABA(A) receptor pharmacology to amyloid precursor protein processing. *J Neurochem* 2008; 106: 392-404.
- [79] Cooney C, Murphy S, Tessema H and Freyne A. Use of low-dose gabapentin for aggressive behavior in vascular and Mixed Vascular/Alzheimer Dementia. *J Neuropsychiatry Clin Neurosci* 2013; 25: 120-125.
- [80] Adam F, Bordenave L, Sessler DI and Chauvin M. Effects of a single 1200-mg preoperative dose of gabapentin on anxiety and memory. *Ann Fr Anesth Reanim* 2012; 31: e223-227.
- [81] Lin TW and Kuo YM. Exercise benefits brain function: the monoamine connection. *Brain Sci* 2013; 3: 39-53.
- [82] Chen CP, Eastwood SL, Hope T, McDonald B, Francis PT and Esiri MM. Immunocytochemical study of the dorsal and median raphe nuclei in patients with Alzheimer's disease prospectively assessed for behavioural changes. *Neuropathol Appl Neurobiol* 2000; 26: 347-355.
- [83] Cochet M, Donneger R, Cassier E, Gaven F, Lichtenthaler SF, Marin P, Bockeaert J, Dumuis A and Claeyssen S. 5-HT₄ receptors constitutively promote the non-amyloidogenic pathway of APP cleavage and interact with ADAM10. *ACS Chem Neurosci* 2013; 4: 130-140.
- [84] Hedlund PB and Sutcliffe JG. Functional, molecular and pharmacological advances in 5-HT₇ receptor research. *Trends Pharmacol Sci* 2004; 25: 481-486.
- [85] Ramirez MJ. 5-HT₆ receptors and Alzheimer's disease. *Alzheimers Res Ther* 2013; 5: 15.
- [86] Madjid N, Tottie EE, Lutgen M, Meister B, Sandin J, Kuzmin A, Stiedl O and Ogren SO. 5-Hydroxytryptamine 1A receptor blockade facilitates aversive learning in mice: interactions with cholinergic and glutamatergic mechanisms. *J Pharmacol Exp Ther* 2006; 316: 581-591.
- [87] Schechter LE, Smith DL, Rosenzweig-Lipson S, Sukoff SJ, Dawson LA, Marquis K, Jones D, Piesla M, Andree T, Nawoschik S, Harder JA, Womack MD, Buccafusco J, Terry AV, Hoebel B, Rada P, Kelly M, Abou-Gharbia M, Barrett JE and Childers W. Lecozotan (SRA-333): a selective serotonin 1A receptor antagonist that enhances the stimulated release of glutamate and acetylcholine in the hippocampus and possesses cognitive-enhancing properties. *J Pharmacol Exp Ther* 2005; 314: 1274-1289.
- [88] Sumiyoshi T, Park S, Jayathilake K, Roy A, Ertugrul A and Meltzer HY. Effect of buspirone, a serotonin1A partial agonist, on cognitive function in schizophrenia: a randomized, double-blind, placebo-controlled study. *Schizophr Res* 2007; 95: 158-168.
- [89] Miller LG, Thompson ML, Byrnes JJ, Greenblatt DJ and Shemer A. Kinetics, brain uptake, and receptor binding of tandospirone and its metabolite 1-(2-pyrimidinyl)-piperazine. *J Clin Psychopharmacol* 1992; 12: 341-345.
- [90] Ramirez MJ, Lai MK, Tordera RM and Francis PT. Serotonergic therapies for cognitive symptoms in Alzheimer's disease: rationale and current status. *Drugs* 2014; 74: 729-736.
- [91] Maher-Edwards G, Zvartau-Hind M, Hunter AJ, Gold M, Hopton G, Jacobs G, Davy M and Williams P. Double-blind, controlled phase II study of a 5-HT₆ receptor antagonist, SB-742457, in Alzheimer's disease. *Curr Alzheimer Res* 2010; 7: 374-385.
- [92] Maher-Edwards G, Dixon R, Hunter J, Gold M, Hopton G, Jacobs G, Hunter J and Williams P. SB-742457 and donepezil in Alzheimer disease: a randomized, placebo-controlled study. *Int J Geriatr Psychiatry* 2011; 26: 536-544.
- [93] Bezprozvanny I. The rise and fall of Dimebon. *Drug News Perspect* 2010; 23: 518-523.
- [94] MacMillan KS, Naidoo J, Liang J, Melito L, Williams NS, Morlock L, Huntington PJ, Estill SJ, Longgood J, Becker GL, McKnight SL, Pieper AA, De Brabander JK and Ready JM. Development of proneurogenic, neuroprotective small molecules. *J Am Chem Soc* 2011; 133: 1428-1437.
- [95] Kendall I, Sloten HA, Codony X, Burgueno J, Pauwels PJ, Vela JM and Fone KC. E-6801, a 5-HT₆ receptor agonist, improves recognition memory by combined modulation of cholinergic and glutamatergic neurotransmission in the rat. *Psychopharmacology (Berl)* 2011; 213: 413-430.

Therapeutics attenuating A β in Alzheimer's disease

- [96] Nagy CF, Kumar D, Perdomo CA, Wason S, Cullen EI and Pratt RD. Concurrent administration of donepezil HCl and sertraline HCl in healthy volunteers: assessment of pharmacokinetic changes and safety following single and multiple oral doses. *Br J Clin Pharmacol* 2004; 58 Suppl 1: 25-33.
- [97] Keowkase R, Aboukhatwa M and Luo Y. Fluoxetine protects against amyloid-beta toxicity, in part via daf-16 mediated cell signaling pathway, in *Caenorhabditis elegans*. *Neuropharmacology* 2010; 59: 358-365.
- [98] Kohler CA, da Silva WC, Benetti F and Bonini JS. Histaminergic mechanisms for modulation of memory systems. *Neural Plast* 2011; 2011: 328602.
- [99] Dai H, Kaneko K, Kato H, Fujii S, Jing Y, Xu A, Sakurai E, Kato M, Okamura N, Kuramasu A and Yanai K. Selective cognitive dysfunction in mice lacking histamine H1 and H2 receptors. *Neurosci Res* 2007; 57: 306-313.
- [100] Threlfell S, Cragg SJ, Kallo I, Turi GF, Coen CW and Greenfield SA. Histamine H3 receptors inhibit serotonin release in substantia nigra pars reticulata. *J Neurosci* 2004; 24: 8704-8710.
- [101] Philippu A and Prast H. Importance of histamine in modulatory processes, locomotion and memory. *Behav Brain Res* 2001; 124: 151-159.
- [102] Naddafi F and Mirshafiey A. The neglected role of histamine in Alzheimer's disease. *Am J Alzheimers Dis Other Demen* 2013; 28: 327-336.
- [103] Geng S, Gao YD, Yang J, Zou JJ and Guo W. Potential role of store-operated Ca²⁺ entry in Th2 response induced by histamine in human monocyte-derived dendritic cells. *Int Immunopharmacol* 2012; 12: 358-367.
- [104] Mizuguchi H, Ono S, Hattori M and Fukui H. Inverse agonistic activity of antihistamines and suppression of histamine H1 receptor gene expression. *J Pharmacol Sci* 2012; 118: 117-121.
- [105] Motawaj M, Peoc'h K, Callebert J and Arrang JM. CSF levels of the histamine metabolite tele-methylhistamine are only slightly decreased in Alzheimer's disease. *J Alzheimers Dis* 2010; 22: 861-871.
- [106] Kharmate G, Liu Z, Patterson E and Khan MM. Histamine affects STAT6 phosphorylation via its effects on IL-4 secretion: role of H1 receptors in the regulation of IL-4 production. *Int Immunopharmacol* 2007; 7: 277-286.
- [107] Baronio D, Gonchoroski T, Castro K, Zanatta G, Gottfried C and Riesgo R. Histaminergic system in brain disorders: lessons from the translational approach and future perspectives. *Ann Gen Psychiatry* 2014; 13: 34.
- [108] Passani MB and Blandina P. Cognitive implications for H3 and 5-HT3 receptor modulation of cortical cholinergic function: a parallel story. *Methods Find Exp Clin Pharmacol* 1998; 20: 725-733.
- [109] Bermejo P, Martin-Aragon S, Benedi J, Susin C, Felici E, Gil P, Ribera JM and Villar AM. Differences of peripheral inflammatory markers between mild cognitive impairment and Alzheimer's disease. *Immunol Lett* 2008; 117: 198-202.
- [110] Bonotis K, Krikki E, Holeva V, Aggouridaki C, Costa V and Baloyannis S. Systemic immune aberrations in Alzheimer's disease patients. *J Neuroimmunol* 2008; 193: 183-187.
- [111] Rivera-Oliver M and Diaz-Rios M. Using caffeine and other adenosine receptor antagonists and agonists as therapeutic tools against neurodegenerative diseases: a review. *Life Sci* 2014; 101: 1-9.
- [112] Gahr M, Nowak DA, Connemann BJ and Schonfeldt-Lecuona C. Cerebral Amyloid Angiopathy—a disease with implications for neurology and psychiatry. *Brain Res* 2013; 1519: 19-30.
- [113] Cupino TL and Zabel MK. Alzheimer's silent partner: cerebral amyloid angiopathy. *Transl Stroke Res* 2014; 5: 330-337.
- [114] Zeitlin R, Patel S, Burgess S, Arendash GW and Echeverria V. Caffeine induces beneficial changes in PKA signaling and JNK and ERK activities in the striatum and cortex of Alzheimer's transgenic mice. *Brain Res* 2011; 1417: 127-136.
- [115] Gelber RP, Petrovitch H, Masaki KH, Ross GW and White LR. Coffee intake in midlife and risk of dementia and its neuropathologic correlates. *J Alzheimers Dis* 2011; 23: 607-615.
- [116] Melani A, Pantoni L, Bordoni F, Gianfriddo M, Bianchi L, Vannucchi MG, Bertorelli R, Monopoli A and Pedata F. The selective A2A receptor antagonist SCH 58261 reduces striatal transmitter outflow, turning behavior and ischemic brain damage induced by permanent focal ischemia in the rat. *Brain Res* 2003; 959: 243-250.
- [117] Azizi G, Navabi SS, Al-Shukaili A, Seyedzadeh MH, Yazdani R and Mirshafiey A. The Role of Inflammatory Mediators in the Pathogenesis of Alzheimer's Disease. *Sultan Qaboos Univ Med J* 2015; 15: e305-316.
- [118] Rubio-Perez JM and Morillas-Ruiz JM. A review: inflammatory process in Alzheimer's disease, role of cytokines. *ScientificWorldJournal* 2012; 2012: 756357.
- [119] Meraz-Rios MA, Toral-Rios D, Franco-Bocanegra D, Villeda-Hernandez J and Campos-Pena V. Inflammatory process in Alzheimer's Disease. *Front Integr Neurosci* 2013; 7: 59.
- [120] Ribizzi G, Fiordoro S, Barocci S, Ferrari E and Megna M. Cytokine polymorphisms and Alzheimer's

Therapeutics attenuating A β in Alzheimer's disease

- mer disease: possible associations. *Neurol Sci* 2010; 31: 321-325.
- [121] Lee SJ, Drabik K, Van Wagoner NJ, Lee S, Choi C, Dong Y and Benveniste EN. ICAM-1-induced expression of proinflammatory cytokines in astrocytes: involvement of extracellular signal-regulated kinase and p38 mitogen-activated protein kinase pathways. *J Immunol* 2000; 165: 4658-4666.
- [122] Greer MM. Inflammation and the pathophysiology of Alzheimer's disease. *Dialogues Clin Neurosci* 2000; 2: 233-239.
- [123] Yeh ET. A new perspective on the biology of C-reactive protein. *Circ Res* 2005; 97: 609-611.
- [124] Guo JT, Yu J, Grass D, de Beer FC and Kindy MS. Inflammation-dependent cerebral deposition of serum amyloid A protein in a mouse model of amyloidosis. *J Neurosci* 2002; 22: 5900-5909.
- [125] Akiyama H, Barger S, Barnum S, Bradt B, Bauer J, Cole GM, Cooper NR, Eikelenboom P, Emmerling M, Fiebich BL, Finch CE, Frautschy S, Griffin WS, Hampel H, Hull M, Landreth G, Lue L, Mrak R, Mackenzie IR, McGeer PL, O'Banion MK, Pachter J, Pasinetti G, Plata-Salaman C, Rogers J, Rydel R, Shen Y, Streit W, Strommeyer R, Tooyoma I, Van Muiswinkel FL, Veerhuis R, Walker D, Webster S, Wegrzyniak B, Wenk G and Wyss-Coray T. Inflammation and Alzheimer's disease. *Neurobiol Aging* 2000; 21: 383-421.
- [126] Richardson C, Gard PR, Klugman A, Isaac M and Tabet N. Blood pro-inflammatory cytokines in Alzheimer's disease in relation to the use of acetylcholinesterase inhibitors. *Int J Geriatr Psychiatry* 2013; 28: 1312-1317.
- [127] Colton CA, Chernyshev ON, Gilbert DL and Vitek MP. Microglial contribution to oxidative stress in Alzheimer's disease. *Ann N Y Acad Sci* 2000; 899: 292-307.
- [128] Giulian D, Haverkamp LJ, Yu J, Karshin W, Tom D, Li J, Kazanskaia A, Kirkpatrick J and Roher AE. The HHQK domain of beta-amyloid provides a structural basis for the immunopathology of Alzheimer's disease. *J Biol Chem* 1998; 273: 29719-29726.
- [129] Koistinaho M, Kettunen MI, Goldsteins G, Keinänen R, Salminen A, Ort M, Bures J, Liu D, Kauppinen RA, Higgins LS and Koistinaho J. Beta-amyloid precursor protein transgenic mice that harbor diffuse A β deposits but do not form plaques show increased ischemic vulnerability: role of inflammation. *Proc Natl Acad Sci U S A* 2002; 99: 1610-1615.
- [130] Shen Y, Yang L and Li R. What does complement do in Alzheimer's disease? Old molecules with new insights. *Transl Neurodegener* 2013; 2: 21.
- [131] Lue LF and Walker DG. Modeling Alzheimer's disease immune therapy mechanisms: interactions of human postmortem microglia with antibody-opsonized amyloid beta peptide. *J Neurosci Res* 2002; 70: 599-610.
- [132] Szekely CA and Zandi PP. Non-steroidal anti-inflammatory drugs and Alzheimer's disease: the epidemiological evidence. *CNS Neurol Disord Drug Targets* 2010; 9: 132-139.
- [133] Hoozemans JJ, Veerhuis R, Janssen I, Roze-muller AJ and Eikelenboom P. Interleukin-1 β induced cyclooxygenase 2 expression and prostaglandin E2 secretion by human neuroblastoma cells: implications for Alzheimer's disease. *Exp Gerontol* 2001; 36: 559-570.
- [134] Blom MA, van Twillert MG, de Vries SC, Engels F, Finch CE, Veerhuis R and Eikelenboom P. NSAIDs inhibit the IL-1 β -induced IL-6 release from human post-mortem astrocytes: the involvement of prostaglandin E2. *Brain Res* 1997; 777: 210-218.
- [135] Choi SH, Aid S, Caracciolo L, Minami SS, Niikura T, Matsuoka Y, Turner RS, Mattson MP and Bosetti F. Cyclooxygenase-1 inhibition reduces amyloid pathology and improves memory deficits in a mouse model of Alzheimer's disease. *J Neurochem* 2013; 124: 59-68.
- [136] Eriksen JL, Sagi SA, Smith TE, Weggen S, Das P, McLendon DC, Ozols VV, Jessing KW, Zavitz KH, Koo EH and Golde TE. NSAIDs and enantiomers of flurbiprofen target gamma-secretase and lower A β 42 in vivo. *J Clin Invest* 2003; 112: 440-449.
- [137] Jantzen PT, Connor KE, DiCarlo G, Wenk GL, Wallace JL, Rojiani AM, Coppola D, Morgan D and Gordon MN. Microglial activation and beta-amyloid deposit reduction caused by a nitric oxide-releasing nonsteroidal anti-inflammatory drug in amyloid precursor protein plus presenilin-1 transgenic mice. *J Neurosci* 2002; 22: 2246-2254.
- [138] Cole GM, Morihara T, Lim GP, Yang F, Begum A and Frautschy SA. NSAID and antioxidant prevention of Alzheimer's disease: lessons from in vitro and animal models. *Ann N Y Acad Sci* 2004; 1035: 68-84.
- [139] Papadopoulos P, Rosa-Neto P, Rochford J and Hamel E. Pioglitazone improves reversal learning and exerts mixed cerebrovascular effects in a mouse model of Alzheimer's disease with combined amyloid-beta and cerebrovascular pathology. *PLoS One* 2013; 8: e68612.
- [140] Nenov MN, Laezza F, Haidacher SJ, Zhao Y, Sadygov RG, Starkey JM, Spratt H, Luxon BA, Dineley KT and Denner L. Cognitive enhancing treatment with a PPAR γ agonist normalizes dentate granule cell presynaptic function in Tg2576 APP mice. *J Neurosci* 2014; 34: 1028-1036.

Therapeutics attenuating A β in Alzheimer's disease

- [141] Sastre M, Dewachter I, Rossner S, Bogdanovic N, Rosen E, Borghgraef P, Evert BO, Dumitrescu-Ozimek L, Thal DR, Landreth G, Walter J, Klockgether T, van Leuven F and Heneka MT. Nonsteroidal anti-inflammatory drugs repress beta-secretase gene promoter activity by the activation of PPAR γ . *Proc Natl Acad Sci U S A* 2006; 103: 443-448.
- [142] Kubo T, Yamaguchi A, Iwata N and Yamashita T. The therapeutic effects of Rho-ROCK inhibitors on CNS disorders. *Ther Clin Risk Manag* 2008; 4: 605-615.
- [143] Rogers JT and Lahiri DK. Metal and inflammatory targets for Alzheimer's disease. *Curr Drug Targets* 2004; 5: 535-551.
- [144] Bleumink GS, Feenstra J, Sturkenboom MC and Stricker BH. Nonsteroidal anti-inflammatory drugs and heart failure. *Drugs* 2003; 63: 525-534.
- [145] Whelton A and Hamilton CW. Nonsteroidal anti-inflammatory drugs: effects on kidney function. *J Clin Pharmacol* 1991; 31: 588-598.
- [146] Satoh H and Takeuchi K. Management of NSAID/aspirin-induced small intestinal damage by GI-sparing NSAIDs, anti-ulcer drugs and food constituents. *Curr Med Chem* 2012; 19: 82-89.
- [147] Imbimbo BP, Solfrizzi V and Panza F. Are NSAIDs useful to treat Alzheimer's disease or mild cognitive impairment? *Front Aging Neurosci* 2010; 2.
- [148] Stoltenberg M, Bruhn M, Sondergaard C, Doering P, West MJ, Larsen A, Troncoso JC and Danscher G. Immersion autometallographic tracing of zinc ions in Alzheimer beta-amyloid plaques. *Histochem Cell Biol* 2005; 123: 605-611.
- [149] Lovell MA, Robertson JD, Teesdale WJ, Campbell JL and Markesbery WR. Copper, iron and zinc in Alzheimer's disease senile plaques. *J Neurol Sci* 1998; 158: 47-52.
- [150] Danscher G, Jensen KB, Frederickson CJ, Kemp K, Andreassen A, Juhl S, Stoltenberg M and Ravid R. Increased amount of zinc in the hippocampus and amygdala of Alzheimer's diseased brains: a proton-induced X-ray emission spectroscopic analysis of cryostat sections from autopsy material. *J Neurosci Methods* 1997; 76: 53-59.
- [151] Miller LM, Wang Q, Telivala TP, Smith RJ, Lanzirotti A and Miklossy J. Synchrotron-based infrared and X-ray imaging shows focalized accumulation of Cu and Zn co-localized with beta-amyloid deposits in Alzheimer's disease. *J Struct Biol* 2006; 155: 30-37.
- [152] Yumoto S, Kakimi S, Ohsaki A and Ishikawa A. Demonstration of aluminum in amyloid fibers in the cores of senile plaques in the brains of patients with Alzheimer's disease. *J Inorg Biochem* 2009; 103: 1579-1584.
- [153] Yang X, Yang Y, Li G, Wang J and Yang ES. Coenzyme Q10 attenuates beta-amyloid pathology in the aged transgenic mice with Alzheimer presenilin 1 mutation. *J Mol Neurosci* 2008; 34: 165-171.
- [154] Perry G, Taddeo MA, Petersen RB, Castellani RJ, Harris PL, Siedlak SL, Cash AD, Liu Q, Nunomura A, Atwood CS and Smith MA. Adventitiously-bound redox active iron and copper are at the center of oxidative damage in Alzheimer disease. *Biometals* 2003; 16: 77-81.
- [155] Velez-Pardo C, Ospina GG and Jimenez del Rio M. A β [25-35] peptide and iron promote apoptosis in lymphocytes by an oxidative stress mechanism: involvement of H₂O₂, caspase-3, NF- κ B, p53 and c-Jun. *Neurotoxicology* 2002; 23: 351-365.
- [156] Ghribi O, Golovko MY, Larsen B, Schrag M and Murphy EJ. Deposition of iron and beta-amyloid plaques is associated with cortical cellular damage in rabbits fed with long-term cholesterol-enriched diets. *J Neurochem* 2006; 99: 438-449.
- [157] Lu PH, Lee GJ, Raven EP, Tingus K, Khoo T, Thompson PM and Bartzokis G. Age-related slowing in cognitive processing speed is associated with myelin integrity in a very healthy elderly sample. *J Clin Exp Neuropsychol* 2011; 33: 1059-1068.
- [158] Thomson AM, Rogers JT and Leedman PJ. Iron-regulatory proteins, iron-responsive elements and ferritin mRNA translation. *Int J Biochem Cell Biol* 1999; 31: 1139-1152.
- [159] Pinero DJ, Hu J and Connor JR. Alterations in the interaction between iron regulatory proteins and their iron responsive element in normal and Alzheimer's diseased brains. *Cell Mol Biol (Noisy-le-grand)* 2000; 46: 761-776.
- [160] Silvestri L and Camaschella C. A potential pathogenetic role of iron in Alzheimer's disease. *J Cell Mol Med* 2008; 12: 1548-1550.
- [161] Atwood CS, Huang X, Khatri A, Scarpa RC, Kim YS, Moir RD, Tanzi RE, Roher AE and Bush AI. Copper catalyzed oxidation of Alzheimer A β . *Cell Mol Biol (Noisy-le-grand)* 2000; 46: 777-783.
- [162] Atwood CS, Moir RD, Huang X, Scarpa RC, Bacarra NM, Romano DM, Hartshorn MA, Tanzi RE and Bush AI. Dramatic aggregation of Alzheimer abeta by Cu(II) is induced by conditions representing physiological acidosis. *J Biol Chem* 1998; 273: 12817-12826.
- [163] Hayashi T, Shishido N, Nakayama K, Nunomura A, Smith MA, Perry G and Nakamura M. Lipid peroxidation and 4-hydroxy-2-nonenal formation by copper ion bound to amyloid-beta peptide. *Free Radic Biol Med* 2007; 43: 1552-1559.

Therapeutics attenuating A β in Alzheimer's disease

- [164] Suh SW, Jensen KB, Jensen MS, Silva DS, Kesslak PJ, Danscher G and Frederickson CJ. Histochemically-reactive zinc in amyloid plaques, angiopathy, and degenerating neurons of Alzheimer's diseased brains. *Brain Res* 2000; 852: 274-278.
- [165] Bush AI, Multhaup G, Moir RD, Williamson TG, Small DH, Rumble B, Pollwein P, Beyreuther K and Masters CL. A novel zinc(II) binding site modulates the function of the beta A4 amyloid protein precursor of Alzheimer's disease. *J Biol Chem* 1993; 268: 16109-16112.
- [166] Kambe T, Yamaguchi-Iwai Y, Sasaki R and Nagao M. Overview of mammalian zinc transporters. *Cell Mol Life Sci* 2004; 61: 49-68.
- [167] Multhaup G, Mechler H and Masters CL. Characterization of the high affinity heparin binding site of the Alzheimer's disease beta A4 amyloid precursor protein (APP) and its enhancement by zinc(II). *J Mol Recognit* 1995; 8: 247-257.
- [168] Cuajungco MP and Lees GJ. Zinc metabolism in the brain: relevance to human neurodegenerative disorders. *Neurobiol Dis* 1997; 4: 137-169.
- [169] Pratico D, Uryu K, Sung S, Tang S, Trojanowski JQ and Lee VM. Aluminum modulates brain amyloidosis through oxidative stress in APP transgenic mice. *FASEB J* 2002; 16: 1138-1140.
- [170] Polizzi S, Pira E, Ferrara M, Bugiani M, Papaleo A, Albera R and Palmi S. Neurotoxic effects of aluminium among foundry workers and Alzheimer's disease. *Neurotoxicology* 2002; 23: 761-774.
- [171] Hegde ML, Anitha S, Latha KS, Mustak MS, Stein R, Ravid R and Rao KS. First evidence for helical transitions in supercoiled DNA by amyloid Beta Peptide (1-42) and aluminum: a new insight in understanding Alzheimer's disease. *J Mol Neurosci* 2004; 22: 19-31.
- [172] Amit T, Avramovich-Tirosh Y, Youdim MB and Mandel S. Targeting multiple Alzheimer's disease etiologies with multimodal neuroprotective and neurorestorative iron chelators. *FASEB J* 2008; 22: 1296-1305.
- [173] Hegde ML, Bharathi P, Suram A, Venugopal C, Jagannathan R, Poddar P, Srinivas P, Sambamurti K, Rao KJ, Scancar J, Messori L, Zecca L and Zatta P. Challenges associated with metal chelation therapy in Alzheimer's disease. *J Alzheimers Dis* 2009; 17: 457-468.
- [174] Mandel S, Amit T, Bar-Am O and Youdim MB. Iron dysregulation in Alzheimer's disease: multimodal brain permeable iron chelating drugs, possessing neuroprotective-neurorescue and amyloid precursor protein-processing regulatory activities as therapeutic agents. *Prog Neurobiol* 2007; 82: 348-360.
- [175] Mecocci P and Polidori MC. Antioxidant clinical trials in mild cognitive impairment and Alzheimer's disease. *Biochim Biophys Acta* 2012; 1822: 631-638.
- [176] Yatin SM, Varadarajan S and Butterfield DA. Vitamin E Prevents Alzheimer's Amyloid beta-Peptide (1-42)-Induced Neuronal Protein Oxidation and Reactive Oxygen Species Production. *J Alzheimers Dis* 2000; 2: 123-131.
- [177] Sung S, Yao Y, Uryu K, Yang H, Lee VM, Trojanowski JQ and Pratico D. Early vitamin E supplementation in young but not aged mice reduces Abeta levels and amyloid deposition in a transgenic model of Alzheimer's disease. *FASEB J* 2004; 18: 323-325.
- [178] Yao Y, Chinnici C, Tang H, Trojanowski JQ, Lee VM and Pratico D. Brain inflammation and oxidative stress in a transgenic mouse model of Alzheimer-like brain amyloidosis. *J Neuroinflammation* 2004; 1: 21.
- [179] Sano M, Ernesto C, Thomas RG, Klauber MR, Schafer K, Grundman M, Woodbury P, Growdon J, Cotman CW, Pfeiffer E, Schneider LS and Thal LJ. A controlled trial of selegiline, alpha-tocopherol, or both as treatment for Alzheimer's disease. The Alzheimer's Disease Cooperative Study. *N Engl J Med* 1997; 336: 1216-1222.
- [180] Zandi PP, Anthony JC, Khachaturian AS, Stone SV, Gustafson D, Tschanz JT, Norton MC, Welsh-Bohmer KA, Breitner JC; Cache County Study Group. Reduced risk of Alzheimer disease in users of antioxidant vitamin supplements: the Cache County Study. *Arch Neurol* 2004; 61: 82-88.
- [181] Dysken MW, Sano M, Asthana S, Vertrees JE, Pallaki M, Llorente M, Love S, Schellenberg GD, McCarten JR, Malphurs J, Prieto S, Chen P, Loreck DJ, Trapp G, Bakshi RS, Mintzer JE, Heidebrink JL, Vidal-Cardona A, Arroyo LM, Cruz AR, Zachariah S, Kowall NW, Chopra MP, Craft S, Thielke S, Turvey CL, Woodman C, Monnell KA, Gordon K, Tomaska J, Segal Y, Peduzzi PN and Guarino PD. Effect of vitamin E and memantine on functional decline in Alzheimer disease: the TEAM-AD VA cooperative randomized trial. *JAMA* 2014; 311: 33-44.
- [182] Petersen RC, Thomas RG, Grundman M, Bennett D, Doody R, Ferris S, Galasko D, Jin S, Kaye J, Levey A, Pfeiffer E, Sano M, van Dyck CH, Thal LJ; Alzheimer's Disease Cooperative Study Group. Vitamin E and donepezil for the treatment of mild cognitive impairment. *N Engl J Med* 2005; 352: 2379-2388.
- [183] Lee J, Boo JH and Ryu H. The failure of mitochondria leads to neurodegeneration: Do mitochondria need a jump start? *Adv Drug Deliv Rev* 2009; 61: 1316-1323.
- [184] Gutzmann H and Hadler D. Sustained efficacy and safety of idebenone in the treatment of

Therapeutics attenuating A β in Alzheimer's disease

- Alzheimer's disease: update on a 2-year double-blind multicentre study. *J Neural Transm Suppl* 1998; 54: 301-310.
- [185] Thal LJ, Grundman M, Berg J, Ernstrom K, Margolin R, Pfeiffer E, Weiner MF, Zamrini E and Thomas RG. Idebenone treatment fails to slow cognitive decline in Alzheimer's disease. *Neurology* 2003; 61: 1498-1502.
- [186] Siedlak SL, Casadesus G, Webber KM, Pappolla MA, Atwood CS, Smith MA and Perry G. Chronic antioxidant therapy reduces oxidative stress in a mouse model of Alzheimer's disease. *Free Radic Res* 2009; 43: 156-164.
- [187] Pocernich CB, Lange ML, Sultana R and Butterfield DA. Nutritional approaches to modulate oxidative stress in Alzheimer's disease. *Curr Alzheimer Res* 2011; 8: 452-469.
- [188] Shinto L, Quinn J, Montine T, Dodge HH, Woodward W, Baldauf-Wagner S, Waichunas D, Bumgarner L, Bourdette D, Silbert L and Kaye J. A randomized placebo-controlled pilot trial of omega-3 fatty acids and alpha lipoic acid in Alzheimer's disease. *J Alzheimers Dis* 2014; 38: 111-120.
- [189] Ono K, Hirohata M and Yamada M. Alpha-lipoic acid exhibits anti-amyloidogenicity for beta-amyloid fibrils in vitro. *Biochem Biophys Res Commun* 2006; 341: 1046-1052.
- [190] McManus MJ, Murphy MP and Franklin JL. The mitochondria-targeted antioxidant MitoQ prevents loss of spatial memory retention and early neuropathology in a transgenic mouse model of Alzheimer's disease. *J Neurosci* 2011; 31: 15703-15715.
- [191] Kapay NA, Isaev NK, Stelmashook EV, Popova OV, Zorov DB, Skrebitsky VG and Skulachev VP. In vivo injected mitochondria-targeted plastoquinone antioxidant SkQR1 prevents beta-amyloid-induced decay of long-term potentiation in rat hippocampal slices. *Biochemistry (Mosc)* 2011; 76: 1367-1370.
- [192] Calkins MJ, Manczak M, Mao P, Shirendeb U and Reddy PH. Impaired mitochondrial biogenesis, defective axonal transport of mitochondria, abnormal mitochondrial dynamics and synaptic degeneration in a mouse model of Alzheimer's disease. *Hum Mol Genet* 2011; 20: 4515-4529.
- [193] Chernyak BV, Antonenko YN, Domnina LV, Ivanova OY, Lyamzaev KG, Pustovidko AV, Rokitskaya TI, Severina II, Simonyan RA, Trendeleva TA and Zvyagil'skaya RA. Novel penetrating cations for targeting mitochondria. *Curr Pharm Des* 2013; 19: 2795-2806.
- [194] Anand R, Gill KD and Mahdi AA. Therapeutics of Alzheimer's disease: Past, present and future. *Neuropharmacology* 2014; 76 Pt A: 27-50.
- [195] Robinson RA, Joshi G, Huang Q, Sultana R, Baker AS, Cai J, Pierce W, St Clair DK, Marksbury WR and Butterfield DA. Proteomic analysis of brain proteins in APP/PS-1 human double mutant knock-in mice with increasing amyloid beta-peptide deposition: insights into the effects of in vivo treatment with N-acetylcysteine as a potential therapeutic intervention in mild cognitive impairment and Alzheimer's disease. *Proteomics* 2011; 11: 4243-4256.
- [196] Butterfield DA and Poon HF. The senescence-accelerated prone mouse (SAMP8): a model of age-related cognitive decline with relevance to alterations of the gene expression and protein abnormalities in Alzheimer's disease. *Exp Gerontol* 2005; 40: 774-783.
- [197] Parachikova A, Green KN, Hendrix C and LaFerla FM. Formulation of a medical food cocktail for Alzheimer's disease: beneficial effects on cognition and neuropathology in a mouse model of the disease. *PLoS One* 2010; 5: e14015.
- [198] Javed H, Khan MM, Ahmad A, Vaibhav K, Ahmad ME, Khan A, Ashafaq M, Islam F, Siddiqui MS, Safhi MM and Islam F. Rutin prevents cognitive impairments by ameliorating oxidative stress and neuroinflammation in rat model of sporadic dementia of Alzheimer type. *Neuroscience* 2012; 210: 340-352.
- [199] Vijayapadma V, Ramyaa P, Pavithra D and Krishnasamy R. Protective effect of lutein against benzo(a)pyrene-induced oxidative stress in human erythrocytes. *Toxicol Ind Health* 2014; 30: 284-293.
- [200] Yang F, Lim GP, Begum AN, Ubada OJ, Simmons MR, Ambegaokar SS, Chen PP, Kayed R, Glabe CG, Frautschy SA and Cole GM. Curcumin inhibits formation of amyloid beta oligomers and fibrils, binds plaques, and reduces amyloid in vivo. *J Biol Chem* 2005; 280: 5892-5901.
- [201] Ringman JM, Frautschy SA, Teng E, Begum AN, Bardens J, Beigi M, Gylys KH, Badmaev V, Heath DD, Apostolova LG, Porter V, Vanek Z, Marshall GA, Helleman G, Sugar C, Masterman DL, Montine TJ, Cummings JL and Cole GM. Oral curcumin for Alzheimer's disease: tolerability and efficacy in a 24-week randomized, double blind, placebo-controlled study. *Alzheimers Res Ther* 2012; 4: 43.
- [202] Joseph JA, Denisova NA, Arendash G, Gordon M, Diamond D, Shukitt-Hale B and Morgan D. Blueberry supplementation enhances signaling and prevents behavioral deficits in an Alzheimer disease model. *Nutr Neurosci* 2003; 6: 153-162.
- [203] Ho L, Chen LH, Wang J, Zhao W, Talcott ST, Ono K, Teplow D, Humala N, Cheng A, Percival SS, Ferruzzi M, Janle E, Dickstein DL and Pasinetti GM. Heterogeneity in red wine polyphenolic contents differentially influences Alzheimer's disease-type neuropathology and cognitive deterioration. *J Alzheimers Dis* 2009; 16: 59-72.

Therapeutics attenuating A β in Alzheimer's disease

- [204] Orgogozo JM, Dartigues JF, Lafont S, Letenneur L, Commenges D, Salamon R, Renaud S and Breteler MB. Wine consumption and dementia in the elderly: a prospective community study in the Bordeaux area. *Rev Neurol (Paris)* 1997; 153: 185-192.
- [205] Zhou J, Zhang S, Zhao X and Wei T. Melatonin impairs NADPH oxidase assembly and decreases superoxide anion production in microglia exposed to amyloid-beta1-42. *J Pineal Res* 2008; 45: 157-165.
- [206] He P, Ouyang X, Zhou S, Yin W, Tang C, Laudon M and Tian S. A novel melatonin agonist Neu-P11 facilitates memory performance and improves cognitive impairment in a rat model of Alzheimer' disease. *Horm Behav* 2013; 64: 1-7.
- [207] Lu P, Mamiya T, Lu LL, Mouri A, Zou L, Nagai T, Hiramatsu M, Ikejima T and Nabeshima T. Silibinin prevents amyloid beta peptide-induced memory impairment and oxidative stress in mice. *Br J Pharmacol* 2009; 157: 1270-1277.
- [208] Stackman RW, Eckenstein F, Frei B, Kulhanek D, Nowlin J and Quinn JF. Prevention of age-related spatial memory deficits in a transgenic mouse model of Alzheimer's disease by chronic Ginkgo biloba treatment. *Exp Neurol* 2003; 184: 510-520.
- [209] Prasanthi JR, Dasari B, Marwarha G, Larson T, Chen X, Geiger JD and Ghribi O. Caffeine protects against oxidative stress and Alzheimer's disease-like pathology in rabbit hippocampus induced by cholesterol-enriched diet. *Free Radic Biol Med* 2010; 49: 1212-1220.
- [210] Cao C, Cirrito JR, Lin X, Wang L, Verges DK, Dickson A, Mamcarz M, Zhang C, Mori T, Arendash GW, Holtzman DM and Potter H. Caffeine suppresses amyloid-beta levels in plasma and brain of Alzheimer's disease transgenic mice. *J Alzheimers Dis* 2009; 17: 681-697.
- [211] Kim TI, Lee YK, Park SG, Choi IS, Ban JO, Park HK, Nam SY, Yun YW, Han SB, Oh KW and Hong JT. I-Theanine, an amino acid in green tea, attenuates beta-amyloid-induced cognitive dysfunction and neurotoxicity: reduction in oxidative damage and inactivation of ERK/p38 kinase and NF-kappaB pathways. *Free Radic Biol Med* 2009; 47: 1601-1610.
- [212] Iuvone T, De Filippis D, Esposito G, D'Amico A and Izzo AA. The spice sage and its active ingredient rosmarinic acid protect PC12 cells from amyloid-beta peptide-induced neurotoxicity. *J Pharmacol Exp Ther* 2006; 317: 1143-1149.
- [213] Zhao BL, Li XJ, He RG, Cheng SJ and Xin WJ. Scavenging effect of extracts of green tea and natural antioxidants on active oxygen radicals. *Cell Biophys* 1989; 14: 175-185.
- [214] Goodman Y, Steiner MR, Steiner SM and Mattson MP. Nordihydroguaiaretic acid protects hippocampal neurons against amyloid beta-peptide toxicity, and attenuates free radical and calcium accumulation. *Brain Res* 1994; 654: 171-176.
- [215] Ramsey CP, Glass CA, Montgomery MB, Lindl KA, Ritson GP, Chia LA, Hamilton RL, Chu CT and Jordan-Sciutto KL. Expression of Nrf2 in neurodegenerative diseases. *J Neuropathol Exp Neurol* 2007; 66: 75-85.
- [216] Dumont M, Stack C, Elipenhli C, Jainuddin S, Gerges M, Starkova N, Calingasan NY, Yang L, Tampellini D, Starkov AA, Chan RB, Di Paolo G, Pujol A and Beal MF. Bezafibrate administration improves behavioral deficits and tau pathology in P301S mice. *Hum Mol Genet* 2012; 21: 5091-5105.
- [217] Kanninen K, Malm TM, Jyrkkanen HK, Goldsteins G, Keksa-Goldsteine V, Tanila H, Yamamoto M, Yla-Herttuala S, Levonen AL and Koistinaho J. Nuclear factor erythroid 2-related factor 2 protects against beta amyloid. *Mol Cell Neurosci* 2008; 39: 302-313.
- [218] Ehehalt R, Keller P, Haass C, Thiele C and Simons K. Amyloidogenic processing of the Alzheimer beta-amyloid precursor protein depends on lipid rafts. *J Cell Biol* 2003; 160: 113-123.
- [219] Wong BX, Hung YH, Bush AI and Duce JA. Metals and cholesterol: two sides of the same coin in Alzheimer's disease pathology. *Front Aging Neurosci* 2014; 6: 91.
- [220] Pierrot N, Tyteca D, D'Auria L, Dewachter I, Gailly P, Hendrickx A, Tasiaux B, Haylani LE, Muls N, N'Kuli F, Laquerriere A, Demoulin JB, Champion D, Brion JP, Courtroy PJ, Kienlen-Campard P and Octave JN. Amyloid precursor protein controls cholesterol turnover needed for neuronal activity. *EMBO Mol Med* 2013; 5: 608-625.
- [221] Mukherjee P, Faber AC, Shelton LM, Baek RC, Chiles TC and Seyfried TN. Thematic review series: sphingolipids. Ganglioside GM3 suppresses the proangiogenic effects of vascular endothelial growth factor and ganglioside GD1a. *J Lipid Res* 2008; 49: 929-938.
- [222] Hicks DA, Nalivaeva NN and Turner AJ. Lipid rafts and Alzheimer's disease: protein-lipid interactions and perturbation of signaling. *Front Physiol* 2012; 3: 189.
- [223] Liu CC, Kanekiyo T, Xu H and Bu G. Apolipoprotein E and Alzheimer disease: risk, mechanisms and therapy. *Nat Rev Neurol* 2013; 9: 106-118.
- [224] Zhang H, Wu LM and Wu J. Cross-talk between apolipoprotein E and cytokines. *Mediators Inflamm* 2011; 2011: 949072.
- [225] Takeda S, Sato N and Morishita R. Systemic inflammation, blood-brain barrier vulnerability

Therapeutics attenuating A β in Alzheimer's disease

- and cognitive/non-cognitive symptoms in Alzheimer disease: relevance to pathogenesis and therapy. *Front Aging Neurosci* 2014; 6: 171.
- [226] Phan HT, Hata T, Morita M, Yoda T, Hamada T, Vestergaard MC and Takagi M. The effect of oxysterols on the interaction of Alzheimer's amyloid beta with model membranes. *Biochim Biophys Acta* 2013; 1828: 2487-2495.
- [227] Vatassery GT, Smith WE, Quach HT and Lai JC. In vitro oxidation of vitamin E, vitamin C, thiols and cholesterol in rat brain mitochondria incubated with free radicals. *Neurochem Int* 1995; 26: 527-535.
- [228] Arciello M, Gori M and Balsano C. Mitochondrial dysfunctions and altered metals homeostasis: new weapons to counteract HCV-related oxidative stress. *Oxid Med Cell Longev* 2013; 2013: 971024.
- [229] Shinohara M, Sato N, Shimamura M, Kurinami H, Hamasaki T, Chatterjee A, Rakugi H and Morishita R. Possible modification of Alzheimer's disease by statins in midlife: interactions with genetic and non-genetic risk factors. *Front Aging Neurosci* 2014; 6: 71.
- [230] Kou J, Song M, Pattanayak A, Lim JE, Yang J, Cao D, Li L and Fukuchi K. Combined treatment of Abeta immunization with statin in a mouse model of Alzheimer's disease. *J Neuroimmunol* 2012; 244: 70-83.
- [231] Buxbaum JD, Cullen EI and Friedhoff LT. Pharmacological concentrations of the HMG-CoA reductase inhibitor lovastatin decrease the formation of the Alzheimer beta-amyloid peptide in vitro and in patients. *Front Biosci* 2002; 7: a50-59.
- [232] Friedhoff LT, Cullen EI, Geoghegan NS and Buxbaum JD. Treatment with controlled-release lovastatin decreases serum concentrations of human beta-amyloid (A beta) peptide. *Int J Neuropsychopharmacol* 2001; 4: 127-130.
- [233] Zhang LL, Sui HJ, Liang B, Wang HM, Qu WH, Yu SX and Jin Y. Atorvastatin prevents amyloid-beta peptide oligomer-induced synaptotoxicity and memory dysfunction in rats through a p38 MAPK-dependent pathway. *Acta Pharmacol Sin* 2014; 35: 716-726.
- [234] Cordle A and Landreth G. 3-Hydroxy-3-methylglutaryl-coenzyme A reductase inhibitors attenuate beta-amyloid-induced microglial inflammatory responses. *J Neurosci* 2005; 25: 299-307.
- [235] Vaughan CJ and Delanty N. Neuroprotective properties of statins in cerebral ischemia and stroke. *Stroke* 1999; 30: 1969-1973.
- [236] Butterfield DA, Barone E and Mancuso C. Cholesterol-independent neuroprotective and neurotoxic activities of statins: perspectives for statin use in Alzheimer disease and other age-related neurodegenerative disorders. *Pharmacol Res* 2011; 64: 180-186.
- [237] Barone E, Di Domenico F and Butterfield DA. Statins more than cholesterol lowering agents in Alzheimer disease: their pleiotropic functions as potential therapeutic targets. *Biochem Pharmacol* 2014; 88: 605-616.
- [238] Mancuso C, Santangelo R and Calabrese V. The heme oxygenase/biliverdin reductase system: a potential drug target in Alzheimer's disease. *J Biol Regul Homeost Agents* 2013; 27: 75-87.
- [239] Ehret MJ and Chamberlin KW. Current Practices in the Treatment of Alzheimer Disease: Where is the Evidence After the Phase III Trials? *Clin Ther* 2015; 37: 1604-1616.