

Review Article

Height and risk of melanoma: a systematic review and meta-analysis

Dao-Jiang Yu^{1*}, Xiu-Jie Li^{1*}, Anne Morice², Li-Jun Wu¹, Wei Sun¹, Tian-Lan Zhao¹

¹Department of Plastic Surgery, The Second Affiliated Hospital of Soochow University, No.1055, Sanxiang Road, Suzhou, Jiangsu Province, China; ²Department of Plastic and Maxilla-Facial Surgery, Necker Children Hospital, No.149 Serves Street 75015, Paris, France. *Equal contributors.

Received May 27, 2017; Accepted March 28, 2018; Epub May 15, 2018; Published May 30, 2018

Abstract: This systematic review and meta-analysis of prospective cohort studies aimed to comprehensively estimate the relationship between height and melanoma risk. Prospective cohort studies were identified through literature search in Pubmed and EMBASE. The summary relative risk (RR) with 95% confidence interval (95% CI) was calculated using random-effects meta-analysis. Twelve studies with a total of 4,723,739 participants and 20,049 melanoma cases were included. When compared with individuals with the lowest category of height, individuals with the top category of height had increased risk of melanoma (Random-effects RR = 1.46, 95% CI 1.24 to 1.73; $P < 0.001$). Per 10-cm increment in the height was positively correlated with increased melanoma risk (Random-effects RR = 1.27, 95% CI 1.19 to 1.35; $P < 0.001$). Subgroup analysis by gender showed that RRs of melanoma risk associated with per 10-cm increment in height in women and men were 1.27 (95% CI 1.18 to 1.36; $P < 0.001$) and 1.52 (95% CI 1.01 to 2.29; $P = 0.04$), respectively. There is strong evidence for the relationship between height and melanoma risk.

Keywords: Height, melanoma, association, risk factor

Introduction

Melanoma is a common malignant disease occurring in the melanocytes [1, 2]. Melanoma accounts for only about 2% of all skin cancers, but it the main cause of deaths associated with skin cancer [2, 3]. There is also an obvious increase in the incidence of melanoma in recent years [4]. There were more than 76,000 estimated newly diagnosed cases of melanoma and more than 10,000 estimated deaths caused by melanoma in United States in 2016 [3]. Ultraviolet light or sunlight exposure has been recognized as the main risk factor of melanoma, and several other risk factors are also reported, such as fair complexion, family history, and obesity [5-9]. However, the etiology of melanoma remains largely unknown, and more risk factors associated with melanoma need to be identified, which may be helpful for the prevention and screening of melanoma [6, 10-13].

Human height is considered to be determined by both genetic and environmental factors, and

it is indeed associated with circulating insulin-like growth factor I (IGF-1) levels during childhood [14-16]. Both experimental and epidemiologic data have suggest that high IGF-1 level is intensively correlated with elevated risk of cancer [17-21]. Some studies demonstrated an association between height and risk of several types of cancer, such as breast cancer and colorectal cancer [22-27]. In contrast, the results of epidemiological studies investigating the relationship between height and melanoma were controversial and inconclusive. Some studies reported that higher height was a risk factor of melanoma, but other studies didn't. We therefore conducted this meta-analysis.

Materials and methods

Literature search and selection of eligible studies

Literature search was performed in Pubmed and EMBASE databases up to Oct 20, 2016. The following search strategy was used: (mela-

Is height a risk factor for melanoma?

Table 1. Characteristics of prospective studies on height and melanoma risk

Study	Location (Study period)	Study design	Participants	Follow-up	Number of cases	Gender (female/male)	Age (median, range) years	Level of height	Adjustment for covariates	Study quality
Thune I 1993 [42]	Norway (1963-1975)	Prospective cohort study	1.3 million individuals	17.6 years	4958	2814/2144	NA	H vs L	Age, birth cohort, geographic region and BMI.	8
Veierod MB 1997 [43]	Norway (1977-1983)	Prospective cohort study	25,708 men and 25,049 women aged 16-56 years	12.4 years	108	61/47	52 (29 - 64)	H vs L	County of residence, gender, age at inclusion and attained age.	8
Freedman DM 2003 [34]	USA (1983-1989)	Prospective cohort study	68,588 white subjects	10 years	207	159/48	NA	H vs L	Age, weight, gender, alcohol intake, smoking, skin pigmentation, hair color, personal history of non-melanoma skin cancer, decade began work as a technologist, education, and proxy measures for residential childhood and adult sunlight exposure.	8
Batty GD 2010 [33]	Asia Pacific Cohort Studies Collaboration	Prospective cohort study	506648 study participants	6.5 years	88	25/63	NA	Per 6 cm increment	Age, study, and year of birth.	8
Green J 2011 [35]	UK (1996-2001)	Prospective cohort study	1297124 women	9.4 years	3583	NA	NA	Per 10 cm increment	Age, region, socioeconomic status, smoking, alcohol intake, body-mass index, strenuous exercise, age at menarche, parity, and age at first birth.	9
Kabat GC 2013 [37]	Canada (1980-1985)	Prospective cohort study	90,000 women	16.2 years	327	NA	NA	Per 10 cm increment	Age at entry, menopausal status, BMI, and years of education.	9
Kabat G 2013 [36]	USA (1993-1998)	Prospective cohort study	144,701 women participating in the Women's Health Initiative	12 years	1169	NA	NA	Per 10 cm increment	Age, alcohol, pack-years, hormone therapy, education, ethnicity, BMI, and randomization status.	9
Kvaskoff M 2014 [39]	France (1989-1991)	Prospective cohort study	98,995 women	18 years	589	0/589	NA	H vs L	Age, number of naevi, freckling, skin and hair colour, skin sensitivity to sun exposure, residential sun exposure, and physical activity.	9
Kabat GC 2014 [38]	USA (1995-1996)	Prospective cohort study	288,683 men and 192,514 women enrolled in the National Institutes of Health-AARP Diet and Health Study	10.5 years	4780	1224/3556	NA	Per 10 cm increment	Age at entry, education, race, smoking status, and body mass index.	9
Wiren S 2014 [44]	Austria, Norway, and Sweden	Prospective cohort study	585,928 participants from seven cohorts	12.7 years	1989	NA	NA	Per 5 cm increment	Date of birth and age at health examination, and stratified for sub-cohort within the model.	9

Is height a risk factor for melanoma?

Lahmann PH 2016 [40]	Australia (1992-1996)	Prospective cohort study	1171 Australian men and women	16 years	28	17/11	NA	Per 5 cm increment	Age, treatment allocation, skin cancer history, elastosis of the neck, and smoking status.	7
Meyle KD 2016 [41]	Denmark (1978)	Prospective cohort study	316,193 individuals from the Copenhagen School Health Records Register	29 years	2223	1205/1018	NA	Per height z-score	Birth cohort, age, and gender.	8

(BMI: body mass index; NA: not available).

Is height a risk factor for melanoma?




Figure 1. Individuals with the top categories of height had increased risk of melanoma.

noma or skin cancer) AND (height OR body size) AND (prospective or cohort or longitudinal). No limitation in language was used. Additional studies were found by looking through the references of retrieved reviews or articles.

Prospective cohort study which evaluated the association between height and melanoma risk and reported risk estimates of melanoma risk were included. Studies which failed to meet any item of the selection criteria above were excluded.

Data extraction

Extracted information included first author, country, follow-up time, adjusted confounding factors, and adjusted RRs with 95% CIs. Any differences in extracted data were settled by consensus. RRs of melanoma associated with the top categories of height or per 10-cm increment in height were used in the meta-analysis. For studies reported RRs of melanoma associated with per 5-cm or per 6-cm increment in height, the RRs of melanoma associated with per 10-cm in height were calculated through the following formula [28]: $RR_{\text{per 10-cm}} = RR_{\text{(original)}} \times \frac{\text{Increment (10)}}{\text{Increment (original)}}$. We assessed the studies' quality by Newcastle-Ottawa Scale (NOS) [29]. Studies with a quality score no less than 6 were considered as high-quality studies.

Statistical analysis

The RRs from included studies were pooled using random-effect model of DerSimonian-Lard method [30]. The RRs of melanoma associated with the top categories of height or per 10-cm increment in height were pooled separately. Heterogeneity was evaluated with the I^2 statistic, and I^2 more than 50% showed obvious heterogeneity between studies [31]. Sensitivity analysis was then performed by excluding single study by turns. Subgroup analysis was performed by gender. To qualitatively assess publication bias, both funnel plot and Egger's test was utilized [32].

Results

Characteristics of included studies

Of 359 abstracts identified from PubMed and EMBASE databases, 24 studies were firstly included and assessed for final inclusion by reading full-texts [33-44]. Twelve studies were then excluded [45-55], and the left 12 prospective studies were finally included in the current meta-analysis [33-44]. Batty et al's study reported a prospective study of participants from 38 cohort studies [33], and Wiren et al's study reported a prospective study of participants from seven cohort studies [44]. Either

Is height a risk factor for melanoma?




Figure 2. Increased height (per 10-cm increment) was positively associated with an increased melanoma risk.

Batty et al's study or Wiren et al's study was regarded as one prospective study in the meta-analysis. Overall, 12 prospective cohort studies contained a total of 4,723,739 participants and 20,049 with cases melanoma [33-44].

Table 1 summarized the characteristics of included studies (**Table 1**). Most studies were performed in Caucasian populations, such as USA, UK, Canada, and Norway (**Table 1**). Four studies reported RRs of melanoma associated with the top categories of height compared with the lowest category of height, and the other 8 studies reported RRs of melanoma associated with per certain increment in height (**Table 1**). All included studies performed multivariate regression analyses by adjusting for potential confounding factors, but the factors varied obviously across different studies. The number of participants ranged from 1171 to 1.3 million, while the time of follow-up ranged from 6.5 years to 25 years (**Table 1**). All studies had a NOS score of more than 6 points, and thus had high-quality.

Meta-analysis

When compared with individuals with the lowest categories of height, individuals with the

top categories of height had increased risk of melanoma (Random-effects model, $I^2 = 60.9\%$, $P = 0.029$; $RR = 1.46$, 95% CI 1.24 to 1.73; $P < 0.001$) (**Figure 1**). Increased height (per 10-cm increment) was positively associated with an increased melanoma risk (Random-effects model, $I^2 = 64.6\%$, $P = 0.002$; $RR = 1.27$, 95% CI 1.19 to 1.35; $P < 0.001$) (**Figure 2**). Subgroup analysis by gender showed that RRs of melanoma risk associated with per 10-cm increment in height were 1.27 (95% CI 1.18 to 1.36; $P < 0.001$; $I^2 = 59.5\%$, $P = 0.016$) in female and 1.52 (95% CI 1.01 to 2.29; $P = 0.04$; $I^2 = 72.8\%$, $P = 0.025$) in male, respectively (**Figure 3**).

There was also no obvious asymmetry in the funnel plot (**Figure 4**), and the P value of Egger's regression test was 0.27, which indicated lack of publication bias in the meta-analysis.

Discussion

The findings in the meta-analysis suggest an obvious association between height and melanoma risk. IGF-1 is a hormone which has key roles in childhood growth and many metabolic effects in adults [56, 57]. IGF-1 mainly produces its by binding to the insulin-like growth factor 1 receptor (IGF1R), which has been found on

Is height a risk factor for melanoma?




Figure 3. Increased height (per 10-cm increment) was positively associated with an increased melanoma risk in both women and men.




Figure 4. Funnel plot in the meta-analysis of studies reporting RRs of melanoma associated with per certain increment of height.

the cells of many tissues [56, 58-60]. Besides, IGF signaling pathway has been shown to have key roles in the development and progression

of many cancers [18, 59, 61]. Both experimental and epidemiologic data have suggest that high IGF-1 level is intensively correlated with elevated risk of cancer, and it has long been regarded as a tumor-promoting factor [18, 59, 61, 62]. Human height is intensively associated with circulating IGF-1 levels during childhood. Children with higher levels of circulating IGF-1 will grow taller than those with lower levels of circulating IGF-1 [63]. On the contrary, individuals with taller height would had higher levels of circulating IGF-1, and these individuals exposing to higher lev-

els of circulating IGF-1 and therefore suffer from increased risk of diseases caused by higher levels of IGF-1 [63-65]. Therefore, there is

Is height a risk factor for melanoma?

also a possible association between height and melanoma risk. However, the above association has not been directly supported by data from experimental studies. In addition, an epidemiologic study even reported an inverse relationship between IGF-1 concentration and melanoma risk [50]. Therefore, the exact biologic mechanisms underlying the association between height and melanoma risk is still largely unclear, and further studies exploring the mechanisms underlying the relationship between height and melanoma risk are also needed.

There were also several limitations in this study. There was obvious heterogeneity among those included studies. The heterogeneity may come from the different sample size, various adjusted confounding factors, and different demographic characteristics of participants. However, all included studies provided adjusted risk estimates, which could decrease the risk of bias caused by the difference in the design of included studies. Secondly, most studies were carried out in European countries and North America. The conclusion in the meta-analysis could not be generalized to Asians or Africans. More cohort studies from non-Caucasians are necessary to further elucidate the association. Thirdly, the present study only considered published literatures, and it's possible for the existence of publication bias which could affect the exact impact of height on melanoma risk. However, neither funnel plot nor Egger's test showed evidence of publication bias. Finally, IGF-1 has been considered as an important factor underlying the biologic mechanism underlying the association between height and melanoma risk. However, it's still unclear whether height can still increase the risk of melanoma after adjusting for IGF-1 levels in human bodies.

In summary, there is strong evidence for the relationship between height and melanoma risk.

Acknowledgements

This research was supported by Research and Innovation Project for College Graduates of Jiangsu Province [SJLX15_0581]; Guidance Program of Municipal Science and Technology of Suzhou [SYSD2015092]; Clinical Application Projects of Technology Research and Demon-

stration of Jiangsu [BL2014041]; Natural Science Foundation of Jiangsu Province [BK2017-0354].

Disclosure of conflict of interest

None.

Address correspondence to: Tian-Lan Zhao, Department of Plastic Surgery, The Second Affiliated Hospital of Soochow University, No.1055, Sanxiang Road, Suzhou 215004, Jiangsu Province, China. Tel: +86-0512-83986538; E-mail: zhaotlps@sohu.com

References

- [1] Wick MR. Cutaneous melanoma: a current overview. *Semin Diagn Pathol* 2016; 33: 225-241.
- [2] Eggermont AM, Spatz A and Robert C. Cutaneous melanoma. *Lancet* 2014; 383: 816-827.
- [3] Siegel RL, Miller KD and Jemal A. Cancer statistics, 2016. *CA Cancer J Clin* 2016; 66: 7-30.
- [4] Simard EP, Ward EM, Siegel R and Jemal A. Cancers with increasing incidence trends in the United States: 1999 through 2008. *CA Cancer J Clin* 2012; 62: 118-128.
- [5] Sergentanis TN, Antoniadis AG, Gogas HJ, Antonopoulos CN, Adami HO, Ekblom A and Petridou ET. Obesity and risk of malignant melanoma: a meta-analysis of cohort and case-control studies. *Eur J Cancer* 2013; 49: 642-657.
- [6] Jiang AJ, Rambhatla PV and Eide MJ. Socioeconomic and lifestyle factors and melanoma: a systematic review. *Br J Dermatol* 2015; 172: 885-915.
- [7] Praestegaard C, Kjaer SK, Christensen J, Tjonneland A, Halkjaer J and Jensen A. Obesity and risks for malignant melanoma and non-melanoma skin cancer: results from a large Danish prospective cohort study. *J Invest Dermatol* 2015; 135: 901-904.
- [8] Caini S, Cattaruzza S, Bendinelli B, Tosti G, Masala G, Gnagnarella P, Assedi M, Stanganelli I, Palli D and Gandini S. Coffee, tea and caffeine intake and the risk of non-melanoma skin cancer: a review of the literature and meta-analysis. *Eur J Nutr* 2017; 56: 1-12.
- [9] Lindqvist PG, Epstein E, Nielsen K, Landin-Olsson M, Ingvar C and Olsson H. Avoidance of sun exposure as a risk factor for major causes of death: a competing risk analysis of the Melanoma in Southern Sweden cohort. *J Intern Med* 2016; 280: 375.
- [10] Berwick M, Buller DB, Cust A, Gallagher R, Lee TK, Meyskens F, Pandey S, Thomas NE, Veierod MB and Ward S. Melanoma epidemiology and prevention. *Cancer Treat Res* 2016; 167: 17-49.

Is height a risk factor for melanoma?

- [11] Demirsoy S, Martin S, Maes H and Agostinis P. Adapt, recycle, and move on: proteostasis and trafficking mechanisms in melanoma. *Front Oncol* 2016; 6: 240.
- [12] Hawkes JE, Truong A and Meyer LJ. Genetic predisposition to melanoma. *Semin Oncol* 2016; 43: 591-597.
- [13] Tripp MK, Watson M, Balk SJ, Swetter SM and Gershenwald JE. State of the science on prevention and screening to reduce melanoma incidence and mortality: the time is now. *CA Cancer J Clin* 2016; [Epub ahead of print].
- [14] Bäckeljauw PF, Kuntze J, Frane J, Calikoglu AS and Chernausek SD. Adult and near-adult height in patients with severe insulin-like growth factor-I deficiency after long-term therapy with recombinant human insulin-like growth factor-I. *Horm Res Paediatr* 2013; 80: 47-56.
- [15] Kristrom B, Lundberg E, Jonsson B and Albertsson-Wikland K. IGF-1 and growth response to adult height in a randomized GH treatment trial in short non-GH-deficient children. *J Clin Endocrinol Metab* 2014; 99: 2917-2924.
- [16] Bogin B, Hermanussen M, Blum WF and Assmann C. Sex, sport, IGF-1 and the community effect in height hypothesis. *Int J Environ Res Public Health* 2015; 12: 4816-4832.
- [17] Gunnell D. Height, insulin-like growth factors and cancer risk. *Growth Horm IGF Res* 2000; 10 Suppl A: S39-40.
- [18] Rota LM and Wood TL. Crosstalk of the insulin-like growth factor receptor with the wnt signaling pathway in breast cancer. *Front Endocrinol (Lausanne)* 2015; 6: 92.
- [19] Werner H, Sarfstein R, LeRoith D and Bruchim I. Insulin-like growth factor 1 signaling axis meets p53 genome protection pathways. *Front Oncol* 2016; 6: 159.
- [20] Gunnell D, Okasha M, Smith GD, Oliver SE, Sandhu J and Holly JM. Height, leg length, and cancer risk: a systematic review. *Epidemiol Rev* 2001; 23: 313-342.
- [21] Ochnik AM and Baxter RC. Combination therapy approaches to target insulin-like growth factor receptor signaling in breast cancer. *Endocr Relat Cancer* 2016; 23: R513-R536.
- [22] Aune D, Vieira AR, Chan DS, Navarro Rosenblatt DA, Vieira R, Greenwood DC, Cade JE, Burley VJ and Norat T. Height and pancreatic cancer risk: a systematic review and meta-analysis of cohort studies. *Cancer Causes Control* 2012; 23: 1213-1222.
- [23] Schouten LJ, Rivera C, Hunter DJ, Spiegelman D, Adami HO, Arslan A, Beeson WL, van den Brandt PA, Buring JE, Folsom AR, Fraser GE, Freudenheim JL, Goldbohm RA, Hankinson SE, Lacey JV Jr, Leitzmann M, Lukanova A, Marshall JR, Miller AB, Patel AV, Rodriguez C, Rohan TE, Ross JA, Wolk A, Zhang SM and Smith-Warner SA. Height, body mass index, and ovarian cancer: a pooled analysis of 12 cohort studies. *Cancer Epidemiol Biomarkers Prev* 2008; 17: 902-912.
- [24] Boursi B, Haynes K, Mamtani R and Yang YX. Height as an independent anthropomorphic risk factor for colorectal cancer. *Eur J Gastroenterol Hepatol* 2014; 26: 1422-1427.
- [25] Aarestrup J, Gamborg M, Cook MB and Baker JL. Childhood height increases the risk of prostate cancer mortality. *Eur J Cancer* 2015; 51: 1340-1345.
- [26] Aarestrup J, Gamborg M, Ulrich LG, Sorensen TI and Baker JL. Childhood body mass index and height and risk of histologic subtypes of endometrial cancer. *Int J Obes (Lond)* 2016; 40: 1096-1102.
- [27] Khankari NK, Shu XO, Wen W, Kraft P, Lindstrom S, Peters U, Schildkraut J, Schumacher F, Bofetta P, Risch A, Bickeboller H, Amos CI, Easton D, Eeles RA, Gruber SB, Haiman CA, Hunter DJ, Chanock SJ, Pierce BL and Zheng W. Association between adult height and risk of colorectal, lung, and prostate cancer: results from meta-analyses of prospective studies and Mendelian randomization analyses. *PLoS Med* 2016; 13: e1002118.
- [28] Shah AS, Langrish JP, Nair H, McAllister DA, Hunter AL, Donaldson K, Newby DE and Mills NL. Global association of air pollution and heart failure: a systematic review and meta-analysis. *Lancet* 2013; 382: 1039-1048.
- [29] Gláucia FC, Marcos RDS, Tatiani OF and Ana R. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomized studies. 2013;
- [30] DerSimonian R and Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7: 177-188.
- [31] Higgins JP and Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; 21: 1539-1558.
- [32] Egger M, Davey Smith G, Schneider M and Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; 315: 629-634.
- [33] Freedman DM, Sigurdson A, Doody MM, Rao RS and Linet MS. Risk of melanoma in relation to smoking, alcohol intake, and other factors in a large occupational cohort. *Cancer Causes Control* 2003; 14: 847-857.
- [34] Green J, Cairns BJ, Casabonne D, Wright FL, Reeves G and Beral V. Height and cancer incidence in the million women study: prospective cohort, and meta-analysis of prospective studies of height and total cancer risk. *Lancet Oncol* 2011; 12: 785-794.
- [35] Batty GD, Barzi F, Woodward M, Jamrozik K, Woo J, Kim HC, Ueshima H and Huxley RR. Adult height and cancer mortality in Asia: the

Is height a risk factor for melanoma?

- Asia Pacific Cohort Studies Collaboration. *Annals of Oncology* 2010; 21: 646-654.
- [36] Kabat GC, Anderson ML, Heo M, Hosgood HD, 3rd, Kamensky V, Bea JW, Hou L, Lane DS, Wactawski-Wende J, Manson JE and Rohan TE. Adult stature and risk of cancer at different anatomic sites in a cohort of postmenopausal women. *Cancer Epidemiol Biomarkers Prev* 2013; 22: 1353-1363.
- [37] Kabat GC, Heo M, Kamensky V, Miller AB and Rohan TE. Adult height in relation to risk of cancer in a cohort of Canadian women. *Int J Cancer* 2013; 132: 1125-1132.
- [38] Kabat GC, Kim MY, Hollenbeck AR and Rohan TE. Attained height, sex, and risk of cancer at different anatomic sites in the NIH-AARP diet and health study. *Cancer Causes Control* 2014; 25: 1697-1706.
- [39] Kvaskoff M, Bijon A, Mesrine S, Vilier A, Clavel-Chapelon F and Boutron-Ruault MC. Anthropometric features and cutaneous melanoma risk: a prospective cohort study in French women. *Cancer Epidemiol* 2014; 38: 357-363.
- [40] Lahmann PH, Hughes MC, Williams GM and Green AC. A prospective study of measured body size and height and risk of keratinocyte cancers and melanoma. *Cancer Epidemiol* 2016; 40: 119-125.
- [41] Meyle KD, Gamborg M, Holmich LR and Baker JL. Associations between childhood height and morphologically different variants of melanoma in adulthood. *Eur J Cancer* 2016; 67: 99-105.
- [42] Thune I, Olsen A, Albrektsen G and Tretli S. Cutaneous malignant melanoma: association with height, weight and body-surface area. a prospective study in Norway. *Int J Cancer* 1993; 55: 555-561.
- [43] Veierod MB, Thelle DS and Laake P. Diet and risk of cutaneous malignant melanoma: a prospective study of 50,757 Norwegian men and women. *Int J Cancer* 1997; 71: 600-604.
- [44] Wiren S, Haggstrom C, Ulmer H, Manjer J, Bjorge T, Nagel G, Johansen D, Hallmans G, Engeland A, Concin H, Jonsson H, Selmer R, Tretli S, Stocks T and Stattin P. Pooled cohort study on height and risk of cancer and cancer death. *Cancer Causes Control* 2014; 25: 151-159.
- [45] Hebert PR, Ajani U, Cook NR, Lee IM, Chan KS and Hennekens CH. Adult height and incidence of cancer in male physicians (United States). *Cancer Causes Control* 1997; 8: 591-597.
- [46] Feskanich D, Hunter DJ, Willett WC, Spiegelman D, Stampfer MJ, Speizer FE and Colditz GA. Oral contraceptive use and risk of melanoma in premenopausal women. *Br J Cancer* 1999; 81: 918-923.
- [47] Shors AR, Solomon C, McTiernan A and White E. Melanoma risk in relation to height, weight, and exercise (United States). *Cancer Causes Control* 2001; 12: 599-606.
- [48] Gallus S, Naldi L, Martin L, Martinelli M and La Vecchia C. Anthropometric measures and risk of cutaneous malignant melanoma: a case-control study from Italy. *Melanoma Res* 2006; 16: 83-87.
- [49] Olsen CM, Green AC, Zens MS, Stukel TA, Bataille V, Berwick M, Elwood JM, Gallagher R, Holly EA, Kirkpatrick C, Mack T, Osterlind A, Rosso S, Swerdlow AJ and Karagas MR. Anthropometric factors and risk of melanoma in women: a pooled analysis. *Int J Cancer* 2008; 122: 1100-1108.
- [50] Park SL, Setiawan VW, Kanetsky PA, Zhang ZF, Wilkens LR, Kolonel LN and Le Marchand L. Serum insulin-like growth factor-I and insulin-like growth factor binding protein-3 levels with risk of malignant melanoma. *Cancer Causes Control* 2011; 22: 1267-1275.
- [51] Walter RB, Brasky TM, Buckley SA, Potter JD and White E. Height as an explanatory factor for sex differences in human cancer. *J Natl Cancer Inst* 2013; 105: 860-868.
- [52] Hallberg O. Cancer and body height. *Pathophysiology* 2014; 21: 177-181.
- [53] Jiang Y, Marshall RJ, Walpole SC, Prieto-Merino D, Liu DX and Perry JK. An international ecological study of adult height in relation to cancer incidence for 24 anatomical sites. *Cancer Causes Control* 2015; 26: 493-499.
- [54] Ribero S, Glass D, Aviv A, Spector TD and Bataille V. Height and bone mineral density are associated with naevus count supporting the importance of growth in melanoma susceptibility. *PLoS One* 2015; 10: e0116863.
- [55] Li X, Liang L, Feng YA, De Vivo I, Giovannucci E, Tang JY and Han J. Height, height-related SNPs, and risk of non-melanoma skin cancer. *Br J Cancer* 2017; 116: 134-140.
- [56] Livingstone C and Borai A. Insulin-like growth factor-II: its role in metabolic and endocrine disease. *Clin Endocrinol (Oxf)* 2014; 80: 773-781.
- [57] Jing Z, Hou X, Wang Y, Yang G, Wang B, Tian X and Zhao S. Association between insulin-like growth factor-1 and cardiovascular disease risk: Evidence from a meta-analysis. *Int J Cardiol* 2015; 198: 1-5.
- [58] Tripaldi R, Stuppia L and Alberti S. Human height genes and cancer. *Biochimica Et Biophysica Acta* 2013; 1836: 27-41.
- [59] Danielpour D and Song K. Cross-talk between IGF-I and TGF-beta signaling pathways. *Cytokine Growth Factor Rev* 2006; 17: 59-74.
- [60] Cox OT, O'Shea S, Tresse E, Bustamante-Garrido M, Kiran-Deevi R and O'Connor R. IGF-1 receptor and adhesion signaling: an important axis in determining cancer cell phenotype and therapy resistance. *Front Endocrinol (Lausanne)* 2015; 6: 106.

Is height a risk factor for melanoma?

- [61] Klement RJ and Fink MK. Dietary and pharmacological modification of the insulin/IGF-1 system: exploiting the full repertoire against cancer. *Oncogenesis* 2016; 5: e193.
- [62] Vaziri-Gohar A and Houston KD. GPER1-mediated IGFBP-1 induction modulates IGF-1-dependent signaling in tamoxifen-treated breast cancer cells. *Mol Cell Endocrinol* 2016; 422: 160-171.
- [63] Janghorbani M, Momeni F and Dehghani M. Hip circumference, height and risk of type 2 diabetes: systematic review and meta-analysis. *Obes Rev* 2012; 13: 1172-1181.
- [64] Perkins JM, Subramanian SV, Smith GD and Özaltın E. Adult height, nutrition, and population health. *Nutrition Reviews* 2016; 74: 149.
- [65] Stefan N, Haring HU, Hu FB and Schulze MB. Divergent associations of height with cardio-metabolic disease and cancer: epidemiology, pathophysiology, and global implications. *Lancet Diabetes Endocrinol* 2016; 4: 457-467.