

Original Article

Comparison of sedative effects between dexmedetomidine and propofol in painless artificial abortion

Wenyong Peng, Wenlong Tu, Duoqia Xu, Huayan Lv, Danyan Zhu, Zhijian Lan

Department of Anesthesiology, Jinhua Municipal Central Hospital, Jinhua Hospital, Zhejiang University School of Medicine, Zhejiang Province, China

Received June 2, 2018; Accepted July 15, 2018; Epub September 15, 2018; Published September 30, 2018

Abstract: Objective: The aim of this study was to compare the sedative effects of dexmedetomidine and propofol in painless artificial abortion patients, discuss the safety and comfort of the anesthesia method, and to provide relevant evidence for clinical anesthesia of painless abortion. Methods: Using the random number table method, this study selected 200 patients that underwent abortions in the Department of Anesthesiology of Jinhua Municipal Central Hospital, from January 2015 to December 2017. American Society of Anesthesiologists (ASA) grades I to II were included, ages 18 to 40. A hundred cases of dexmedetomidine injections were used for induction and maintenance of anesthesia (Group D). A hundred cases of propofol injections were used for induction and maintenance of anesthesia (Group P). Heart rate (HR), mean arterial pressure (MAP), oxygen saturation (SpO₂), modified observer's assessment of alertness/sedation scale (MOAA/S), and incidence of respiratory depression, hypotension, sinus bradycardia, and other complications during surgery were compared between the two groups in T₀ (before anesthesia), T₁ (2 minutes after medication), T₂ (uterus dilatation), and T₃ (5 minutes after surgery), respectively. Results: There were no significant differences in MAP, HR, SpO₂, and MOAA/S scores between group D and group P at the time of T₀ (P>0.05). In group D, MAP and HR were lower than those in group P during T₁-T₃, but SpO₂ was significantly higher than that in group P (all P<0.0001). MOAA/S scores in group D were significantly lower than those in group P during T₁-T₃ (all P<0.0001). However, incidence of respiratory depression in group P was significantly higher than that of group D. Incidence of intraoperative hypotension and sinus bradycardia in group D were significantly higher than that of group P (all P<0.0001). Conclusion: In painless artificial abortions, the sedative effects of dexmedetomidine are significantly better than propofol and there is no respiratory depression. The hemodynamics of propofol are relatively more stable.

Keywords: Dexmedetomidine, propofol, sedation, hemodynamics, respiratory depression

Introduction

Artificial abortion is a common surgery in gynecology. According to statistics, the number of artificial abortions in China is about 10 million per year [1]. As this affects the reproductive health of women, artificial abortions have become a national medical concern [2]. To make patients painless, safe, and comfortable during surgery and to reduce intraoperative and post-operative complications, seeking more ideal methods of anesthesia is essential for clinical anesthesia.

Artificial abortions are generally completed within 10 to 20 minutes. There is little trauma

to the patient but they require high quality anesthesia for pain, safety, comfort, and efficiency. Although the operation time is short, hypotension, decrease of heart rate or even cardiac arrest, intraoperative awareness caused by ineffective analgesia during the operation, and other abnormal phenomena can be life-threatening for patients. Therefore, seeking an anesthesia method with less pain, better sedation, and less adverse reactions after surgery is critical for clinical anesthesia [3, 4].

Painless artificial abortion is a painless surgery with intravenous general anesthesia and suction abortion. There is no pain during the procedure [5]. The most common intravenous anes-

Sedative effects of dexmedetomidine and propofol in painless abortion

Table 1. Comparison of general conditions between the two groups of patients

	Group D	Group P	X ² /t value	P value
Number of cases	100	100		
ASA I	45	50	0.5010	0.4790
ASA II	55	50		
Age	25.2 ± 5.8	28.6 ± 4.7	1.8750	0.0622
Weight (kg)	55.4 ± 7.8	56.3 ± 4.8	1.1160	0.2656
Duration of Pregnancy (d)	54.0 ± 4.8	55.0 ± 5.2	1.4130	0.1592
Operation time (min)	8.2 ± 2.1	7.9 ± 2.6	0.8976	0.3705

Note: Group D: dexmedetomidine group; group P: propofol group; ASA, American Society of Anesthesiologists.

Table 2. Comparison of MAP, HR, and SpO₂ values at different time points in the two groups of patients

		Group D	Group P	T value	P value
MAP (mmHg)	T ₀	123.8 ± 10.3	122.5 ± 9.8	0.9144	0.3616
	T ₁	85.3 ± 4.5	103.6 ± 5.2	36.7905	<0.0001
	T ₂	92.4 ± 3.5	105.3 ± 2.6	29.5868	<0.0001
	T ₃	100.4 ± 4.5	115.5 ± 2.9	28.2058	<0.0001
HR (cpm)	T ₀	70.4 ± 5.6	71.5 ± 3.2	1.7055	0.0897
	T ₁	56.2 ± 3.2	65.2 ± 2.3	22.8379	<0.0001
	T ₂	61.5 ± 3.2	68.3 ± 2.6	16.4924	<0.0001
	T ₃	60.3 ± 2.7	70.3 ± 2.8	25.7087	<0.0001
SpO ₂ (%)	T ₀	99.3 ± 0.7	99.4 ± 0.6	1.0846	0.2794
	T ₁	99.5 ± 0.5	90.4 ± 0.5	128.6930	<0.0001
	T ₂	99.0 ± 0.2	89.4 ± 0.4	214.6620	<0.0001
	T ₃	98.3 ± 1.2	92.7 ± 0.5	43.0770	<0.0001

Note: Group D: dexmedetomidine group; group P: propofol group; T₀: before anesthesia; T₁: 2 min after medication; T₂: uterus dilatation; T₃: 5 min after surgery; HR: heart rate; MAP: mean arterial pressure; SpO₂: oxygen saturation.

thetic used in painless abortion is propofol. Although it has the advantages of rapid onset, short duration of action, and quick recovery, it has weak analgesic effects that could cause body movement during surgery [6]. Dexmedetomidine is an α₂ adrenoceptor-agonist that blocks sympathetic nerves and has strong effects of sedation and hypnosis. It has been widely used in the field of anesthesia [7]. Ishibashi et al. found that during continuous epidural anesthesia of percutaneous nephrolithotomy lithotripsy, dexmedetomidine had obvious characteristics of conscious sedation with less irritability and body movement. Maintenance of intraoperative hemodynamic stability makes the surgery more successful [8].

Dexmedetomidine has rarely been reported in the sedation application of painless artificial

abortion. Therefore, this study compared sedation and hemodynamic effects of propofol and dexmedetomidine in human abortions to provide a basis for clinical anesthesia.

Materials and methods

General information

Using the random number table method, this study selected 200 patients with artificial abortions, from January 2015 to December 2017, in the Department of Anesthesiology of Jinhua Municipal Central Hospital, ASA I-II grade. All patients signed informed consent and the study was approved by the Ethics Committee of Jinhua Municipal of Central Hospital.

Inclusion criteria

Preoperative diagnosis was intrauterine pregnancy. Pregnancy time 30-60 days (gestational sac ≤20 mm). Age was 18 to 40 years old, with no vaginal birth history, and it was the first painless artificial abortion.

Exclusion criteria

Abnormal cardiopulmonary function, abnormal liver and kidney function, mental disorders, and operation times were significantly longer than 20 minutes or significant bleeding during surgery.

Grouping method

Dexmedetomidine group (group D): 100 cases of dexmedetomidine injections for induction and maintenance of anesthesia. Dexmedetomidine hydrochloride injections purchased from Jiangsu Hengrui Medicine Co, Ltd.

Propofol group (P group): 100 cases of propofol injections were used for induction and maintenance of anesthesia. Propofol medium/long-chain fat emulsion injections purchased from Beijing Fresenius Kabi Pharmaceutical Co, Ltd.

Sedative effects of dexmedetomidine and propofol in painless abortion




Figure 1. Comparison of mean arterial pressure, heart rates, and oxygen saturation at four time points in two groups of patients. Group D: dexmedetomidine group; group P: propofol group; T₀: before anesthesia; T₁: 2 min after medication; T₂: uterus dilatation; T₃: 5 min after surgery; MOAA/S: the modified observer's assessment of alertness/sedation scale; HR: heart rate; MAP: mean arterial pressure; SpO₂: oxygen saturation; *P<0.05.

Anesthesia method

Abrasia for 8 hours and water-deprivation for 4 hours before anesthesia were routine for all patients. After admission to the operation room, the venous access of patients was opened. Lactated Ringer's solution was maintained intravenously and titrated slowly. They connected the right upper cuff to monitor blood pressure, monitored heart rate by electrocardiogram, monitored oxygen saturation by pulse oximetry, and maintained airway patency and oxygen with a mask by oxygen flow rate 3.0 L/min. After lithotomy positioning, skin preparation, and draping, patients in group D were given a slow injection of dexmedetomidine by 1.0 µg/kg, gradually entering a sleep state. When the eyelash reflex and response of call disappeared, the abortion was performed. They transfused continuous intravenous infusion to patients at a speed of 0.7 µg/kg/h until the end of surgery. In group P, propofol was induced intravenously at a dose of 0.6 mg/kg, followed by an intravenous infusion to patients at a speed of 3.0 µg/kg/h until the end of the surgery. If patients had physical movement or moaning during surgery, propofol or dexme-

detomidine was added until patients became calm.

Observation indicators

General condition of patients: Patient preoperative age, weight, gestational age, and operation times were observed and recorded.

Observation time points: Time points included T₀ (before anesthesia), T₁ (2 minutes after medication), T₂ (uterus dilatation), and T₃ (5 minutes after surgery).

Main indicators: Hemodynamics: Recorded heart rate, mean arterial pressure, and oxygen saturation at all four time points.

MOAA/S scores: 5 points, completely awake, normal response to normal call; 4 points, responded slowly to normal call; 3 points, no response to normal call, responded to repeated loud calls; 2 points, no response to repeated loud calls, response to a pat on the body; 1 point, no response to a pat on the body, but responded to noxious stimulation; 0 points, no response to noxious stimulation.

Sedative effects of dexmedetomidine and propofol in painless abortion

Table 3. Comparison of improved MOAA/S scores at different time points in the two groups of patients

	Group D	Group P	T value	P value
T ₀	5.00	5.00		
T ₁	0.50 ± 0.04	2.30 ± 0.05	281.1000	<0.0001
T ₂	0.10 ± 0.03	1.20 ± 0.04	220.0000	<0.0001
T ₃	3.50 ± 0.08	4.70 ± 0.10	93.7000	<0.0001

Note: Group D: dexmedetomidine group; group P: propofol group; T₀: before anesthesia; T₁: 2 min after medication; T₂: uterus dilatation; T₃: 5 min after surgery; MOAA/S: the modified observer's assessment of alertness/sedation scale.




Figure 2. Comparison of MOAA/S sedation scores in the two groups of patients. Group D: dexmedetomidine group; group P: propofol group; T₀: before anesthesia; T₁: 2 min after medication; T₂: uterus dilatation; T₃: 5 min after surgery; MOAA/S: the modified observer's assessment of alertness/sedation scale; *P<0.05.

Secondary observation indicators: Adverse reactions: Hypotension, sinus bradycardia, and respiratory depression during the operation were observed and recorded.

Hypotension: systolic blood pressure <90 mmHg or less than 30% of basal blood pressure.

Sinus bradycardia: heart rate <60 beats/min, less than 30% of the basal heart rate.

Respiratory inhibition: Respiratory frequency <8 times/min, oxygen saturation <90%. If respiratory inhibition occurred during the operation, the mask was immediately applied with a pressurized oxygen supply.

Statistical analysis

All data were statistically analyzed using SPSS17.0 software. Graph Pad Prism 5 was used for picture drawing. Measured data are expressed as mean ± sd. Differences between the 2 groups in baseline measurements were

compared using a two-sample independent t-test. Comparison of count data used a double-side Chi-square test (X²) or Fisher's Exact Test. Significance level is defined as α=0.05 and there are statistically significant differences when P<0.05.

Results

Analysis of general information

There were no significant differences between group D and group P in age, ASA grade, and operation time of patients (P>0.05) as shown in **Table 1**.

Hemodynamic comparison between the two groups at four time points

At T₀ (before anesthesia), there were no significant differences in mean arterial pressure, heart rate, and oxygen saturation between the two groups (P>0.05) as shown in **Table 2** and **Figure 1**.

At T₁ (2 minutes after medication), T₂ (uterus dilatation), and T₃ (5 minutes after surgery), mean arterial pressure and heart rates in group D were significantly lower than those in group P. Oxygen saturation was significantly decreased in group P, while there was no significant change in group D. Differences between the two groups at T₁, T₂ and T₃ were statistically significant (all P<0.0001) as shown in **Table 2** and **Figure 1**.

Comparison of MOAA/S scores in the two groups of patients

MOAA/S scores in group D at T₁-T₃ were lower than those in group P and differences were statistically significant (P<0.0001) as shown in **Table 3** and **Figure 2**.

Comparison of intraoperative complications between the two groups

The number of patients with respiratory inhibition in group P was significantly higher than in group D (P<0.0001). Patients with intraoperative hypotension and sinus bradycardia in group D were significantly more than those in group P (both P<0.0001) as shown in **Table 4**.

Discussion

Painless artificial abortion is a new, safe, and effective intravenous general anesthetic. Pre-

Sedative effects of dexmedetomidine and propofol in painless abortion

Table 4. Comparison of intraoperative complications between the two groups

	Group D	Group P	X ² value	P value
Hypotension	36	5	29.4830	<0.0001
Sinus bradycardia	48	10	35.0660	<0.0001
Respiratory inhibition	8	38	25.4090	<0.0001

Note: Group D: dexmedetomidine group; group P: propofol group.

gnant women can enter a sleep state within 30 seconds and surgeons can complete the operation in a short time. Although the operation is short and quick, uterus dilatation operations could bring great physical and mental harm to patients [9]. Intravenous sedative drug infusions during the operation can eliminate patient tension and anxiety, enable hemodynamic stability, and reduce the effects of postoperative complications. Many studies have shown that the use of dexmedetomidine and propofol can maintain a safe and effective conduct of artificial abortions [10, 11].

Some studies have found that dexmedetomidine has significant effects on the cardiovascular system. In the early stage of injection of dexmedetomidine, a transient two-phase cardiovascular reaction appears on the body, a dose-dependent reaction. It has been reported in the literature that a dose of 1 µg/kg dexmedetomidine could cause transient elevation of blood pressure and reflex heart rate reduction. These were more common in young patients or healthy volunteers [12]. This may be due to a direct reduction of peripheral resistance, inhibition of the extraction of calcium ions by the endoplasmic reticulum, inhibition of myocardial contractility, inhibition of the reaction of cyclic baroreceptors to hypotension, inhibition of vasomotor centers, and release of norepinephrine from sympathetic nerve endings. Through the mechanisms above, blood pressure drops and heart rates slow down during intravenous injections of dexmedetomidine [13, 14]. This present study found that intraoperative and postoperative hemodynamics in the propofol group were more stable than the dexmedetomidine group, consistent with previous results.

The present study found that the sedative effects of dexmedetomidine were superior to propofol during induction and maintenance of anesthesia and the postoperative period. Dexmedetomidine is a highly effective α₂ ad-

renergic receptor agonist. Its main mechanism is to regulate arousal and sleep by acting on the A₂AR in the pons and the locus coeruleus of medulla. Locus coeruleus (LC) located in brainstem is a brain nucleus rich in adrenergic receptors that plays a major role in regulating arousal [15, 16]. Dexmedetomidine binds to the α₂-adreno-

receptor on the cell membrane of norepinephrine-producing neurons on LC, inhibits the activity of AC, decreases the content of cAMP in cells, accelerates intracellular anabolic process, and produces sedative and hypnosis effects [17]. Animal experiments have found that sedating doses of dexmedetomidine can inhibit the release of NE from LC [11]. The ventrolateral preoptic nucleus (VLPO) loses its control of NE and releases γ-aminobutyric acid (GABA) and galanin. These two neurotransmitters also inhibit LC and the tuberomammillary nucleus (TMN) in medial thalamus, causing a decrease in TMN histamine release and producing a hypnotic effect [18]. The acting site of dexmedetomidine analgesia is also regarded to be in the spinal dorsal horn [19]. Propofol acts on the central GABA to produce a certain sedative effect, with rapid onset, short duration of action, no accumulation, and rapid recovery [20, 21]. However, propofol has poor analgesia, large intravenous stimulation, and a high dose requirement. Rapid injections or large doses may cause adverse reactions such as respiratory and blood circulation inhibition. Moreover, changes in mean arterial pressure and heart rates during induction, unavoidable intraoperative limb movement, and low levels of sedation are notable defects of propofol.

Some studies have found that dexmedetomidine can produce deeper sedation with less effects on respiration, even at 15 times the therapeutic dose of plasma concentrations [22]. The possible mechanism is that dexmedetomidine retains the body's awakening to hypercapnia, but the threshold of suffocation is reduced [23, 24]. Therefore, compared with an injection of propofol, dexmedetomidine can be safely used for extraction of endotracheal tubes. Although there is no respiratory inhibition, dexmedetomidine was initially approved by the FDA only for patients with initial intubation mechanical ventilation. By October 2008, the FDA approved sedation for non-intubated

Sedative effects of dexmedetomidine and propofol in painless abortion

patients [25]. This study found that incidence of respiratory inhibition in the propofol group was significantly higher than that of the dexmedetomidine group.

There were some limitations to the present study. Sedative effects and hemodynamics of dexmedetomidine and propofol on painless artificial abortion were compared, but no further study was made regarding underlying mechanisms. The next study will examine, in depth, how dexmedetomidine and propofol can improve the mechanism of intraoperative and postoperative sedation and circulation by regulating the nervous system and circulatory system.

In conclusion, in gynecological painless artificial abortion surgeries, the sedative effects of dexmedetomidine are significantly better than propofol. There is no respiratory inhibition with dexmedetomidine, whereas the hemodynamics of propofol are relatively more stable.

Acknowledgements

This work was supported by the Clinical Research Fund of Zhejiang Province Medical Association (2012ZYC-A82).

Disclosure of conflict of interest

None.

Address correspondence to: Zhijian Lan, Department of Anesthesiology, Jinhua Municipal Central Hospital, Jinhua Hospital, Zhejiang University School of Medicine, No.365 Renmin East Road, Jinhua 321000, Zhejiang Province, China. Tel: +86-13586999948; E-mail: lanzhijian12sd@163.com

References

- [1] Li XA, Tang WC, Zeng XR, Zhang JQ and Zou SJ. Application effect of propofol combined with fentanyl in painless artificial abortion. *China Modern Medicine* 2016; 23: 135-137.
- [2] Wang F, XU Rui and Geng ZL. Comparison of anesthetic effects of propofol combined with fentanyl, lornoxicam or nefopam in painless artificial abortion. *Journal of The Fourth Military Medical University* 2007; 28: 1875-1877.
- [3] Wang S. Clinical observation of indolent artificial abortion operation with the application of misoprostol. *Chinese Journal of Ethnomedicine Ethnopharmacy* 2011; 155: 191-204.
- [4] Wu B. Analysis of the effect of painless artificial abortion with misoprostol for termination

- of early pregnancy. *Chinese Community Doctors* 2014; 30: 8-10.
- [5] Fan W, Xue H and Sun Y. Efficacy of dexmedetomidine in attenuating stress response to artificial abortion. *Jiangsu Medical Journal* 2016; 42: 183-185.
- [6] Zhou XM, Yu NH and Li YF. Dezocine and fentanyl combined with propofol for clinical comparative analysis of painless artificial abortion anesthesia. *Journal of Hunan Normal University* 2017; 14: 37-40.
- [7] Jakob SM, Ruokonen E, Grounds RM, Sarapohja T, Garratt C, Pocock SJ, Bratty JR, Takala J; Dexmedetomidine for Long-Term Sedation Investigators. Dexmedetomidine vs midazolam or propofol for sedation during prolonged mechanical ventilation: two randomized controlled trials. *JAMA* 2012; 307: 1151-1160.
- [8] Ishibashi C, Hayashida M, Sugasawa Y, Yamaguchi K, Tomita N, Kajiyama Y and Inada E. Effects of dexmedetomidine on hemodynamics and respiration in intubated, spontaneously breathing patients after endoscopic submucosal dissection for cervical esophageal or pharyngeal cancer. *J Anesth* 2016; 30: 628-636.
- [9] Zhang XH and Er LM. Clinical application of propofol combined with fentanyl intravenous anesthesia in painless artificial abortion. *Chinese Journal of Medicinal Guide* 2015; 17: 288-289.
- [10] Li Y, Gu PF, Shang Y and Li YC. Analgesic effect of dexmedetomidine combined with propofol in artificial abortion surgery. *Chinese Journal of New Drugs* 2011; 20: 1885-1891.
- [11] Zhang ZW and Wang YH. Comparative study on dexmedetomidine and fentanyl combined with propofol for anaesthesia in painless abortion. *Drugs Clinic* 2017; 32: 1085-1089.
- [12] Shank ES, Sheridan RL, Ryan CM, Keaney TJ and Martyn JA. Hemodynamic responses to dexmedetomidine in critically injured intubated pediatric burned patients: a preliminary study. *J Burn Care Res* 2013; 34: 311-317.
- [13] Wit F, Vliet AL, Wilde RB, Jansen JR, Vuyk J, Aarts LP, Jonge E, Veelo DP and Geerts BF. The effect of propofol on haemodynamics: cardiac output, venous return, mean systemic filling pressure, and vascular resistances. *Br J Anaesth* 2016; 116: 784-789.
- [14] Łasińska-Kowara M, Kardel-Reszkiewicz E, Owczuk R. Effects of sevoflurane versus target-controlled infusion of propofol on haemodynamics during elective breast surgery in healthy women. *Anestezjol Intens Ter* 2009; 41: 135-139.
- [15] Djaiani G, Silverton N, Fedorko L, Carroll J, Styra R, Rao V and Katznelson R. Dexmedetomidine versus propofol sedation reduces delirium after cardiac surgery: a randomized con-

Sedative effects of dexmedetomidine and propofol in painless abortion

- trolled trial. *Anesthesiology* 2015; 124: 362-368.
- [16] Li BL, Ni J, Huang JX, Zhang N, Song XR and Yuen VM. Intranasal dexmedetomidine for sedation in children undergoing transthoracic echocardiography study—a prospective observational study. *Paediatric Anaesthesia* 2015; 25: 891-896.
- [17] Wang D, Cao Y, Yi L, Li Y and Tao H. Corrective effect of norepinephrine on hypotension induced by dexmedetomidine in critically ill patients. *Int J Clin Pharmacol Ther* 2016; 54: 987-991.
- [18] Wang XF, Luo XL, Liu WC, Hou BC, Huang J, Zhan YP and Chen SB. Effect of dexmedetomidine priming on convulsion reaction induced by lidocaine. *Medicine* 2016; 95: e4781.
- [19] Scibelli G, Maio L, Sasso M, Lanza A and Savoia G. Dexmedetomidine: current role in burn ICU. *Translational Medicine* 2017; 16: 1-10.
- [20] Germann AL, Shin DJ, Manion BD, Edge CJ, Smith EH, Franks NP, Evers AS and Akk G. Activation and modulation of recombinant glycine and GABA receptors by 4-halogenated analogues of propofol: actions of halogenated propofol analogues. *Br J Pharmacol* 2016; 173: 3110-3120.
- [21] Ziemba A M and Forman SA. Correction for inhibition leads to an allosteric co-agonist model for pentobarbital modulation and activation of $\alpha 1\beta 3\gamma 2\delta$ GABA receptors. *PLoS One* 2016; 11: e0154031.
- [22] Skrupky L P, Drewry AM, Wessman B, Field RR, Fagley RE, Varghese L, Lieu A, Olatunde J, Micek ST, Kollef MH and Boyle WA. Clinical effectiveness of a sedation protocol minimizing benzodiazepine infusions and favoring early dexmedetomidine: a before-after study. *Critical Care* 2015; 19: 1-12.
- [23] Guo TZ, Jiang JY, Buttermann AE and Maze M. Dexmedetomidine injection into the locus ceruleus produces antinociception. *Anesthesiology* 1996; 84: 873-881.
- [24] Zhan-Ying G, Chang-Ming W, Shuai T, Lin-Lin T, Yu-Feng H. Comparison of effects of different doses dexmedetomidine on inhibiting tracheal intubation-evoked haemodynamic response in the elderly patients. *Journal of Clinical Diagnostic Research Jcdr* 2015; 9: UC10-103.
- [25] Zhou Q. Effect analysis of doxapram and dexmedetomidine in the prevention of postoperation chill of esophageal cancer treated with radical surgery under general anesthesia. *China Modern Medicine* 2015; 22: 123-125.