

Original Article

Liushen Pills inhibit inflammation and growth of esophageal cancer by promoting the inflammatory cell infiltration

Hui Zhang¹, Jingwen Jiang¹, Xuewu Chen¹, Fengting Zhu¹, Lei Fu²

Departments of ¹Oncology, ²Dermatology, Hainan Hospital of Traditional Chinese Medicine, Haikou 570203, Hainan, China

Received October 11, 2019; Accepted November 26, 2019; Epub February 15, 2020; Published February 28, 2020

Abstract: Background: Liushen pills (LSP) is a Chinese traditional medicine recorded by the Chinese medical dictionary. This study is aimed to investigate the effect of LSP on TNF- α /IL-1 β and iNOS signaling pathways and their roles in the migration and invasion of esophageal cancer. Methods: By establishing the nude mice xenograft model of esophageal cancer, we evaluated the effect of LSP on esophageal cancer. The levels of chemokines were detected by Western blot and ELISA. Then, the role of LSP in cell proliferation, apoptosis, migration and invasion were validated by preparing the drug-containing serum. Results: LSP inhibited the tumor growth of esophageal cancer. LSP promoted the inflammatory cells infiltration around the tumor and induced the cell necrosis. LSP treatment significantly inhibited the levels of TLR4, Myd88, TNF- α , IKK- α , p-IKK α , IKK β , p-IKK β , iNOS, COX-2, IL-1 β and IL-6 in esophageal cancer tissues. LSP serum induced cell apoptosis, and suppressed cell migration and invasion. Conclusion: LSP improves the microenvironment by inhibiting the inflammation and its contribution to metastasis via inhibition of esophageal cancer.

Keywords: Chinese medicine, Liushen pills, inflammation, metastasis, esophageal cancer

Introduction

Esophageal cancer is a malignant tumor that occurs in the epithelium of esophageal mucosa, which is one of the most common malignant tumors in the world [1]. The incidence of esophageal cancer in China ranks fifth among all the malignant tumors and the mortality rate ranks fourth [2]. Thus, it is one of the diseases that seriously threaten human life and health.

In 1889, Stephen Paget proposed the hypothesis of the metastatic “seed and soil”. He believed that the growth of cancer cells must rely on the surrounding tissues to provide a specific environment [3]. In the process of tumorigenesis and development, the dynamic balance between the tumor cells and the surrounding tissue environment is constantly destroyed, causing the cell proliferation, differentiation, apoptosis, and expression and secretion of cytokines in cell surface are discarded, while the external tumor environment conducive to their own growth is constantly estab-

lished, which leads to malignant transformation in the whole process of tumorigenesis and progression [4]. The tumor environment has the characteristics of hypoxia, low mesenchymal pressure, proteolytic fermentation and a large number of growth factors [3, 4]. It is the survival environment in which tumor cells grow and proliferate, and induces neovascularization, inhibits immune inflammatory reactions and inoculates cancer stem cells to facilitate the migration, invasion and metastasis of tumor cells [5-7]. In the tumor environment, a large number of cellular molecules (i.e. iNOS), growth factors, chemokines (i.e. TNF- α , IL-1 β , and IL-6) and various proteolytic enzymes produced by the immune inflammatory response constituted the inflammatory microenvironment of tumors are particularly important [8, 9]. In recent years, more and more attention has been paid to the role of inflammatory microenvironment in the process of carcinogenesis [10, 11]. Antineoplastic drugs targeting tumor microenvironment such as non-steroidal anti-inflammatory drug COX-2 inhibitor celecoxib

have been emerging [12]. Therefore, exploring the mechanism of inflammatory microenvironment of tumors can provide a novel strategy for the treatment of tumors.

Liushen Pill (LSP) is recorded by the Chinese medical dictionary. It is composed of six medicines including bezoar, fragrance, pearl, borneol, toad venom and realgar, which is derived from Leiyun Shangyufeng Tang Fang [13]. LSP has shown good outcome in the treatment of liver cancer, gastric cancer, colon cancer, cervical cancer, and esophageal cancer [14]. Previous studies have shown that LSP inhibits the growth of esophageal cancer by inhibiting angiogenesis [13]. In this study, we aimed to investigate the effect of LSP on TNF- α /IL-1 β and iNOS signaling pathways and their roles in the migration and invasion of esophageal cancer. The findings indicate that TNF- α pathway in the inflammatory microenvironment was involved in the metastasis of esophageal cancer.

Materials and methods

Cell culture

Human esophageal cancer cell line Eca-109 (ATCC, USA) was cultured in 50 mL flask. The cells were cultured in high glucose DMEM medium (Hyclone, USA) containing 10% newborn calf serum (Hyclone, USA) in a constant temperature incubator with 37°C, 5% CO₂ and saturated humidity.

Preparation of LPS solution

Liushen pills were thoroughly ground in a mortar and adjusted to 0.48 mg/mL with three distilled water for gastric lavage. The solutions were separately packed and stored at 4°C. Before use, the solution was warmed to room temperature and well mixed.

Animals and modeling

A total of 10 healthy, male, BALB/c nude mice, aged 6-8 weeks and weighted 18-22 g were purchased from Carvins Laboratory Animal Co., Ltd. (Changzhou, China). For drug-serum preparation, a total of 10 SD, male, rats weighted 200 g were randomly divided into normal serum group and LSP serum group. The mice and rats were fed with food and water freely, ventilated for 24 h in SPF animal room with 24-28°C, humidity 40% -50%. After 1 week of adaptive

feeding, the xenograft mice model was performed under anesthetized by intraperitoneal injection of chloral hydrate (400 mg/kg), and the rat drug-containing serum was prepared under anesthetized with sodium pentobarbital (50 mg/kg). At the end of experiments, the mice and rats were euthanized by increasing CO₂ levels to 33% over 1 min. Death was verified by decapitation. This study was approved by the Ethics Committee of Hainan Hospital of Traditional Chinese Medicine (no.2018.1.13, Haikou, China).

Xenograft model of esophageal cancer in nude mice

Healthy male BALB/c mice aged 6-8 weeks old and weighted 18-22 g were kept with isolator. The mice were anesthetized by intraperitoneal injection of chloral hydrate. 0.2 mL of human esophageal cancer cell suspension (1×10^7 /mL) was extracted and inserted into the axillary subcutaneously. LSP treatment group was intragastrically administered with 4 mL LSP once a day for 20 days (n=5). The model group was intragastrically administered with normal saline (n=5). The tumor size was measured in vitro with Vernier caliper every 5 days, and the volume of tumor was calculated and the growth curve was drawn. After 20 days, the mice were sacrificed by spinal dislocation and the tumor were removed. After fixed with polyformaldehyde, the tumor tissue were dehydrated by gradient alcohol, embedded in paraffin, sectioned for routinely hematoxylin-eosin (HE) staining. The levels of TLR4, Myd88, TNF- α , IKK- α , p-IKK α , IKK β , p-IKK β , iNOS and COX-2 were detected by Western blot. The levels of IL-1 β and IL-6 were detected by ELISA.

Preparation of drug-containing serum

The SD male rats weighted 200 g were randomly divided into normal serum group and LSP serum group (n=5). The rats in LSP serum group were given 4 mL of LSP solution intragastrically twice a day for 3 days, while the normal serum group was given the same amount of saline daily. The rats were fasted for 12 h before the last intragastric administration and anesthetized with sodium pentobarbital after the last gavage for 1 h. Blood was collected from the posterior abdominal aorta and ventricle. Serum was aseptically separated, in activated and sterilized, and then placed in a vial and stored

at -20°C. The esophageal cancer cell was cultured using the normal serum and LSP serum for 48 h, respectively.

MTT assay

The cell proliferation was determined by MTT assay. The cells were inoculated in 96-well flat-bottom culture plate with 200 µL/well (2×10^4 cells per well) and cultured for 24 h. Then, the culture medium was removed and replaced with the normal serum and LSP serum, respectively for 24, 48, and 72 h. Four hours before the end of treatment, 20 µL MTT (5 mg/mL) was added into each well. At the end of treatment, 200 µL dimethyl sulfoxide was added to each well for 5 min. The absorbance was measured at 490 nm. Finally, the growth inhibition rate was calculated.

Detection of apoptosis by flow cytometry

The cells treated with normal serum and LSP serum were collected and suspended in binding buffer to adjust to 1×10^7 /ml. 100 µL cells were taken and mixed with 5 µL annexin V-FITC and 5 µL PI for 15 min at room temperature in the dark. After adding 400 µL of binding buffer, the apoptosis was detected with a flow cytometer. The flow cytometry data was obtained by FACS Calibur flow cytometry and CELLQuest software, and the experimental data were analyzed using Winmidi 218 software. The normal cells are located at the lower left, Annexin V (-) and PI (-). The early apoptotic cells are located at the lower right, Annexin V (+) and PI (-). The late apoptotic cells are located at the upper right, Annexin V (+) and PI (+).

ELISA

Levels of IL-1β and IL-6 were measured using the commercially available enzyme-linked immunosorbent assay (ELISA) kits according to the manufacturer's instructions (Raybiotech Inc., GA, USA). The intra-assay and inter-assay coefficients of variation were less than 10% in enzyme immunoassays.

Transwell assay for migration and invasion

The cell migration and invasion were performed with QCM-96-well cell migration and invasion assay kits (Chemicon International, Temecula, CA, USA). The tumor cells were stained by

Crystal violet and photographed under $\times 100$ microscopy (Olympus, Japan).

Western blot

Rabbit anti-TLR4, Myd88, TNF-α, IKK-α, p-IKKα, IKKβ, p-IKKβ, iNOS, COX-2 (Abcam, 1:1000) and 1:3000 dilution of anti-rabbit IgG secondary antibody (ZSGB-BIO, China, 1:1000) were used to determine the protein expression. Rabbit anti-GAPDH (ZSGB-BIO, China, 1:1000) was used as an internal control. Images were acquired with a ChemiDoc Touch imager (Bio-Rad, USA) and quantification was done using Quantity One 4.4.0 software (Bio-Rad, USA).

Statistical analysis

All data are presented as the mean \pm standard deviation of at least 3 independent experiments. Graph construction and statistical analysis were performed using SPSS 17.0 and GraphPad Prism 5.0. Differences between groups were evaluated with Student's t test. *P* values <0.05 were considered to be statistically significant.

Results

Effect of LSP on the growth of esophageal cancer

According to the change in transplanted human esophageal cancer in nude mice (**Figure 1A**), the growth curve was drawn (**Figure 1C**). LSP significantly inhibited the growth of esophageal cancer after 10 days treatment ($P < 0.05$). The subcutaneous tumors were exfoliated (**Figure 1B**). The weight of tumors in model group were significantly higher than LSP treatment group ($P < 0.001$) (**Figure 1D**). The tumors in model group were incomplete in envelope, while the LSP treatment tumors were intact in envelope (**Figure 1B**). These findings suggested that LSP inhibited the progression of esophageal cancer.

Histopathological observation of tumor tissues

In the model group, the cancer cells invaded the deep muscular layer, and the nucleus of the cancer cells was large and hyperchromatic (**Figure 2**). In the LSP group, inflammatory cells infiltrated around the tumor and the cancer cells invaded the muscular layer of the line




Figure 1. Effect of LSP on the growth of esophageal cancer. A. The xenograft model of esophageal cancer after treatment of LSP for 20 days. B. The tumors. C. The growth curve of esophageal cancer. D. The weight of tumors. * $P < 0.05$, *** $P < 0.001$.




Figure 2. The histopathological change in tumor tissues in xenograft model after LSP treatment (100x). The scale bar is 100 μm.

(Figure 2). The abnormality of the tumor cells was obvious, and the necrosis surface was small. It showed hemorrhagic foci in LSP treated tumor tissues (Figure 2).

Effect of LSP on levels of TLR4, Myd88, TNF- α , IKK- α , p-IKK α , IKK β , p-IKK β , iNOS, COX-2, IL-1 β and IL-6

The levels of TLR4, Myd88, TNF- α , IKK- α , p-IKK α , IKK β , p-IKK β , iNOS and COX-2 were detected by Western blot (Figure 3A-C). The levels of IL-1 β and IL-6 were detected by ELISA (Figure 3D). LSP treatment significantly inhibited the levels of TLR4, Myd88, TNF- α , IKK- α , p-IKK α , IKK β , p-IKK β , iNOS, COX-2, IL-1 β and

IL-6 in esophageal cancer tissues.

LSP serum induced cell apoptosis, and suppressed cell migration and invasion of esophageal cancer cells.

To further analyze the role of LSP in esophageal cancer cells, human esophageal cancer cells (line Eca-109) were treated with LSP serum (Figure 4). Compared with normal serum, LSP serum inhibited the proliferation of esophageal cancer cells at 48 h and 72 h ($P < 0.05$). The apoptotic rate of the esophageal cancer cells was significantly increased. The apoptotic rate was mostly increased in the early apoptotic cells (Figure 4B). The migration and invasion of esophageal cells were inhibited by LSP treatment. Thus, the results suggested that LSP induces early apoptosis, and inhibited the migration and invasion of esophageal cancer cells.

Discussion

Metastasis are essential characteristics of malignant tumors and the leading causes of death in tumor patients [15, 16]. The occurrence, development, invasion and metastasis of tumors are complex processes involving cell-cell communication and multi-gene changes in microenvironment [17, 18]. Tumor invasion and metastasis are two different stages in the same process [19]. They are closely linked to each other. Invasion is the basis of metastasis, and metastasis is the consequence of invasion. The movement and migration of cancer cells ultimately achieve the invasion and metastasis of tumors [20]. In this study, we found that LSP promoted inflammatory cell infiltration, but inhibited the esophageal cancer cell migration, invasion and metastasis.

TNF- α , IL-6 and NF- κ B are important cytokines in the inflammatory microenvironment of tu-




Figure 3. Levels of TLR4, Myd88, TNF- α , IKK- α , p-IKK α , IKK β , p-IKK β , iNOS, COX-2, IL-1 β and IL-6 in xenograft model after LSP treatment. TLR4, Myd88, TNF- α , IKK- α , p-IKK α , IKK β , p-IKK β , iNOS and COX-2 levels in tumor tissues in xenograft model after LSP treatment were performed by Western blot. The levels of IL-1 β and IL-6 were detected by ELISA. ***P<0.001.




Figure 4. Effect of drug-containing serum on esophageal cancer cells. Serum was aseptically separated from SD rats after LSP or normal saline administration intragastrically twice a day for 3 days. (A) Cell proliferation was detected by MTT assay. (B) Cell apoptosis was detected by flow cytometry. (C) Migration and (D) invasion were detected by Transwell assays.

mors [21, 22]. IL-6 is an inflammatory factor, which is generally expressed in various tumor microenvironments and inflammatory diseases [23, 24]. It participates in the occurrence of various tumors, such as hepatocellular carcinoma, lung cancer, oral cancer, gall bladder carcinomas, gynecologic cancer and esophageal cancer [23-25]. Its function is to affect cell cycle, regulate angiogenesis and so on by activating multiple signaling pathways such as

noma, lung cancer, oral cancer, gall bladder carcinomas, gynecologic cancer and esophageal cancer [23-25]. Its function is to affect cell cycle, regulate angiogenesis and so on by activating multiple signaling pathways such as

JAKs/STAT3 [22]. TNF- α is an essential factor in the initiation of chronic inflammation and is one of the most influential inflammatory mediators in the early stage of inflammation [26]. TNF- α can activate the NF- κ B pathway and triggers the transcription of many inflammatory cytokine genes, leading to tumorigenesis [27, 28]. It was found that TNF- α was expressed in tumor cells and stromal cells of breast cancer, prostate cancer, bladder cancer, colon cancer, liver cancer, lymphoma and leukemia, and the prognosis of patients with positive TNF- α was poor [26]. TNF- α could induce epithelial-mesenchymal transition through NF- κ B pathway mediated Snail stabilization, which can significantly promote the migration and invasion of tumor cells [28, 29]. TNF- α and IL-1 β are not only the downstream target genes of NF- κ B, but also the promoter of NF- κ B. After activation of NF- κ B pathway by TNF- α and IL-1 β , NF- κ B can be further activated, and the cycle is repeated, so that the inflammatory response is continuously expanded [30]. Most of the inflammatory reactions are self-limiting, but this process of inflammation self-amplification leads to excessive autologous tissue damage, which promotes the occurrence and development of tumors [31, 32]. In this study, we found that LSP treatment significantly inhibited the levels of inflammatory factors including TLR4, Myd88, TNF- α , IKK- α , p-IKK α , IKK β , p-IKK β , iNOS, COX-2, IL-1 β and IL-6 in esophageal cancer tissues.

At present, the main methods of cancer treatment, such as surgery, radiotherapy and chemotherapy, cause damage to tissues and cells and then cause inflammation. This inflammatory environment is beneficial to tumor cells and increases the possibility of metastasis. Therapeutic effect will inevitably be discounted. Therefore, improving the microenvironment for treating inflammation can also improve the therapeutic effect and inhibit the occurrence and metastasis of tumors to a certain extent, which is of great significance for the treatment of tumors. LSP promoted the cell apoptosis but inhibited the inflammation, showing a promising role in improving the microenvironment for inhibition of tumor metastasis. However, due to the complexity of the inflammatory microenvironment, the role and mechanism of the inflammatory microenvironment are not yet fully elucidated and the establishment of tumor microenvironment model is still lack of certain standards. Nevertheless, we could conclude that

LSP improves the microenvironment by inhibiting the inflammation and contribution to metastasis via inhibition of esophageal cancer.

Acknowledgements

The Natural science foundation of hainan province (Project No. 817338); Hainan province in 2019-2021 provincial key subject construction project of traditional Chinese medicine (The Oncology Department) (NO. S100111.401).

Disclosure of conflict of interest

None.

Address correspondence to: Dr. Lei Fu, Department of Dermatology, Hainan Hospital of Traditional Chinese Medicine, No. 48, Heping North Road, Meilan District, Haikou 570203, Hainan Province, China. Tel: +86-0898-66203809; E-mail: layfu52143@163.com

References

- [1] Lin Y, Totsuka Y, Shan B, Wang C, Wei W, Qiao Y, Kikuchi S, Inoue M, Tanaka H and He Y. Esophageal cancer in high-risk areas of China: research progress and challenges. *Ann Epidemiol* 2017; 27: 215-221.
- [2] Chen W, Zheng R, Zeng H, Zhang S and He J. Annual report on status of cancer in China, 2011. *Chin J Cancer Res* 2015; 27: 2-12.
- [3] de Groot AE, Roy S, Brown JS, Pienta KJ and Amend SR. Revisiting seed and soil: examining the primary tumor and cancer cell foraging in metastasis. *Mol Cancer Res* 2017; 15: 361-370.
- [4] Fidler IJ. The pathogenesis of cancer metastasis: the 'seed and soil' hypothesis revisited. *Nat Rev Cancer* 2003; 3: 453-458.
- [5] Vander Linden C and Corbet C. Reconciling environment-mediated metabolic heterogeneity with the oncogene-driven cancer paradigm in precision oncology. *Semin Cell Dev Biol* 2020; 98: 202-210.
- [6] Looi CK, Chung FF, Leong CO, Wong SF, Rosli R and Mai CW. Therapeutic challenges and current immunomodulatory strategies in targeting the immunosuppressive pancreatic tumor microenvironment. *J Exp Clin Cancer Res* 2019; 38: 162.
- [7] Hinshaw DC and Shevde LA. The tumor microenvironment innately modulates cancer progression. *Cancer Res* 2019; 79:4557-4566.
- [8] Mantovani A. Molecular pathways linking inflammation and cancer. *Curr Mol Med* 2010; 10: 369-373.

- [9] Vannini F, Kashfi K and Nath N. The dual role of iNOS in cancer. *Redox Biol* 2015; 6: 334-343.
- [10] Quail DF and Joyce JA. Microenvironmental regulation of tumor progression and metastasis. *Nat Med* 2013; 19: 1423-1437.
- [11] Borsig L, Wolf MJ, Roblek M, Lorentzen A and Heikenwalder M. Inflammatory chemokines and metastasis—tracing the accessory. *Oncogene* 2014; 33: 3217-3224.
- [12] Wang YP, Wang QY, Li CH and Li XW. COX-2 inhibition by celecoxib in epithelial ovarian cancer attenuates E-cadherin suppression through reduced snail nuclear translocation. *Chem Biol Interact* 2018; 292: 24-29.
- [13] Bi YF, Mao JY and Liu CX. Current status of study on impacts of Chinese herbal drugs and their preparations on pharmacokinetics. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 2008; 28: 662-665.
- [14] Yang Q, Zhou X, Zhang M, Bi L, Miao S, Cao W, Xie Y, Sun J, Tang H, Li Y, Miao Q and Wang S. Angel of human health: current research updates in toad medicine. *Am J Transl Res* 2015; 7: 1-14.
- [15] Zeng Y. Cell biology, biophysics, and mechanobiology: from the basics to clinics. *Cell Mol Biol (Noisy-le-grand)* 2017; 63: 1-2.
- [16] Zeng Y, Yao X, Chen L, Yan Z, Liu J, Zhang Y, Feng T, Wu J and Liu X. Sphingosine-1-phosphate induced epithelial-mesenchymal transition of hepatocellular carcinoma via an MMP-7/syndecan-1/TGF-beta autocrine loop. *Oncotarget* 2016; 7: 63324-63337.
- [17] Zeng Y, Yao X, Liu X, He X, Li L, Liu X, Yan Z, Wu J and Fu B. Anti-angiogenesis triggers exosomes release from endothelial cells to promote tumor vasculogenesis. *J Extracell Vesicles* 2019; 8: 1629865.
- [18] Steeg PS. Targeting metastasis. *Nat Rev Cancer* 2016; 16: 201-218.
- [19] Fidler IJ and Kripke ML. The challenge of targeting metastasis. *Cancer Metastasis Rev* 2015; 34: 635-641.
- [20] Lintz M, Munoz A and Reinhart-King CA. The mechanics of single cell and collective migration of tumor cells. *J Biomech Eng* 2017; 139: 021005.
- [21] Huang C, Yang X, Huang J, Liu X, Yang X, Jin H, Huang Q, Li L and Zhou R. Porcine beta-defensin 2 provides protection against bacterial infection by a direct bactericidal activity and alleviates inflammation via interference with the TLR4/NF-kappaB pathway. *Front Immunol* 2019; 10: 1673.
- [22] Lee SB, Lee WS, Shin JS, Jang DS and Lee KT. Xanthotoxin suppresses LPS-induced expression of iNOS, COX-2, TNF-alpha, and IL-6 via AP-1, NF-kappaB, and JAK-STAT inactivation in RAW 264.7 macrophages. *Int Immunopharmacol* 2017; 49: 21-29.
- [23] Yang J, Wang J and Luo J. Decreased IL-6 induces sensitivity of hepatocellular carcinoma cells to sorafenib. *Pathol Res Pract* 2019; 215: 152565.
- [24] Vinocha A, Grover RK and Deepak R. Clinical significance of interleukin-6 in diagnosis of lung, oral, esophageal, and gall bladder carcinomas. *J Cancer Res Ther* 2018; 14: S758-S760.
- [25] Kacel EL, Kirsch JL, Sannes TS, Patidar S, Postupack R, Jensen S, Wong S, Garey S, Dodd S, Ulfing CM, McCrae CS, Robinson ME, Castagno J, Schultz GS and Pereira DB. Interleukin-6 and body mass index, tobacco use, and sleep in gynecologic cancers. *Health Psychol* 2019; 38: 866-877.
- [26] Gupta M, Babic A, Beck AH and Terry K. TNF-alpha expression, risk factors, and inflammatory exposures in ovarian cancer: evidence for an inflammatory pathway of ovarian carcinogenesis? *Hum Pathol* 2016; 54: 82-91.
- [27] Nicolini A, Ferrari P, Diodati L and Carpi A. Alterations of signaling pathways related to the immune system in breast cancer: new perspectives in patient management. *Int J Mol Sci* 2018; 19: 2733.
- [28] Annibaldi A and Meier P. Checkpoints in TNF-induced cell death: implications in inflammation and cancer. *Trends Mol Med* 2018; 24: 49-65.
- [29] Shen J, Xiao Z, Zhao Q, Li M, Wu X, Zhang L, Hu W and Cho CH. Anti-cancer therapy with TNFalpha and IFNgamma: a comprehensive review. *Cell Prolif* 2018; 51: e12441.
- [30] Vendramini-Costa DB and Carvalho JE. Molecular link mechanisms between inflammation and cancer. *Curr Pharm Des* 2012; 18: 3831-3852.
- [31] Lu N and Malesmud CJ. Extracellular signal-regulated kinase: a regulator of cell growth, inflammation, chondrocyte and bone cell receptor-mediated gene expression. *Int J Mol Sci* 2019; 20.
- [32] Kay J, Thadhani E, Samson L and Engelward B. Inflammation-induced DNA damage, mutations and cancer. *DNA Repair (Amst)* 2019; 83: 102673.