

Erratum

Polydatin regulates Wnt/ β -catenin signaling to inhibit proliferation and migration of hepatocellular carcinoma cells: Int J Clin Exp Med. 2019; 12(11): 12670-12678

Jian Xu^{1,2}, Wei Zhang^{1,2}, Changkang Wu^{1,2}, Jingdong Li^{1,2}

¹Department I of Hepatobiliary Surgery, Affiliated Hospital of North Sichuan Medical College, Nanchong, Sichuan, China; ²Hepatobiliary, Pancreatic and Intestinal Diseases Research Institute of North Sichuan Medical College, Nanchong, Sichuan, China

Received July 7, 2022; Accepted August 30, 2023; Epub November 15, 2023; Published November 30, 2023

In this article, we found some problems in **Figure 9**. Therefore, we would like to publish this Erratum to reflect this change. We apologize for these problems.

The corrected **Figure 9** is as follows.

Address correspondence to: Jingdong Li, Department I of Hepatobiliary Surgery, Affiliated Hospital

of North Sichuan Medical College, Nanchong, Sichuan, China; Hepatobiliary, Pancreatic and Intestinal Diseases Research Institute of North Sichuan Medical College, No.63 Wenhua Road, Nanchong 637000, Sichuan, China. Tel: +86-158-81750153; E-mail: jingdonglii@163.com

Polydatin targets Wnt/ β -catenin to inhibit cancer growth




Figure 9. (A) Effect of Polydatin β -catenin protein and downstream target genes in SMMC-7721 cells. (B) Effect of Polydatin on protein expressions of β -catenin and downstream target genes in HepG2 cells. A. control; B. 25 μ g/ml; C. 50 μ g/ml; D. 100 μ g/ml. (C) Effect of Resveratrol+SB-216763I on protein expressions of β -catenin and downstream target genes in SMMC-7721 cells. (D) Effect of Resveratrol+SB-216763I on protein expressions of β -catenin and downstream target genes in HepG2 cells.