

Original Article

Reduced beta 2 glycoprotein I improve diabetic nephropathy via inhibiting TGF- β 1-p38 MAPK pathway

Tong Wang, Si-Si Chen, Rui Chen, De-Min Yu, Pei Yu

2011 Collaborative Innovation Center of Tianjin for Medical Epigenetics, The Key Laboratory of Hormones and Development (Ministry of Health), Metabolic Diseases Hospital & Tianjin Institute of Endocrinology, Tianjin Medical University, Tianjin 300070, China

Received February 27, 2015; Accepted May 2, 2015; Epub May 15, 2015; Published May 30, 2015

Abstract: Purpose: Beta 2 glycoprotein I (β 2GPI) has been shown the positive effect on diabetic atherosclerosis and retinal neovascularization. β 2GPI can be reduced by thioredoxin-1, resulting in the reduced state of β 2GPI. The possible protective effects of β 2GPI and reduced β 2GPI on diabetic nephropathy (DN) are not fully elucidated. The purpose of this study was to test a hypothesis that β 2GPI and reduced β 2GPI would improve DN in streptozotocin (STZ) induced diabetic mice and high-glucose (HG) exposed rat mesangial cell (RMC). Methods: The STZ-induced Balb/c mice and HG exposed RMCs were administrated with β 2-GPI and reduced β 2-GPI at different time and concentrations gradient respectively. The changes of glomerular structure and expression of collagen IV, TGF- β 1, p38 MAPK and phospho-p38 MAPK in renal cortical and mesangial cells were observed by immunohistochemical techniques, quantitative real-time PCR and western blot with or without the treatment of β 2-GPI and reduced β 2-GPI. Results: β 2GPI and reduced β 2GPI improved early clinical and pathological changes of DN in STZ-diabetic mice. Treatment with β 2GPI and reduced β 2GPI in the STZ-diabetic mice and HG exposed RMCs resulted in decrease expression levels of TGF- β 1 and collagen IV, with concomitant decrease in phospho-p38 MAPK expression. Conclusions: β 2GPI and reduced β 2GPI improved renal structural damage and kidney function. The renoprotective and antifibrosis effects of β 2GPI and reduced β 2GPI on DN were closely associated with suppressing the activation of the TGF- β 1-p38 MAPK pathway.

Keywords: Beta 2 glycoprotein I, diabetic nephropathy, TGF- β 1

Introduction

Diabetic nephropathy (DN) is the major microvascular complication of diabetes mellitus (DM) and is the most common cause of end-stage renal disease worldwide [1]. Clinically, DN is characterized by a reduction in the glomerular filtration rate and increasing proteinuria that eventually leads to progressive renal failure. Pathologically, DN is characterized by the glomerular mesangium hypertrophy caused by a proliferation of mesangial cells and excessive accumulation of extracellular matrix (ECM), which ultimately progresses to kidney fibrosis and glomerulosclerosis [2].

Transforming growth factor- β 1 (TGF- β 1) is a fibrogenic and inflammatory cytokine that plays a vital role in glomerulosclerosis and interstitial fibrosis in various renal diseases, including DN. The abnormal production of TGF- β 1 induced by

hyperglycemia causes an excessive accumulation of ECM proteins, such as collagen and fibronectin, which ultimately leads to mesangial expansion and glomerular basement membrane thickening. The mitogen-activated protein kinases (MAPK), a family of serine/threonine kinases, regulate intracellular signal transduction in response to various extracellular stimuli. MAPK subfamilies contain extracellular signal-regulated kinase (ERK) 1/2, c-Jun NH2-terminal kinase (JNK), and p38 mitogen-activated protein kinase (p38 MAPK) [3]. Recently, some evidence suggests that the activation of the p38 MAPK pathway, but not the canonical Smad signaling pathway, mediates TGF- β 1-induced collagen and fibronectin expression during the development of DN [4].

Beta 2 glycoprotein I (β 2GPI) is a 50-kDa protein that was first described in 1961[5], and since 1990 β 2GPI has been identified as the

Reduced β 2GPI improve diabetic nephropathy

most prominent antigen in antiphospholipid syndrome [6]. β 2GPI is predominantly synthesized in hepatocytes, and its circulating plasma concentration is variable (50-500 μ g/mL, 1-10 μ M) [7]. β 2GPI consists of five repeating amino acid domains. Domains I-IV include four cysteines each and have the conserved sequences, domain V has an extra 20 amino acid tail with a unique cysteine termination and is aberrant [8]. Recently, it has been shown that the disulfide bond between Cys288 to Cys326 in the domain V can be reduced by thioredoxin-1 (TRX-1) or protein disulfide isomerase (PDI) [9], resulting in the reduced state of β 2GPI, referred to as reduced β 2GPI.

Although β 2GPI has been shown to be a participant in the autoimmune system [10, 11], vascular thrombosis [12, 13], infectious diseases [14] and others system, the function of β 2GPI remains unclear. It has also reported that β 2GPI is associated with accelerated atherosclerosis and enhanced oxidative stress [15, 16]. High levels of β 2GPI have been reported to decrease the risk of myocardial infarction in elderly men [17]. In addition, reduced β 2GPI was found to protect endothelial cells from oxidative stress-induced injury [18]. We have previously found a positive effect of β 2GPI and reduced β 2GPI in diabetic atherosclerosis and retinal neovascularization [19]. Till now, no studies have examined the relationship between β 2GPI and reduced β 2GPI and DN. In this study, we hypothesize that β 2GPI or reduced β 2GPI has beneficial effects in DN. In addition, we investigated the renoprotective effects of β 2GPI or reduced β 2GPI in the TGF- β 1-p38 MAPK pathway by examining the expression of total and phosphorylated p38 MAPK (p-p38 MAPK), TGF- β 1, and collagen IV in diabetic mice and high glucose exposed rat mesangial cells.

Materials and methods

Materials and reagents

The rat mesangial cell (RMC) line, HBZY-1, was obtained from the American Type Culture Collection (ATCC number: CRL-2573). Fetal bovine serum (FBS), human serum albumin (HSA) and mannitol were purchased from Beijing Solarbio Science & Technology Co., Ltd. (Beijing, China). Streptozotocin (STZ) was purchased from Sigma (Aldrich, USA). Trizol Reagent was purchased from Invitrogen

(Carlsbad, USA). A Reverse Transcription Kit was purchased from Thermo Scientific. SYBR[®] *Premix Ex Taq*[™] reagent was purchased from TaKaRa Biotechnology, Inc. (Otsu, Japan). Antibodies against TGF β 1 and collagen IV were purchased from ABCAM (Cambridge, UK). Antibodies against p38 MAPK and p-p38 MAPK were purchased from Cell Signaling Technology, Inc. (Boston, USA). Antibodies against β -tubulin were purchased from Tianjin Sungene Biotech Co., Ltd. (Tianjin, China). Horseradish peroxidase labelled goat anti-rabbit IgG was purchased from Beijing ComWin Biotech Co., Ltd. (Beijing, China). The biotinylated goat anti-rabbit antibody and the Vectastain Elite ABC Staining Kit were purchased from Vector Laboratories (Burlingame, CA, USA). All other materials were of reagent grade.

Purification of β 2GPI and preparation of reduced β 2GPI

β 2GPI was purified from normal human plasma using previously methods [20]. Plasma β 2GPI was precipitated by 3% perchloric acid and isolated by heparin-sepharose affinity chromatography (HiTrap Heparin, GE Healthcare). LC-MS analysis was used to confirm this protein. The purity of β 2GPI was confirmed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) on a 10% mini-gel. SDS-PAGE analysis of the protein sample showed the same band as the standard sample (**Figure 1**). The bicinchoninic acid (BCA) method was used to determine the concentration of β 2GPI. Reduced β 2GPI was prepared by previously methods [18]. Purified β 2GPI (1 μ M) was reduced by TRX-1 (3.5 μ M) activated with dithiothreitol (DTT, 70 μ M). The thiols of reduced β 2GPI were protected by reduced glutathione. The reduced β 2GPI was verified using a western blot and LC-MS analysis. Western blot analysis showed the new protein maintained the same immunologic activity as β 2GPI. LC-MS analysis verified that domain V of the new protein had free thiols.

Experimental animals for diabetic nephropathy

Female Balb/c mice (8-weeks old, 18-20 g) were obtained from Peking University Laboratory Animal Center (Beijing, China). All animal procedures were performed according to the *Guide for the Care and Use of Laboratory Animals of the National Institutes of Health*. All animals were housed under standard condi-

Reduced β 2GPI improve diabetic nephropathy




Figure 1. Verification of β 2GPI. After purification, the sample fractions were digested by trypsin PNGase F followed by LC-MS analysis and SDS-PAGE analysis. A. The red coloured sequence is the sequence we found. B. SDS-PAGE result revealed that the digested fraction was consistent with standard protein. Thus, we verified that the fractions dly contaed β 2GPI. C. After further purification with heparin-sepharose affinity chromatography, the sample fractions were analyzed with SDS-PAGE. The molecule weight of the sample was calculated according to the relationship between molecular weight of standard protein and relative mobility ratio, and the result was 50 kDa. SP, std: β 2GPI standard protein; M: marker; DE: digested by trypsin PNGase F.

tions with unrestricted food and water. Mice were randomly divided into a diabetes mellitus group ($n=40$) and a normal control group ($n=8$) and were fed a high fat and standard chow diet respectively. After 8 weeks, the DM group was induced by intraperitoneal injection of STZ at 80 mg/kg three times for three days as previously described [21]. The normal control group was injected with an equal amount of sodium citrate buffer. Diabetic mice were confirmed by a plasma glucose concentration ≥ 16.7 mM after 72 hours injection. The STZ-induced diabetic mice were randomly divided into 5 groups ($n=8$) as follows: (a) and (b) β 2GPI or reduced β 2GPI for 3 weeks in which mice were induced by an intravenous injection of 20 μ g of β 2GPI or reduced β 2GPI one time for 3 weeks; (c) and (d) β 2GPI or reduced β 2GPI for 6 weeks in which mice were injected with 20 μ g of β 2GPI or reduced β 2GPI two times for 6 weeks (one time per 3 weeks); (e) diabetic control mice in which mice were injected with equal volumes of PBS two times for 6 weeks. The normal control group was simultaneously injected with equal volumes of PBS two times for 6 weeks also. After STZ or control injections, the blood glu-

cose and body weight were monitored weekly. The mice were housed individually in metabolic cages. The 24 hours urine samples were collected on the day before the end of the experiment and were stored at -80°C . After 3 or 6 weeks of treatment, blood samples were collected from the retro-orbital venous plexus then mice were sacrificed and both kidneys were removed. Serum was collected by centrifugation at 3000 g for 15 min and stored at -80°C . Urine albumin concentration was determined using the mouse albumin ELISA kit (Assaypro, USA) according to the manufacturer's instructions. The serum levels of glucose, urea nitrogen and creatinine were determined using an Automatic Biochemical Analyzer (Hitachi Co., Ltd., Japan). Kidney samples were weighed and frozen in liquid nitrogen and stored at -80°C or fixed in 10% neutral-buffered formalin.

Cell culture and treatment

RMCs between passages 3 and 8 were used in experiments. RMCs were cultured in Dulbecco's modified Eagle Medium (DMEM) containing 5.5

Reduced β 2GPI improve diabetic nephropathy

mM glucose supplemented with 10% FBS and antibiotics, maintained at 37°C in a 95% air, 5% CO₂ atmosphere. RMCs were seeded at a density of 1.0×10⁶ cells/well in six-well plates. When the cells had grown to 70-80% confluence, RMCs were placed in serum-free DMEM for 24 hours and were then divided into eight groups for the next 24 hours as follows: (a) normal glucose in which RMCs were cultured in 5.5 mM glucose DMEM (NG group); (b) mannitol in which RMCs were cultured in medium composed of 19.5 mM mannitol and 5.5 mM glucose DMEM that was used as an osmotic control group (Mannitol group); (c) and (d) low or high-dose HSA in which RMCs were cultured in 25 mM glucose DMEM supplemented with 40 or 80 μ g/ml HSA (HG plus HSA group) that was used as the high glucose control group; (e) and (f) low or high-dose β 2GPI in which RMCs were cultured in 25 mM glucose DMEM supplemented with 40 or 80 μ g/ml β 2GPI (HG plus β 2GPI group); and (g) and (h) low or high-dose reduced β 2GPI in which RMCs were cultured in 25 mM glucose DMEM supplemented with 40 or 80 μ g/ml reduced β 2GPI (HG plus reduced β 2GPI group). Cell lysates and RNA were collected and used for quantitative real-time PCR and western blot analysis.

Morphological and immunohistochemical analysis

The kidneys were embedded in paraffin and were sliced into 4 μ m sections. The tissue slices were stained with periodic acid-Schiff (PAS) for general histological studies. For immunohistochemical analysis, the paraffin sections were deparaffinized and rehydrated. After blockage of endogenous peroxidase and antigen retrieval, the tissue slices were incubated with primary antibodies (TGF- β 1 or collagen IV, diluted 1:200) at 4°C overnight. The slices were then washed with PBS and incubated with a biotinylated second antibody (diluted 1:200) (Vector, Burlingame, CA) at room temperature for 30 min. The slices were washed again with PBS and incubated with a Vectastain ABC kit (Vector, UK) according to the manufacturer's instructions. Immunoreactivity was visualized with the chromogen 3,3'-diaminobenzidine (DAB, Sigma). The slices were counterstained with hematoxylin, dehydrated and mounted. Brown areas were judged as positive staining. Glomerular cross-sectional areas were assessed with light microscopy (Olympus IX-52, Olympus Optical, Tokyo, Japan). Semi-

quantitative analysis of the ratio of positive staining areas to the glomerular areas was evaluated with the image analysis software Image Pro. Plus 6.0 (Media Cybernetics, Inc., Bethesda, MD, USA).

Quantitative real-time RT-PCR

Total RNA was extracted from kidneys or cultured cells with Trizol Regent and was reverse transcribed into cDNA with a Reverse Transcription Kit. RNA was quantitated by OD values at 260 nm and the integrity was verified by ethidium bromide staining of ribosomal 18S and 28S bands on agarose gels. Real-time PCR was performed with the SYBR[®] *Premix Ex Taq*[™] reagent according to the manufacturer's instructions. The following primers were used for TGF- β 1 of STZ-diabetic mice: forward: 5'-AGG GCT ACC ATG CCA ACT TC-3', reverse: 5'-CCA CGT AGT AGA CGA TGG GC-3'; for TGF- β 1 of RMCs: forward: 5'-AGG GCT ACC ATG CCA ACT TC-3', reverse: 5'-CCA CGT AGT AGA CGA TGG GC-3'; for collagen IV of STZ-diabetic mice: forward 5'-TCC AGG CCC CCC TGG AAC TGT-3', reverse 5'-GAG GGC CTG GTT GGC CTG-3'; for collagen IV of RMCs: forward 5'-CCA TCT GTG GAC CAT GGC TT-3', reverse 5'-GCG AAG TTG CAG ACG TTG TT-3'; for GAPDH: forward: 5'-CAA GGT CAT CCA TGA CAA CTT TG-3', reverse: 5'-GTC CAC CAC CCT GTT GCT GTA G-3'. Water, as a replacement for cDNA, was used as negative control. GAPDH was used as an internal control. The mRNA expression levels were quantified with normalization to GAPDH. The results were calculated using the 2^{- $\Delta\Delta$ CT} method.

Western blot

The renal cortex tissue homogenates and RMCs after being washed with ice-cold PBS were lysed with RIPA lysis buffer. After incubation for 30 min on ice, the lysates were centrifuged at 12000 rpm for 15 min at 4°C and the supernatant was collected. The concentration of protein in the supernatant was determined with the Micro BCA[™] Protein Assay Reagent Kit according to the manufacturer's instructions. The samples (30 μ g protein per lane) were separated on a 10% SDS-PAGE and blotted onto a nitrocellulose transfer membrane. The membranes were blocked with 5% nonfat milk in TBST (20 mM Tris-HCl, pH 7.6, 137 mM NaCl and 0.01% Tween-20) for 60 min at room temperature and were then incubated with primary

Reduced β 2GPI improve diabetic nephropathy




Figure 2. Treatment with β 2GPI and reduced β 2GPI prevented mesangial expansion and capillary basement membrane thickening in STZ-induced diabetic mice. Glomerular histopathology analysis by PAS stain. A. The pictures display representative glomeruli of PAS-stained sections with original magnification of $\times 400$. Scale bar = 50 μ m for all microrahs. B. The ratio of PAS-positive staining areas to the glomerular areas was evaluated by a semi-quantitative method. Data are expressed as the mean \pm S.E.M. (n=8 for each group). * $P < 0.05$ vs. NC roup, # $P < 0.05$ vs. DM group. NC: normal control; DM: diabetic model control; β 2GPI: 20 μ g β 2GPI; reduced β 2GPI: 20 μ g reduced β 2GPI.

antibodies (against TGF- β 1, collagen IV, p38 MAPK, p-p38 MAPK and β -tubulin) overnight at 4°C. The membranes were subsequently washed with TBST and incubated with horse-radish peroxidase-conjugated secondary antibodies for 60 min at room temperature. After the incubation, the membranes were washed with TBST. Immunoreactive bands were detected with ECL reagents (Amersham Biosciences, UK) and exposure to Hyperfilm (GE Healthcare, UK). Band densities were measured using BandScan software and quantified by normalization to β -tubulin.

Statistical analysis

All of the results are expressed as the means \pm SEM. Statistical analyses were performed using IBM SPSS Statistics 19.0 software. Statistical significance among multi-groups was determined using one way analysis of vari-

ance and Tukey's test post hoc analysis. Statistical significance between two groups was determined using the unpaired Student's t-test. $P < 0.05$ was considered statistically significant.

Results

β 2GPI and reduced β 2GPI improve biochemical parameters and kidney hypertrophy in STZ-induced diabetic mice

After STZ injection, all mice became diabetic with significantly higher urine albumin, urea nitrogen and creatinine compared with the normal control group ($P < 0.05$) (Table 1). However, the levels of urea nitrogen, creatinine and urine albumin were significantly lower in diabetic mice treated with β 2GPI and reduced β 2GPI compared with untreated diabetic mice ($P < 0.05$), except in mice treated with β 2GPI for 3 weeks.

Reduced β 2GPI improve diabetic nephropathy

Table 1. The effects of β 2GPI and reduced β 2GPI on the biochemical parameters and renal function of STZ-induced diabetic mice at the end of the experimental period. Data are expressed as the mean \pm S.E.M. (n = 8 for each group)

Groups	NC	DM	DM + β 2-GPI		DM + reduced β 2GPI	
			3 wk	6 wk	3 wk	6 wk
Body weight (g)	24.58 \pm 2.53	22.48 \pm 3.11	23.39 \pm 1.65	24.04 \pm 2.83	24.33 \pm 2.76	23.72 \pm 2.42
Kidney weight (g)	134.69 \pm 19.40	263.09 \pm 15.81*	212.02 \pm 22.41*	205.30 \pm 22.69* [#]	210.45 \pm 30.49*	186.34 \pm 15.04 [#]
Kidney/body weight ratio (mg/g)	5.57 \pm 1.40	11.88 \pm 1.93*	9.04 \pm 0.33*	8.56 \pm 0.64* [#]	8.64 \pm 0.59* [#]	7.88 \pm 0.63 [#]
Blood glucose (mmol/L)	6.05 \pm 0.86	25.66 \pm 3.21*	23.89 \pm 6.43*	23.78 \pm 6.21*	24.49 \pm 5.55*	22.40 \pm 4.28*
Serum creatinine (μ mmol/L)	7.29 \pm 1.23	23.64 \pm 5.07*	17.17 \pm 3.06* [#]	15.98 \pm 5.89* [#]	16.14 \pm 3.93* [#]	13.20 \pm 1.71* [#]
Blood urea nitrogen (mmol/L)	5.23 \pm 0.65	12.57 \pm 1.24*	9.75 \pm 1.22*	8.10 \pm 0.84* [#]	8.60 \pm 1.43* [#]	8.33 \pm 1.34* [#]
Urine albumin (μ g/24 h)	27.26 \pm 8.74	230.99 \pm 40.01*	161.41 \pm 31.03*	132.88 \pm 28.61* [#]	140.58 \pm 25.72* [#]	116.88 \pm 18.52* [#]

C: normal control; DM: diabetic model control; β 2GPI: 20 μ g β 2GPI; reduced β 2GPI: 20 μ g reduced β 2GPI. NC and DM were treated with equal volumes of PBS. All the mice were administered via intravenous injection. * P <0.05 vs. NC group. [#] P <0.05 vs. DM group.

Reduced β 2GPI improve diabetic nephropathy




Figure 3. Treatment with β 2GPI and reduced β 2GPI decreased the excessive accumulation of ECM proteins in STZ-induced diabetic mice. A. Immunohistochemical staining of collagen IV in the above groups. Increased expression of collagen IV-positive areas (dark brown) was observed after STZ induction in the DM group and was reduced with β 2GPI and reduced β 2GPI as the treatment increasing. Original magnification: $\times 400$. Scale bar = 50 μ m for all micrographs. NC: normal control; DM: diabetic model control; β 2GPI: 20 μ g β 2GPI; reduced β 2GPI: 20 μ g reduced β 2GPI. B and C. Quantification of collagen IV expression in the kidneys was performed using quantitative real-time RT-PCR and western blot. The mRNA and protein expression of collagen IV in kidneys were significantly increased after STZ induction in the DM group and were reduced with β 2GPI, evidently with reduced β 2GPI for 6 weeks. Data are expressed as the mean \pm S.E.M. (n=8 for each group). * $P < 0.05$ vs. NC group, # $P < 0.05$ vs. DM group. N: normal control; DM: diabetic model control; β 2GPI: 20 μ g β 2GPI; reduced β 2GPI: 20 μ g reduced β 2GPI.

The kidney hypertrophy index (kidney/body weight ratio) increased significantly ($P < 0.05$), while the body weight slightly decreased in diabetic mice compared with the normal control group ($P > 0.05$) (Table 1). However, the degree of kidney hypertrophy index decreased after treatment with β 2GPI and reduced β 2GPI, particularly in mice treated with reduced β 2GPI for 6 weeks.

β 2GPI and reduced β 2GPI improved glomerular morphological changes in STZ-induced diabetic mice

The mesangial expansion and capillary basement membrane thickening were observed in

the diabetic control group after PAS staining (Figure 2). After treatment with β 2GPI and reduced β 2GPI, there was decreased mesangial expansion and capillary basement membrane thickening. As treatment time progressed, the improvements were evident. Particularly, after 6 weeks treatment with reduced β 2GPI, the PAS staining showed nearly normal glomerular structure.

β 2GPI and reduced β 2GPI decreased the excessive accumulation of ECM proteins in STZ-induced diabetic mice

The STZ-induced diabetic mice exhibited an increase in collagen IV-positive areas accumu-

Reduced β 2GPI improve diabetic nephropathy




Figure 4. Treatment with β 2GPI and reduced β 2GPI inhibited the activation of TGF- β 1 and p38 MAPK in STZ-induced diabetic mice. A and B. Quantification of TGF- β 1 expression in the kidneys was performed using quantitative real-time RT-PCR and western blot. The mRNA and protein expression of TGF- β 1 in kidneys were significantly increased after STZ induction in the DM group and were reduced with β 2GPI, evidently with reduced β 2GPI for 6 weeks. C. Quantification of p38MAPK

and phospho-p38 MAPK expression in the kidneys was performed using western blot. The phosphorylation level of p38 MAPK significantly increased in the diabetic control group, whereas β 2GPI and reduced β 2GPI treatment inhibited p38 MAPK phosphorylation. Data are expressed as the mean \pm S.E.M. (n=8 for each group). * P <0.05 vs. group, # P <0.05 vs. DM group. N: normal control; DM: diabetic model control; β 2GPI: 20 μ g β 2GPI; reduced β 2GPI: 20 μ g reduced β 2GPI.

lated in the mesangial matrix and capillary basement membranes compared with the normal control group (Figure 3A). After 3 weeks treatment with β 2GPI and reduced β 2GPI, collagen IV-positive areas decreased slightly compared with the diabetic control group. However, 6 weeks treatment with β 2GPI and reduced β 2GPI exhibited significantly less collagen IV-positive areas. In addition, collagen IV mRNA and protein expression from the diabetic control group were significantly higher than those from the normal control group (Figure 3B and 3C). After β 2GPI and reduced β 2GPI treatment, collagen IV mRNA and protein expression decreased significantly compared with the diabetic control group, particularly after 6 weeks treatment with reduced β 2GPI.

β 2GPI and reduced β 2GPI inhibited the activation of TGF- β 1 and p38 MAPK in STZ-induced diabetic mice

TGF- β 1 mRNA and protein expression were significantly higher in the diabetic control group compared with the normal control group (Figure 4A and 4B). Although TGF- β 1 mRNA expression was higher than the normal control group after treatment, it was lower than the diabetic control group. Meanwhile, the β 2GPI and reduced β 2GPI treated diabetic mice had decreased TGF- β 1 protein expression compared with the diabetic and normal control groups. The reduction of TGF- β 1 expression was most evident in the diabetic

Reduced β 2GPI improve diabetic nephropathy





Figure 5. Treatment with β 2GPI and reduced β 2GPI decreased the expression of collagen IV in high glucose-induced RMCs. A and B. Quantification of collagen IV expression in high glucose-induced RMCs was performed using quantitative real-time RT-PCR and western blot. The mRNA and protein expression of collagen IV were significantly increased after high glucose stimulation. However, the elevated expression levels decreased after β 2GPI and reduced β 2GPI treatment with high glucose stimulation. Data are expressed as the mean \pm S.E.M.. * P <0.05 vs. normal glucose group, # P <0.05 vs. high glucose control group. NG: normal glucose group; Mannitol: osmotic control group; HSA: high glucose with HSA as high glucose control group; β 2GPI: high glucose with β 2GPI; reduced β 2GPI: high glucose with reduced 2GPI.

mice treated with reduced β 2GPI for 6 weeks. In addition, the phosphorylation level of p38

MAPK significantly increased in the diabetic control group, whereas β 2GPI and reduced

Reduced β 2GPI improve diabetic nephropathy



Reduced β 2GPI improve diabetic nephropathy

Figure 6. Treatment with β 2GPI and reduced β 2GPI inhibited the activation of TGF- β 1 and p38 MAPK in high glucose-induced RMs. A and B. Quantification of TGF- β 1 expression in high glucose-induced RMCs was performed using quantitative real-time RT-PCR and western blot. The mRNA and protein expression of TGF- β 1 were significantly increased after high glucose stimulation and were significantly decreased with a high-dose of β 2GPI and reduced β 2GPI compared with a low-dose. C. Quantification of p38MAPK and phospho-p38 MAPK expression in high glucose-induced RMCs was performed using western blot. The phosphorylation level of p38 MAPK significantly increased after high glucose stimulation, whereas β 2GPI and reduced β 2GPI treatment inhibited p38 MAPK phosphorylation. Data are expressed as the mean \pm S.E.M. (n=8 for each group). * P <0.05 v. N group, # P <0.05 vs. DM group. N: normal control; DM: diabetic model control; β 2GPI: 20 μ g β 2GPI; reduced β 2GPI: 20 μ g reduced β 2GPI.

β 2GPI treatment inhibited p38 MAPK phosphorylation (**Figure 4C**). It was noted that the phosphorylation level of p38 MAPK in treated diabetic mice was lower than the normal control group.

β 2GPI and reduced β 2GPI decreased the expression of collagen IV in high glucose-induced RMCs

Collagen IV mRNA and protein expression were significantly increased after high glucose stimulation for 24 hours (**Figure 5A and 5B**). These elevated expression levels decreased after β 2GPI and reduced β 2GPI treatment (P <0.05). As a hyperosmotic control, the mannitol group demonstrated no difference compared with the normal glucose group. Incredibly, the protein expression of collagen IV in β 2GPI and reduced β 2GPI groups were lower than the normal glucose group. Furthermore, collagen IV mRNA expression was slightly lower in the high-dose of β 2GPI and reduced β 2GPI compared with the low-dose groups. Collagen IV protein expression was lower in the high-dose reduced β 2GPI group compared with the low-dose group.

β 2GPI and reduced β 2GPI inhibited the activation of TGF β 1 and p38 MAPK in high glucose-induced RMCs

High glucose stimulation increased TGF- β 1 mRNA and protein expression compared with the normal glucose and mannitol groups (**Figure 6A and 6B**). Although TGF- β 1 mRNA expression was higher than the normal glucose and mannitol group after β 2GPI and reduced β 2GPI treatment, it was lower than the high glucose group. TGF- β 1 mRNA expression levels were significantly decreased with a high-dose of β 2GPI and reduced β 2GPI compared with a low-dose groups. Furthermore, TGF- β 1 protein expression was significantly lower after β 2GPI and reduced β 2GPI treatment compared with

the normal glucose group. In addition, the phosphorylation level of p38 MAPK significantly increased after high glucose stimulation. It was noted that p38 MAPK phosphorylation levels in the high glucose-induced RMCs treated with reduced β 2GPI were lower than the normal glucose group, whereas treatment with β 2GPI slightly inhibited p38 MAPK phosphorylation compared with the normal glucose group (**Figure 6C**).

Discussion

In the present study, we demonstrated that diabetic mice induced by a high-fat diet and STZ exhibited a number of early clinical and pathological characteristics of DN. Treatment with β 2GPI and reduced β 2GPI improved the above changes with time dependently, indicating their renoprotective effects during DN development. *In vivo* and *in vitro* studies have demonstrated that hyperglycemia can induce enhanced synthesis of ECM proteins such as collagen and fibronectin [22-24]. In this study, we determined that collagen IV expression was significantly increased in STZ-induced diabetic mice and high glucose-induced RMCs, which is consistent with previous reports. However, treatment with β 2GPI and specifically reduced β 2GPI effectively decreased collagen IV expression, indicating that β 2GPI and reduced β 2GPI could improve kidney fibrosis and glomerulosclerosis through the suppression of ECM proteins synthesis.

TGF- β 1 has been recognized as an important inflammatory cytokine in DN as previously mentioned. The inhibition of TGF- β 1 expression enhances DN treatment by suppressing matrix accumulation. Therefore, TGF- β 1 has been proposed as an intervention target for DN treatment [25]. In the present study, the STZ-induced diabetic mice and the high glucose-induced RMCs had increased TGF- β 1 mRNA and protein expression. However, treatment with β 2GPI and

Reduced β 2GPI improve diabetic nephropathy

reduced β 2GPI could inhibit TGF- β 1 mRNA and protein expression.

The activation of p38 MAPK, one of the downstream effectors of the TGF- β 1 signaling cascades, has been reported to be involved in the progression of DN [26]. The effects of β 2GPI and reduced β 2GPI on TGF- β 1-p38 MAPK activation in DN have not been elucidated. The phosphorylation levels of p38 MAPK significantly increased in the present study, which were positively correlated with TGF- β 1 protein. The β 2GPI and reduced β 2GPI treatment decreased p38 MAPK phosphorylation levels as well as TGF- β 1 expression in STZ-induced diabetic mice, and similar results were observed in high glucose-induced RMCs.

The proportion of reduced β 2GPI is significantly lower in the antiphospholipid syndrome group than that in healthy individuals [27], which suggests that reduced β 2GPI may play a protective role in our bodies. Reduced β 2GPI was recently found to protect EAhy926 (fusion of HUVECs and the A549 carcinoma cell line) from oxidative stress induced endothelial cell damage and to display increased binding to von Willebrand factor (VWF) than non-reduced β 2GPI *in vitro* [28]. At present, no studies have reported the correlation between β 2GPI and reduced β 2GPI and DN. Our study first indicated that both β 2GPI and reduced β 2GPI improved kidney fibrosis and decreased mesangial cells produce collagen IV by inhibiting TGF- β 1 and p38 MAPK phosphorylation expression. Although reduced β 2GPI have the better effect on kidney fibrosis, β 2GPI have the same renoprotective effect. It is possible that β 2GPI can switch between an oxidized and reduced state under harsh conditions. As described previously, domain V of β 2GPI can be reduced by TRX-1 resulting the functional disulfide bond (Cys288-Cys326) opened, resulting in some functional changes. The opened functional disulfide bond could be anchored to the cells membrane, providing the appropriate interface to react with cell surface proteins such as TGF- β 1, resulting inhibiting the activation of TGF- β 1-p38 MAPK signal pathway. TRX-1 is ubiquitously expressed and is also present on mesangial cells in DN [29]. Given β 2GPI's high concentration in plasma makes it easily available for reactions with TRX-1, resulting in the reduced formation of β 2GPI. Thus, in this study β 2GPI and reduced

β 2GPI have been showed the similar renoprotective and antifibrosis effects.

In summary, the present study confirmed that β 2GPI and reduced β 2GPI improved renal dysfunction and kidney fibrosis as well as decreased collagen IV and TGF- β 1 mRNA and protein expression in STZ-induced diabetic mice and high glucose-induced RMCs. Moreover, the present studies demonstrated that the renoprotective and antifibrosis effects of β 2GPI and reduced β 2GPI in DN were closely associated with suppressing the activation of the TGF- β 1-p38 MAPK pathway. Currently, the function of β 2GPI and reduced β 2GPI has been poorly understood, the definitive molecular mechanism of β 2GPI and reduced β 2GPI inhibition of the TGF- β 1-p38 MAPK pathway in DN needs to be investigated in future.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (No. 30971393 and No. 81070645), the Tianjin natural science fund (No. 10JCYBJC12000).

Disclosure of conflict of interest

None.

Address correspondence to: Pei Yu, 2011 Collaborative Innovation Center of Tianjin for Medical Epigenetics, The Key Laboratory of Hormones and Development (Ministry of Health), Metabolic Diseases Hospital & Tianjin Institute of Endocrinology, Tianjin Medical University, Tianjin 300070, China. Tel: +86 13821008270; Fax: +86 22 23333222; E-mail: peiyudoc@yeah.net

References

- [1] Rossing P. Diabetic nephropathy: worldwide epidemic and effects of current treatment on natural history. *Curr Diab Rep* 2006; 6: 479-483.
- [2] Raptis AE, Viberti G. Pathogenesis of diabetic nephropathy. *Exp Clin Endocrinol Diabetes* 2001; 109 Suppl 2: S424-S437.
- [3] Seger R, Krebs EG. The MAPK signaling cascade. *Faseb J* 1995; 9: 726-735.
- [4] Choi ME, Ding Y, Kim SI. TGF-beta signaling via TAK1 pathway: role in kidney fibrosis. *Semin Nephrol* 2012; 32: 244-252.
- [5] Schultze HE, Heide K, Haupt H. Über ein bisher unbekanntes niedermolekulares Beta2-

Reduced β 2GPI improve diabetic nephropathy

- Globulin des Humanserums, Vol. 48, p 719, 1961.
- [6] McNeil HP, Simpson RJ, Chesterman CN, Krilis SA. Anti-phospholipid antibodies are directed against a complex antigen that includes a lipid-binding inhibitor of coagulation: beta 2-glycoprotein I (apolipoprotein H). *Proc Natl Acad Sci U S A* 1990; 87: 4120-4124.
- [7] Rioche M, Maseyeff R. Synthesis of plasma beta 2 glycoprotein I by human hepatoma cells in tissue culture. *Biomedicine* 1974; 21: 420-423.
- [8] Schwarzenbacher R, Zeth K, Diederichs K, Gries A, Kostner GM, Laggner P, Prassl R. Crystal structure of human beta2-glycoprotein I: implications for phospholipid binding and the antiphospholipid syndrome. *Embo J* 1999; 18: 6228-6239.
- [9] de Groot PG, Meijers JC. beta(2) -Glycoprotein I: evolution, structure and function. *J Thromb Haemost* 2011; 9: 1275-1284.
- [10] Petri M. Update on anti-phospholipid antibodies in SLE: the Hopkins' Lupus Cohort. *Lupus* 2010; 19: 419-423.
- [11] Agar C, de Groot PG, Marquart JA, Meijers JC. Evolutionary conservation of the lipopolysaccharide binding site of beta(2)-glycoprotein I. *Thromb Haemost* 2011; 106: 1069-1075.
- [12] Jankowski M, Vreys I, Wittevrongel C, Boon D, Vermynen J, Hoylaerts MF, Arnout J. Thrombogenicity of beta 2-glycoprotein I-dependent antiphospholipid antibodies in a photochemically induced thrombosis model in the hamster. *Blood* 2003; 101: 157-162.
- [13] Arad A, Proulle V, Furie RA, Furie BC, Furie B. beta(2)-Glycoprotein-1 autoantibodies from patients with antiphospholipid syndrome are sufficient to potentiate arterial thrombus formation in a mouse model. *Blood* 2011; 117: 3453-3459.
- [14] Nilsson M, Wasyluk S, Morgelin M, Olin AI, Meijers JC, Derksen RH, de Groot PG, Herwald H. The antibacterial activity of peptides derived from human beta-2 glycoprotein I is inhibited by protein H and M1 protein from *Streptococcus pyogenes*. *Mol Microbiol* 2008; 67: 482-492.
- [15] Vaarala O. Antiphospholipid antibodies and atherosclerosis. *Lupus* 1996; 5: 442-447.
- [16] Staub HL, Franck M, Ranzolin A, Norman GL, Iverson GM, von Muhlen CA. IgA antibodies to beta2-glycoprotein I and atherosclerosis. *Autoimmun Rev* 2006; 6: 104-106.
- [17] de Laat B, de Groot PG, Derksen RH, Urbanus RT, Mertens K, Rosendaal FR, Doggen CJ. Association between beta2-glycoprotein I plasma levels and the risk of myocardial infarction in older men. *Blood* 2009; 114: 3656-3661.
- [18] Ioannou Y, Zhang JY, Passam FH, Rahgozar S, Qi JC, Giannakopoulos B, Qi M, Yu P, Yu DM, Hogg PJ, Krilis SA. Naturally occurring free thiols within beta 2-glycoprotein I in vivo: nitrosylation, redox modification by endothelial cells, and regulation of oxidative stress-induced cell injury. *Blood* 2010; 116: 1961-1970.
- [19] Yu P, Passam FH, Yu DM, Denyer G, Krilis SA. Beta2-glycoprotein I inhibits vascular endothelial growth factor and basic fibroblast growth factor induced angiogenesis through its amino terminal domain. *J Thromb Haemost* 2008; 6: 1215-1223.
- [20] Lin KY, Wang HH, Lai ST, Pan JP, Chiang AN. beta(2)-glycoprotein I protects J774A.1 macrophages and human coronary artery smooth muscle cells against apoptosis. *J Cell Biochem* 2005; 94: 485-496.
- [21] Zhang R, Zhou SJ, Li CJ, Wang XN, Tang YZ, Chen R, Lv L, Zhao Q, Xing QL, Yu DM, Yu P. C-reactive protein/oxidised low-density lipoprotein/beta2-glycoprotein I complex promotes atherosclerosis in diabetic BALB/c mice via p38mitogen-activated protein kinase signal pathway. *Lipids Health Dis* 2013; 12: 42.
- [22] Noh H, Ha H, Yu MR, Kang SW, Choi KH, Han DS, Lee HY. High glucose increases inducible NO production in cultured rat mesangial cells. Possible role in fibronectin production. *Nephron* 2002; 90: 78-85.
- [23] Yu Y, Lyons TJ. A lethal tetrad in diabetes: hyperglycemia, dyslipidemia, oxidative stress, and endothelial dysfunction. *Am J Med Sci* 2005; 330: 227-232.
- [24] Mahimainathan L, Das F, Venkatesan B, Choudhury GG. Mesangial cell hypertrophy by high glucose is mediated by downregulation of the tumor suppressor PTEN. *Diabetes* 2006; 55: 2115-2125.
- [25] McGowan TA, Zhu Y, Sharma K. Transforming growth factor-beta: a clinical target for the treatment of diabetic nephropathy. *Curr Diab Rep* 2004; 4: 447-454.
- [26] Hayashida T, Poncelet AC, Hubchak SC, Schnaper HW. TGF-beta1 activates MAP kinase in human mesangial cells: a possible role in collagen expression. *Kidney Int* 1999; 56: 1710-1720.
- [27] Ioannou Y, Zhang JY, Qi M, Gao L, Qi JC, Yu DM, Lau H, Sturgess AD, Vlachoyiannopoulos PG, Moutsopoulos HM, Rahman A, Pericleous C, Atsumi T, Koike T, Heritier S, Giannakopoulos B, Krilis SA. Novel assays of thrombogenic pathogenicity in the antiphospholipid syndrome based on the detection of molecular oxidative modification of the major autoantigen beta2-glycoprotein I. *Arthritis Rheum* 2011; 63: 2774-2782.

Reduced β 2GPI improve diabetic nephropathy

- [28] Passam FH, Rahgozar S, Qi M, Raftery MJ, Wong JW, Tanaka K, Ioannou Y, Zhang JY, Gemmell R, Qi JC, Giannakopoulos B, Hughes WE, Hogg PJ, Krilis SA. Redox control of beta2-glycoprotein I-von Willebrand factor interaction by thioredoxin-1. *J Thromb Haemost* 2010; **8**: 1754-1762.
- [29] Advani A, Gilbert RE, Thai K, Gow RM, Langham RG, Cox AJ, Connelly KA, Zhang Y, Herzenberg AM, Christensen PK, Pollock CA, Qi W, Tan SM, Parving HH, Kelly DJ. Expression, localization, and function of the thioredoxin system in diabetic nephropathy. *J Am Soc Nephrol* 2009; **20**: 730-741.