

## Erratum

# MiR-30a-5p suppresses cell growth and enhances apoptosis of hepatocellular carcinoma cells via targeting AEG-1: Int J Clin Exp Pathol. 2015; 8(12): 15632-15641

Rongquan He<sup>1\*</sup>, Lihua Yang<sup>1\*</sup>, Xiaomiao Lin<sup>2</sup>, Xin Chen<sup>3</sup>, Xinggu Lin<sup>4</sup>, Fanglin Wei<sup>3</sup>, Xiaona Liang<sup>3</sup>, Yihuan Luo<sup>3</sup>, Yuzhuang Wu<sup>3</sup>, Tingqing Gan<sup>1</sup>, Yiwu Dang<sup>3</sup>, Gang Chen<sup>3</sup>


<sup>1</sup>Department of Medical Oncology, The First Affiliated Hospital of Guangxi Medical University, Nanning, P. R. China;

<sup>2</sup>Department of Children Rehabilitation Medicine, Guangxi Maternal and Child Health Hospital, Nanning, Guangxi Zhuang Autonomous Region, P. R. China; <sup>3</sup>Department of Pathology, The First Affiliated Hospital of Guangxi Medical University, Nanning, P. R. China; <sup>4</sup>Key Laboratory for Genomic and Personalized Medicine, Guangxi Medical University, Nanning, P. R. China. \*Equal contributors.

Received February 9, 2020; Accepted March 26, 2020; Epub May 1, 2020; Published May 15, 2020

In this article published in IJCEP, the authors reported that they had misspelled the name of a cell line and supplied an incorrect picture for publication. The correct name of the cell line should be 'Hep3B', instead of HepB3. The correct **Figure 5** is shown below. The published results and conclusions are not affected by the errata. The authors sincerely apologize for the errors.

**Address correspondence to:** Drs. Yiwu Dang and Gang Chen, Department of Pathology, First Affiliated Hospital of Guangxi Medical University, 6 Shuangyong Road, Nanning 530021, Guangxi Zhuang Autonomous Region, China. E-mail: dangyiwu@126.com (YWD); chen\_gang\_triones@163.com (GC)



**Figure 5.** HCC HepG2 cell morphology is influenced by miR-30a-5p. HepG2 cells ( $2.5 \times 10^3$  cells per well in 96-well-plate) were cultured for 24 h and transfected with miR-30a-5p inhibitor, miR-30a-5p mimic, AEG-1 siRNA, and their negative controls (200 nM) for another 96 h. Change of morphology was evaluated using Hoechst 33342/propidium iodide (PI) double fluorescent chromatin staining. A. Negative inhibitor control; B. miR-30a-5p inhibitor; C. Negative mimic control; D. miR-30a-5p mimic; E. Scrambled siRNAs; F. AEG-1 siRNAs.