

Original Article

Association of the SPT2 chromatin protein domain containing 1 gene rs17579600 polymorphism and serum lipid traits

Tao Guo¹, Rui-Xing Yin¹, Yuan Bin¹, Rong-Jun Nie¹, Xia Chen¹, Shang-Ling Pan²

¹Department of Cardiology, Institute of Cardiovascular Diseases, The First Affiliated Hospital, Guangxi Medical University, Nanning 530021, Guangxi, China; ²Department of Pathophysiology, School of Premedical Sciences, Guangxi Medical University, Nanning 530021, Guangxi, People's Republic of China

Received August 10, 2015; Accepted September 22, 2015; Epub October 1, 2015; Published October 15, 2015

Abstract: SPT2 chromatin protein domain containing 1 gene (*SPTY2D1*) is a candidate gene for dyslipidemia. The single nucleotide polymorphism (SNP) of rs7934205 near *SPTY2D1* locus was ethnic- and sex-specific associated with serum lipid levels in our previous study. Whether *SPTY2D1* rs17579600 SNP and several environmental factors are associated with serum lipid profiles is unknown. A total of 712 participants of Han and 689 unrelated individuals of Mulao were included. The genotype and allele frequencies of *SPTY2D1* rs17579600 SNP were different between the Han and Mulao populations (TT, 74.3% vs. 55.7%; TC, 17.6% vs. 31.2%, CC, 8.1% vs. 13.1%, $P = 0.028$; T, 83.1% vs. 71.3%; C, 16.9% vs. 28.7%, $P = 0.044$), and between males and females in the both ethnic groups. The levels of serum apolipoprotein (Apo) A1 in Han, triglyceride (TG) in Mulao, and total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), ApoA1 and ApoB in Mulao males were difference among the genotypes. The C allele carriers had higher ApoA1 in Han, lower TG in Mulao, and lower TC, LDL-C and ApoB and higher ApoA1 in Mulao males than the C allele non-carriers. Serum lipid parameters were also associated with several environmental factors in both ethnic groups. The differences suggesting there may be a racial/ethnic- and/or sex-specific association between the *SPTY2D1* rs17579600 SNP and serum lipid parameters in some ethnic groups.

Keywords: Lipids, SPT2 chromatin protein domain containing 1 gene, single nucleotide polymorphism, environmental factors

Introduction

Cardiovascular disease (CVD) is the world's leading cause of mortality, morbidity, disability, functional decline, and healthcare costs [1, 2]. The 2011 overall rate of death attributable to CVD was 229.6 per 100 000 Americans. The death rates were 275.7 for males and 192.3 for females. The rates were 271.9 for white males, 352.4 for black males, 188.1 for white females, and 248.6 for black females [3]. To establish risk status measurement of a standard lipid profile, including total cholesterol (TC) [4], triglycerides (TG) [5], LDL (low-density lipoprotein) cholesterol [6], apolipoprotein (Apo) B [7], HDL (high-density lipoproteins) cholesterol [8], ApoA1 [9] and the ratio ApoA1 to ApoB [10] is recommended from an integral component of approaches to cardiovascular risk prediction. This is well illustrated by the widespread popu-

larity of the metabolic syndrome concept [11], a constellation of risk factors that confers an elevated risk of cardiometabolic anomalies and CVD. Decades of research on common CVD risk factors have established that they often differ between men and women [12], are influenced by age [13] and ethnicity [14], and are modulated by behavioral choices [15], including poor diet [16] and a sedentary lifestyle [17], environmental conditions [18], and one's genetic profile [19, 20]. Even though it is well recognized that all these risk factors taken individually are characterized by a significant genetic component, there are uncertainties about the true magnitude of risk factor clustering, as well as on the role of genetic factors in risk factor clustering in individuals.

Using several twin and family study designs, studies have revealed that there is a significant

genetic component to human variability in CVD risk factors when considered individually [21, 22]. These risks above mentioned factors are all characterized by familial resemblance and significant heritability estimates [23, 24]. All these CVD risk factors have been the target of genome-wide association studies (GWAS) aimed at identifying common single nucleotide polymorphisms (SNPs) and at quantifying how much of the phenotypic variance is actually captured by them [25].

Several GWAS have reported the association of many SNPs near the SPT2 chromatin protein domain containing 1 gene (SPTY2D1; previous symbols & names: SPT2, Suppressor of Ty, domain containing 1 (*S. cerevisiae*), Gene ID: 144108, HGNC ID: 26818, synonyms: DKFZp686i068, FLJ39441, Spt2, locus type: gene with protein product, chromosomal location: 11p15.1) with one or more lipid traits [26-28]. SPT2 is a DNA binding protein with HMG-like domains. Functional domains of the yeast chromatin protein SPT2 can bind four-way junction and crossing DNA structures [29]. It plays a role in chromatin modulations associated with transcription elongation in *Saccharomyces cerevisiae* [30]. Vertebrate SPT2 is a representative of a new class of nucleolar histone chaperones, which associate with chromatin by their DNA-binding activity and function as nucleosome assembly/disassembly factors in the regulation of rDNA transcription [31]. A previous GWAS on plasma-lipid levels has identified the rs10128711 SNP near the SPTY2D1 as TC-related loci in European [28]. A sex-stratified analysis of other variant in SPTY2D1, rs7934205, in our previous study in Chinese has shown that the association between the SPTY2D1 rs7934205 SNP and serum lipid levels might have ethnic- and/or sex-specificity [32]. Whether SPTY2D1 rs17579600 SNP is associated with serum lipid levels or whether it exhibits ethnic and/or sex specific association like the previously reported SPTY2D1 SNPs remains elusive.

Mulao nationality, as one of the minorities (Han is the largest one), is a relatively conservative and isolated minority, and preserves their custom of intra-ethnic marriage. Interestingly, they have their culture of consanguineous marriage to cousins of maternal side, suggesting that the genetic background of Mulao population may be less heterogeneous within the population.

The recent molecular anthropological data showed that Mulao has much closer genetic relationship with the other minorities in Guangxi than with the Han nationality [33]. Height, fat mass, and fat distribution differ substantially between men and women, and these differences may, in part, explain the sex-specific susceptibilities to certain diseases such as CAD [34]. These considerable differences in anthropometry may reflect sex-specific differences in steroid hormone regulation, adipogenesis, lipid storage, muscle metabolism, composition, and contractile speed, skeletal growth and maturation, or lipolysis, and suggest a genetic underpinning [35]. Sexual dimorphism has been demonstrated as the potential of dyslipidemia and CVD risk factors. This study, therefore, was undertaken to detect the association of SPTY2D1 rs17579600 SNP and several environmental factors with serum lipid levels between males and females in the Mulao and Han populations.

Materials and methods

Subjects

The study populations including 712 unrelated subjects (248 males, 34.83% and 464 females, 65.17%) of Han and 689 unrelated participants (222 males, 33.22% and 467 females, 67.78%) of Mulao were randomly selected from our previous stratified randomized samples [36]. All participants were agricultural workers from Luocheng Mulao Autonomous County, Guangxi Zhuang Autonomous region, People's Republic of China. The participants' age ranged from 15 to 80 years with the mean age of 49.02 ± 14.39 years in Han and 48.43 ± 14.58 years in Mulao; respectively. The age distribution and gender ratio were matched between the two groups. All participants were essentially healthy with no history of CVD such as coronary artery disease, stroke, diabetes, hyper- or hypo-thyroids, and chronic renal disease. They were free from medications known to affect serum lipid levels. Informed consent was taken from all participants. The study design was approved by the Ethics Committee of the First Affiliated Hospital, Guangxi Medical University.

Epidemiological survey

The epidemiological survey was carried out using internationally standardized methods,

SPTY2D1 rs17579600 polymorphism and serum lipid traits

following a common protocol [37]. Information on demographics, socioeconomic status, and lifestyle factors was collected with standardized questionnaires. Alcohol consumption was categorized into groups of grams of alcohol per day: < 25 and \geq 25. Smoking status was categorized into groups of cigarettes per day: < 20 and \geq 20. Several parameters such as blood pressure, height, weight, waist circumference, and body mass index (BMI) were measured. The methods of measuring above parameters were referred to previous studies [38].

Biochemical measurements

A fasting venous blood sample of 5 ml was drawn from the participants. The levels of TC, TG, HDL-C and LDL-C in the samples were determined by enzymatic methods with commercially available kits. Serum apolipoprotein (Apo) A1 and ApoB levels were assessed by the immunoturbidimetric immunoassay [39].

Genotyping

Genomic DNA was isolated from peripheral blood leukocytes using the phenol-chloroform method [36-39]. The SPTY2D1 rs17579600 SNP was genotyped by polymerase chain reaction and restriction fragment length polymorphism (PCR-RFLP). PCR amplification was performed using 5'-CAAAGAAATCTCTATCTCAC-3' as the forward and 5'-ACCAGCCTGGCCAA-CATGGT-3' as reversed primer pair. Each amplification reaction was performed in a total volume of 25 μ l, 12.5 μ l of 2 \times Taq PCR MasterMix (constituent: 0.1 U Taq polymerase/ μ l, 500 μ M dNTP each and PCR buffer) and nuclease-free water 8.5 μ l, 20 pmol/L of each primer and 100 ng of genomic DNA, processing started with 7 min of pre-denaturing at 95°C and followed by 50 s of denaturing at 95°C, 45 s of annealing at 60°C and 1 min of elongation at 72°C for 33 cycles. The amplification was completed by a final extension at 72°C for 7 min. Then each restriction enzyme reaction was performed with 10 μ l of amplified DNA, 8 μ l of nuclease-free water, 1 μ l of 10 \times buffer solutions, and 10 U of 'Mob II' enzyme in a total volume of 20 μ l digested at 37°C overnight. After restriction enzyme digestion of the amplified DNA, the digestive products were separated by electrophoresis on 2% agarose gel. The length of each digested DNA fragment was determined by comparing migration of a sample with that of standard DNA marker. Genotypes were scored

by an experienced reader blinded to the epidemiological and lipid results. Six samples (each genotype in two; respectively) detected by the PCR-RFLP were also confirmed by direct sequencing. The PCR products were purified by low melting point gel electrophoresis and phenol extraction, and then the DNA sequences were analyzed using an ABI Prism 3100 (Applied Biosystems) in Shanghai Sangon Biological Engineering Technology & Services Co., Ltd., People's Republic of China.

Diagnostic criteria

The normal values of serum TC, TG, HDL-C, LDL-C, ApoA1 and ApoB levels, and the ratio of ApoA1 to ApoB in our Clinical Science Experiment Center were 3.10-5.17, 0.56-1.70, 1.16-1.42, 2.70-3.10 mmol/L, 1.20-1.60, 0.80-1.05 g/L, and 1.00-2.50; respectively [36-39].

Statistical analysis

The statistical analyses were performed with the statistical software package SPSS 17.0 (SPSS Inc., Chicago, Illinois). The quantitative variables were presented as mean \pm standard deviation (serum TG levels were presented as medians and interquartile ranges). Allele frequency was determined via direct counting, and the Hardy-Weinberg equilibrium was verified with the standard goodness-of-fit test. The genotype distribution between the two groups was analyzed by the chi-square test. General characteristics between two ethnic groups were compared by the Student's unpaired *t*-test. The association between genotypes and serum lipid parameters was tested by analysis of covariance (ANCOVA). Age, sex, BMI, smoking, and alcohol consumption were adjusted for the statistical analysis. Multivariable linear regression analyses with stepwise modeling were used to determine the correlation between genotypes (TT = 1, TC = 2, CC = 3) or alleles (the C allele non-carrier = 1, the C allele carrier = 2) and several environmental factors with serum lipid levels in males and females of Han and Mulao populations. Two sided *P* value < 0.05 was considered statistically significant.

Results

General and biochemical characteristics of the subjects

The comparison of general characteristics and serum lipid levels between the Han and Mulao

SPTY2D1 rs17579600 polymorphism and serum lipid traits

Table 1. Comparison of demographic, lifestyle characteristics and serum lipid levels between the Han and Mulao populations

| Parameter | Han | Mulao | T (χ^2) | P |
|--------------------------------------|--------------|--------------|----------------|-------|
| Number | 712 | 689 | | |
| Male/female | 248/464 | 222/467 | 1.071 | 0.301 |
| Age (years) | 49.02±14.39 | 48.43±14.58 | 0.762 | 0.446 |
| Height (cm) | 154.65±7.48 | 154.03±7.86 | 1.522 | 0.128 |
| Weight (kg) | 53.09±8.74 | 51.30±8.68 | 3.843 | 0.000 |
| Body mass index (kg/m ²) | 22.38±3.41 | 21.41±3.08 | 5.566 | 0.000 |
| waist circumference (cm) | 74.78±7.85 | 73.32±7.88 | 3.479 | 0.001 |
| Cigarette smoking (n%) | | | | |
| Nonsmoker | 547 (76.8) | 558 (81.0) | | |
| < 20 cigarettes/day | 137 (19.2) | 113 (16.4) | 4.211 | 0.122 |
| ≥ 20 cigarettes/day | 28 (3.9) | 18 (2.6) | | |
| Alcohol consumption [n (%)] | | | | |
| Nondrinker | 577 (81.0) | 577 (83.7) | | |
| < 25 g/day | 64 (9.0) | 38 (5.5) | 6.314 | 0.043 |
| ≥ 25 g/day | 71 (10.0) | 74 (10.7) | | |
| Systolic blood pressure (mmHg) | 127.03±18.35 | 116.86±11.79 | 12.380 | 0.000 |
| Diastolic blood pressure (mmHg) | 81.25±10.81 | 75.48±7.31 | 11.738 | 0.000 |
| Pulse pressure (mmHg) | 45.78±13.59 | 41.38±10.02 | 6.908 | 0.000 |
| Blood glucose (mmol/L) | 5.98±1.55 | 5.43±0.78 | 8.099 | 0.000 |
| Total cholesterol (mmol/L) | 4.96±1.09 | 4.95±1.27 | 0.045 | 0.964 |
| Triglyceride (mmol/L) | 1.02 (0.75) | 1.01 (0.76) | -1.852 | 0.064 |
| HDL-C (mmol/L) | 1.75±0.59 | 1.78±0.45 | -1.075 | 0.283 |
| LDL-C (mmol/L) | 2.90±0.89 | 2.85±0.85 | 1.230 | 0.219 |
| Apolipoprotein (Apo) A1 (g/L) | 1.34±0.26 | 1.34±0.38 | -0.080 | 0.937 |
| ApoB (g/L) | 0.95±0.53 | 0.85±0.20 | 4.409 | 0.000 |
| ApoA1/ApoB | 1.66±0.48 | 1.67±0.77 | -0.126 | 0.900 |

HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol. The value of triglyceride was presented as median (interquartile range). The difference between the two ethnic groups was determined by the Wilcoxon-Mann-Whitney test.

populations is summarized in **Table 1**. The levels of body weight, BMI, waist circumference, systolic blood pressure, diastolic blood pressure, pulse pressure, blood glucose and the levels of ApoB were lower in Mulao than in Han ($P < 0.05$ - 0.001), whereas the percentage of excessive alcohol consumption were higher in Mulao than in Han ($P < 0.05$ - 0.001). There were no significant differences in the gender ratio, age structure, body height, the percentage of cigarette smoking, serum TC, TG, HDL-C, LDL-C and ApoA1 levels and the ApoA1/ApoB ratio between the two ethnic groups ($P > 0.05$ for all).

Results of genotyping

After the genomic DNA of the samples was amplified by PCR, the purpose gene of 496-bp

nucleotide sequences could be seen in all samples (**Figure 1**). The genotypes identified were labeled according to the presence or absence of the enzyme restriction sites. Thus, TT genotype is homozygote for the absence of the site (496-bp), TC genotype is heterozygote for the presence and absence of the site (496-, 288- and 208-bp) and CC genotype is homozygote for the presence of the site (288- and 208-bp; **Figure 2**). The TT, CT and CC genotypes detected by PCR-RFLP were also confirmed by direct sequencing (**Figure 3**), respectively.

Genotypic and allelic frequencies

As shown in **Table 2**, the genotype and allele frequencies of SPTY2D1 rs17579600 SNP were different between the Han and Mulao populations (TT, 74.3% vs. 55.7%; TC, 17.6% vs.

SPTY2D1 rs17579600 polymorphism and serum lipid traits




Figure 1. Electrophoresis of PCR products of the samples. Lane M is the 100 bp Marker ladder; lines 1-6 are samples, the 496 bp bands are the target genes.




Figure 2. Genotyping of the *SPTY2D1* rs17579600 SNP. Lane M is the 100 bp Marker Ladder; lanes 1 and 2, TT genotype (496-bp); lanes 3 and 4, TC genotype (496-, 288- and 208-bp); and lanes 5 and 6, CC genotype (288- and 208-bp).

31.2%, CC, 8.1% vs. 13.1%, $P = 0.028$; T, 83.1% vs. 71.3%; C, 16.9% vs. 28.7%, $P = 0.044$). The genotype frequencies of rs17579600 SNP agreed with the Hardy-Weinberg equilibrium in both populations ($P > 0.05$ for each). Gender-subgroup analysis showed that the genotype and allele frequencies of *SPTY2D1* rs17579600 SNP between males and females were different in Han and Mulao. The genotype and allele frequencies were different between Han males and females (TT, 66.1% vs. 78.7%; TC, 27.8% vs. 12.1%, CC, 6.0% vs. 9.3%, $P = 0.017$; T, 80.0% vs. 84.7%; C, 20.0% vs. 15.3%, $P = 0.026$). The genotype and allele frequencies were significantly different between Mulao males and females (TT, 45.9% vs. 60.4%; TC, 31.5% vs. 31.0%, CC, 22.5% vs. 8.6%, $P =$

0.018; T, 61.7% vs. 75.9%; C, 38.3% vs. 24.1%, $P = 0.032$).

Genotypes and serum lipid levels

Tables 3 and **4** describe the association between genotypes and serum lipid levels. Serum ApoA1 levels in Han were different among the genotypes ($P < 0.05$), and the C allele carriers had higher ApoA1 levels than the C allele non-carriers. Serum TG levels in Mulao were different among the genotypes ($P < 0.05$), and the C allele carriers had lower TG levels than the C allele non-carriers. Subgroup analyses showed that serum levels of TC, LDL-C, ApoA1 and ApoB in Mulao males were different among the genotypes ($P < 0.05$ for all); the C allele carriers had higher serum ApoA1 level and lower serum TC, LDL-C and ApoB levels than the C allele non-carriers. In a word, the subjects with the minor C allele have more favorable lipid profiles than those with the C allele non-carriers.

Relative factors for serum lipid parameters

Several environmental factors such as age, gender, height, weight, waist circumference, alcohol consumption and cigarette smoking, and traditional cardiovascular risk factors such as BMI, fasting blood glucose and blood pressure levels were also correlated with serum lipid parameters in the Han and Mulao populations and in males and females of both ethnic groups ($P < 0.05-0.001$, **Tables 5** and **6**).

Discussion

In the current study, we showed that serum lipid profiles were significantly different between Han and Mulao ethnic groups. A significant difference in the genotype and allele frequencies of *SPTY2D1* rs17579600 SNP was also noted between the two ethnic populations. The minor C allele frequencies in Han and Mulao were 16.9% and 28.7% respectively, which were in close proximity to those of Chinese Han Beijing (14.0%) reported in international haplotype map (HapMap) project. According to HapMap data, the minor allele frequency of rs17579600 was 15.1% in Japanese, and 8.8% in Europeans. Apparently, the minor allele frequency was higher in Asian than the Western populations. These findings suggest that genotype and allele frequencies of

SPTY2D1 rs17579600 polymorphism and serum lipid traits




Figure 3. A part of the nucleotide forward sequence of the *SPTY2D1* rs17579600 SNP. A: TT genotype; B: TC genotype; C: CC genotype.

Table 2. Comparison of the genotype and allele frequencies of the *SPTY2D1* rs17579600 SNP between males and females of the Han and Mulao populations

| Group | n | Genotype | | | Allele | |
|----------|-----|------------|------------|-----------|-------------|------------|
| | | TT | TC | CC | T | C |
| Han | 712 | 529 (74.3) | 125 (17.6) | 58 (8.1) | 1183 (83.1) | 241 (16.9) |
| Mulao | 689 | 384 (55.7) | 215 (31.2) | 90 (13.1) | 983 (71.3) | 395 (28.7) |
| χ^2 | - | | 7.132 | | 4.065 | |
| P | - | | 0.028 | | 0.044 | |
| Han | | | | | | |
| Male | 248 | 164 (66.1) | 69 (27.8) | 15 (6.0) | 397 (80.0) | 99 (20.0) |
| Female | 464 | 365 (78.7) | 56 (12.1) | 43 (9.3) | 786 (84.7) | 142 (15.3) |
| χ^2 | - | | 8.166 | | 4.988 | |
| P | - | | 0.017 | | 0.026 | |
| Mulao | | | | | | |
| Male | 222 | 102 (45.9) | 70 (31.5) | 50 (22.5) | 274 (61.7) | 170 (38.3) |
| Female | 467 | 282 (60.4) | 145 (31.0) | 40 (8.6) | 709 (75.9) | 225 (24.1) |
| χ^2 | - | | 7.985 | | 4.582 | |
| P | - | | 0.018 | | 0.032 | |

SPTY2D1 rs17579600 SNP are inconsistent among diverse ethnic groups.

to the differences in genetic backgrounds, dietary habits, and environmental factors

A recent GWAS reported that *SPTY2D1* rs10128711 SNP was the top association SNP with MetS in European ancestry [27, 28]. The minor allele was significantly associated with TC. Another our previous study shown that the *SPTY2D1* rs7934205 SNP minor allele was significantly associated with pleiotropic (one SNP influence many serum lipid traits) effects on serum lipid profiles. In the present study, the *SPTY2D1* rs17579600 SNP was correlated serum ApoA1 levels in Han and TG in Mulao. However, no association with TC was detected either in Han or Mulao population. The reason for this discrepancy is not fully understood. It might be due

SPTY2D1 rs17579600 polymorphism and serum lipid traits

Table 3. Comparison of the genotypes and serum lipid levels in the Han and Mulao populations

| Ethnic/ Genotype | N | TC (mmol/L) | TG (mmol/L) | HDL-C (mmol/L) | LDL-C (mmol/L) | ApoA1 (g/L) | ApoB (g/L) | ApoA1/ ApoB |
|---------------------|-----|----------------|----------------|-------------------|-------------------|----------------|---------------|----------------|
| Han | | | | | | | | |
| TT | 529 | 4.98±1.13 | 1.02 (0.74) | 1.73±0.64 | 2.86±0.89 | 1.32±0.29 | 0.86±0.21 | 1.64±0.40 |
| TC/CC | 183 | 4.90±0.97 | 1.02 (0.75) | 1.76±0.57 | 2.81±0.75 | 1.35±0.25 | 0.83±0.17 | 1.67±0.51 |
| <i>F</i> | | 0.690 | 0.091 | 0.960 | 0.779 | 4.347 | 3.229 | 1.900 |
| <i>P</i> | | 0.502 | 0.763 | 0.328 | 0.378 | 0.037 | 0.073 | 0.169 |
| Mulao | | | | | | | | |
| TT | 384 | 4.98±1.38 | 1.14 (0.80) | 1.78±0.47 | 2.94±0.93 | 1.34±0.41 | 0.96±0.55 | 1.65±0.74 |
| TC/CC | 305 | 4.92±1.13 | 1.07 (0.76) | 1.79±0.41 | 2.88±0.86 | 1.35±0.35 | 0.93±0.50 | 1.68±0.80 |
| <i>F</i> | | 2.292 | 4.172 | 0.041 | 0.017 | 0.000 | 0.671 | 1.278 |
| <i>P</i> | | 0.131 | 0.041 | 0.839 | 0.896 | 0.996 | 0.413 | 0.259 |

TC, total cholesterol; TG, triglyceride; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; ApoA1, apolipoprotein A1; ApoB, apolipoprotein B; ApoA1/ApoB, the ratio of apolipoprotein A1 to apolipoprotein B. The value of TG was presented as median (interquartile range). The difference between the genotypes was determined by the Wilcoxon-Mann-Whitney test.

Table 4. Comparison between the SPTY2D1 rs17579600 genotypes and serum levels in the males and females of the Mulao and Han populations

| Genotype | n | TC (mmol/L) | TG (mmol/L) | HDL-C (mmol/L) | LDL-C (mmol/L) | ApoA1 (g/L) | ApoB (g/L) | ApoA/ ApoB |
|--------------|-----|----------------|----------------|-------------------|-------------------|----------------|---------------|---------------|
| Han/Male | | | | | | | | |
| TT | 102 | 5.04±1.93 | 1.02 (0.76) | 1.70±0.63 | 2.94±0.74 | 1.31±0.47 | 0.98±0.62 | 1.58±0.77 |
| TC/CC | 120 | 5.00±0.90 | 0.97 (0.74) | 1.78±0.40 | 2.79±0.90 | 1.37±0.33 | 0.96±0.48 | 1.62±0.63 |
| <i>F</i> | | 0.322 | 1.754 | 2.330 | 1.618 | 0.124 | 0.055 | 0.099 |
| <i>P</i> | | 0.571 | 0.187 | 0.128 | 0.205 | 0.725 | 0.814 | 0.754 |
| Han/Female | | | | | | | | |
| TT | 282 | 4.96±1.12 | 1.02 (0.78) | 1.79±0.42 | 2.93±1.03 | 1.34±0.36 | 0.95±0.52 | 1.68±0.73 |
| TC/CC | 185 | 4.86±1.25 | 1.01 (0.73) | 1.81±0.40 | 2.91±0.85 | 1.35±0.39 | 0.91±0.52 | 1.73±0.89 |
| <i>F</i> | | 0.372 | 2.552 | 0.312 | 0.245 | 0.065 | 0.243 | 0.294 |
| <i>P</i> | | 0.542 | 0.111 | 0.576 | 0.621 | 0.799 | 0.622 | 0.588 |
| Mulao/Male | | | | | | | | |
| TT | 164 | 5.40±1.17 | 1.31 (0.93) | 1.65±0.42 | 3.02±0.88 | 1.33±0.30 | 0.96±0.22 | 1.54±0.52 |
| TC/CC | 84 | 4.88±0.91 | 1.10 (0.83) | 1.70±0.47 | 2.74±0.79 | 1.40±0.30 | 0.85±0.16 | 1.60±0.40 |
| <i>F</i> | | 9.637 | 0.615 | 3.417 | 8.600 | 4.881 | 6.341 | 0.027 |
| <i>P</i> | | 0.002 | 0.434 | 0.066 | 0.004 | 0.028 | 0.012 | 0.870 |
| Mulao/Female | | | | | | | | |
| TT | 365 | 4.92±1.02 | 0.96 (0.69) | 1.79±0.61 | 2.87±0.71 | 1.32±0.28 | 0.81±0.19 | 1.67±0.39 |
| TC/CC | 99 | 4.79±1.06 | 0.97 (0.75) | 1.80±0.78 | 2.79±0.88 | 1.33±0.22 | 0.81±0.18 | 1.73±0.49 |
| <i>F</i> | | 0.116 | 0.691 | 0.614 | 0.237 | 2.375 | 0.331 | 1.889 |
| <i>P</i> | | 0.733 | 0.406 | 0.434 | 0.627 | 0.124 | 0.565 | 0.170 |

TC, total cholesterol; TG, triglyceride; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; ApoA1, apolipoprotein A1; ApoB, apolipoprotein B; ApoA1/ApoB, the ratio of apolipoprotein A1 to apolipoprotein B. The values of triglyceride were presented as median (interquartile range), and the difference between the TT and TC/CC genotypes was determined by the Wilcoxon-Mann-Whitney test.

between the two ethnic populations and/or simply due to the low power of this study. It is well accepted that ethnic differences in serum

lipid levels were partly due to the differences in the dietary intakes [40]. Diet alone could account for up to 2.5% of the variability on

SPTY2D1 rs17579600 polymorphism and serum lipid traits

Table 5. The risk factors for serum lipid parameters in the Han and Mulao populations

| Lipid parameter | Risk factor | <i>B</i> | <i>Std. error</i> | <i>Beta</i> | <i>t</i> | <i>P</i> |
|-----------------|--------------------------|----------|-------------------|-------------|----------|----------|
| Han and Mulao | | | | | | |
| TC | Ethnic group | 0.137 | 0.067 | 0.058 | 2.037 | 0.042 |
| | Age | 0.011 | 0.002 | -.136 | 4.554 | 0.000 |
| | Alcohol consumption | 0.219 | 0.060 | 0.118 | 3.665 | 0.000 |
| | Waist circumference | 0.027 | 0.006 | 0.180 | 4.199 | 0.000 |
| | Diastolic blood pressure | 0.008 | 0.003 | 0.065 | 2.293 | 0.022 |
| TG | Alcohol consumption | 0.301 | 0.100 | 0.097 | 3.021 | 0.003 |
| | Waist circumference | 0.071 | 0.011 | 0.281 | 6.605 | 0.000 |
| | Waist circumference | 0.070 | 0.009 | 0.297 | 7.779 | 0.000 |
| | Diastolic blood pressure | 0.013 | 0.006 | 0.064 | 2.300 | 0.022 |
| HDL-C | Glucose | 0.091 | 0.043 | 0.058 | 2.121 | 0.034 |
| | Gender | 0.150 | 0.047 | 0.135 | 3.171 | 0.002 |
| | Age | 0.004 | 0.001 | 0.108 | 3.589 | 0.000 |
| | Alcohol consumption | 0.124 | 0.027 | 0.151 | 4.652 | 0.000 |
| | Height | 0.018 | 0.009 | 0.257 | 2.044 | 0.041 |
| | Weight | -0.028 | 0.012 | -0.461 | -2.377 | 0.018 |
| LDL-C | Waist circumference | -0.007 | 0.003 | -0.103 | -2.378 | 0.018 |
| | Ethnic group | 0.132 | 0.050 | 0.075 | 2.646 | 0.008 |
| | Gender | -0.210 | 0.078 | -0.114 | -2.686 | 0.007 |
| | Age | 0.009 | 0.002 | 0.145 | 4.852 | 0.000 |
| | Cigarette smoking | -0.141 | 0.061 | -0.081 | -2.314 | 0.021 |
| ApoA1 | Waist circumference | 0.018 | 0.005 | 0.159 | 3.695 | 0.000 |
| | Gender | 0.112 | 0.029 | 0.162 | 3.821 | 0.000 |
| | Age | 0.002 | 0.001 | 0.106 | 3.527 | 0.000 |
| ApoB | Alcohol consumption | 0.142 | 0.017 | 0.279 | 8.557 | 0.000 |
| | Ethnic group | 0.152 | 0.023 | 0.190 | 6.718 | 0.000 |
| ApoA1/ApoB | Waist circumference | 0.010 | 0.002 | 0.202 | 4.727 | 0.000 |
| | Pulse pressure | 0.002 | 0.001 | 0.075 | 2.727 | 0.006 |
| | Glucose | 0.021 | 0.009 | 0.065 | 2.373 | 0.018 |
| | Gender | 0.213 | 0.057 | 0.158 | 3.771 | 0.000 |
| ApoA1/ApoB | Alcohol consumption | 0.127 | 0.032 | 0.128 | 3.980 | 0.000 |
| | Waist circumference | -0.015 | 0.003 | -0.182 | -4.267 | 0.000 |
| | Pulse pressure | -0.003 | 0.001 | -0.061 | -2.211 | 0.027 |
| Han | | | | | | |
| TC | Gender | -0.394 | 0.131 | -0.173 | -3.018 | 0.003 |
| | Age | 0.010 | 0.003 | 0.127 | 2.978 | 0.003 |
| | Alcohol consumption | 0.294 | 0.076 | 0.172 | 3.887 | 0.000 |
| | Waist circumference | 0.041 | 0.009 | 0.295 | 4.708 | 0.000 |
| | Diastolic blood pressure | 0.013 | 0.004 | 0.134 | 3.583 | 0.000 |
| TG | Age | -0.015 | 0.007 | -0.098 | -2.278 | 0.023 |
| | Cigarette smoking | 0.762 | 0.201 | 0.182 | 3.793 | 0.000 |
| | Weight | 0.092 | 0.040 | 0.916 | 2.303 | 0.022 |
| | Waist circumference | 0.102 | 0.018 | 0.365 | 5.758 | 0.000 |
| HDL-C | Diastolic blood pressure | 0.023 | 0.008 | 0.113 | 2.997 | 0.003 |
| | Glucose | 0.177 | 0.053 | 0.125 | 3.326 | 0.001 |
| | Gender | 0.164 | 0.075 | 0.132 | 2.177 | 0.030 |
| | Age | 0.005 | 0.002 | 0.125 | 2.745 | 0.006 |
| | | | | | | |

SPTY2D1 rs17579600 polymorphism and serum lipid traits

| | | | | | | |
|------------|--------------------------|--------|-------|--------|--------|-------|
| LDL-C | Alcohol consumption | 0.131 | 0.044 | 0.141 | 3.011 | 0.003 |
| | Gender | -0.400 | 0.106 | -0.223 | -3.789 | 0.000 |
| | Age | 0.011 | 0.003 | 0.180 | 4.095 | 0.000 |
| | Cigarette smoking | -0.342 | 0.079 | -0.211 | -4.310 | 0.000 |
| ApoA1 | Waist circumference | 0.027 | 0.007 | 0.251 | 3.890 | 0.000 |
| | Genotype | -0.044 | 0.021 | -0.073 | -2.085 | 0.037 |
| | Age | 0.002 | 0.001 | 0.085 | 1.988 | 0.047 |
| | Cigarette smoking | 0.063 | 0.023 | 0.128 | 2.713 | 0.007 |
| ApoB | Alcohol consumption | 0.146 | 0.018 | 0.359 | 8.138 | 0.000 |
| | Weight | -0.021 | 0.007 | -0.699 | -3.039 | 0.002 |
| | Gender | -0.087 | 0.023 | -0.203 | -3.802 | 0.000 |
| | Age | 0.001 | 0.001 | 0.084 | 2.107 | 0.035 |
| | Alcohol consumption | 0.036 | 0.013 | 0.111 | 2.689 | 0.007 |
| | Waist circumference | 0.009 | 0.002 | 0.357 | 6.094 | 0.000 |
| ApoA1/ApoB | Diastolic blood pressure | 0.002 | 0.001 | 0.099 | 2.823 | 0.005 |
| | Glucose | 0.015 | 0.005 | 0.111 | 3.196 | 0.001 |
| | Gender | 0.241 | 0.056 | 0.238 | 4.301 | 0.000 |
| | Cigarette smoking | 0.134 | 0.042 | 0.147 | 3.189 | 0.001 |
| | Alcohol consumption | 0.113 | 0.032 | 0.149 | 3.491 | 0.001 |
| | Waist circumference | -0.014 | 0.004 | -0.220 | -3.625 | 0.000 |
| Mulao | | | | | | |
| TC | Age | 0.014 | 0.004 | 0.163 | 3.879 | 0.000 |
| | Height | -0.112 | 0.047 | -0.655 | -2.376 | 0.018 |
| | Weight | 0.174 | 0.071 | 1.186 | 2.455 | 0.014 |
| | Body mass index | -0.383 | 0.168 | -0.927 | -2.278 | 0.023 |
| TG | Genotype | -0.272 | 0.133 | -0.078 | -2.042 | 0.041 |
| | Cigarette smoking | -0.363 | 0.182 | -0.099 | -1.996 | 0.046 |
| | Alcohol consumption | 0.378 | 0.123 | 0.140 | 3.065 | 0.002 |
| | Height | -0.158 | 0.063 | -0.679 | -2.499 | 0.013 |
| | Weight | 0.252 | 0.095 | 1.260 | 2.643 | 0.008 |
| | Body mass index | -0.582 | 0.226 | -1.032 | -2.571 | 0.010 |
| | Waist circumference | 0.040 | 0.013 | 0.182 | 3.182 | 0.002 |
| | Glucose | -0.172 | 0.083 | -0.077 | -2.077 | 0.038 |
| HDL-C | Gender | 0.128 | 0.058 | 0.135 | 2.223 | 0.027 |
| | Age | 0.003 | 0.001 | 0.100 | 2.412 | 0.016 |
| | Alcohol consumption | 0.116 | 0.032 | 0.167 | 3.671 | 0.000 |
| | Waist circumference | -0.007 | 0.003 | -0.128 | -2.245 | 0.025 |
| LDL-C | Age | 0.008 | 0.003 | 0.131 | 3.122 | 0.002 |
| ApoA1 | Gender | 0.156 | 0.051 | 0.190 | 3.056 | 0.002 |
| | Age | 0.003 | 0.001 | 0.117 | 2.762 | 0.006 |
| ApoB | Alcohol consumption | 0.136 | 0.028 | 0.228 | 4.899 | 0.000 |
| | Waist circumference | 0.011 | 0.004 | 0.166 | 2.866 | 0.004 |
| | Pulse pressure | 0.005 | 0.002 | 0.104 | 2.750 | 0.006 |
| ApoA1/ApoB | Alcohol consumption | 0.140 | 0.055 | 0.117 | 2.537 | 0.011 |
| | Waist circumference | -0.015 | 0.006 | -0.159 | -2.748 | 0.006 |
| | Pulse pressure | -0.006 | 0.003 | -0.079 | -2.099 | 0.036 |

TC, total cholesterol; TG, triglyceride; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; ApoA1, apolipoprotein A1; ApoB, apolipoprotein B; ApoA1/ApoB, the ratio of apolipoprotein A1 to apolipoprotein B.

SPTY2D1 rs17579600 polymorphism and serum lipid traits

Table 6. The risk factors for serum lipid parameters in the males and females of the Han and Mulao populations

| Lipid parameter | Risk factor | B | Std. error | Beta | t | P |
|-----------------|--------------------------|--------|------------|--------|--------|-------|
| Han/male | | | | | | |
| TC | Genotype | -0.268 | 0.101 | -0.116 | -2.651 | 0.008 |
| | Alcohol consumption | 0.222 | 0.074 | 0.140 | 2.998 | 0.003 |
| | Waist circumference | 0.050 | 0.012 | 0.370 | 4.255 | 0.000 |
| | Diastolic blood pressure | 0.032 | 0.005 | 0.311 | 6.842 | 0.000 |
| | Glucose | 0.102 | 0.031 | 0.150 | 3.262 | 0.001 |
| TG | Cigarette smoking | 0.862 | 0.236 | 0.172 | 3.650 | 0.000 |
| | Waist circumference | 0.227 | 0.036 | 0.562 | 6.344 | 0.000 |
| | Diastolic blood pressure | 0.035 | 0.014 | 0.116 | 2.503 | 0.013 |
| | Glucose | 0.300 | 0.095 | 0.148 | 3.174 | 0.002 |
| HDL-C | Age | 0.004 | 0.002 | 0.115 | 2.264 | 0.024 |
| | Cigarette smoking | 0.081 | 0.031 | 0.120 | 2.646 | 0.008 |
| | Alcohol consumption | 0.113 | 0.029 | 0.179 | 3.919 | 0.000 |
| | Height | 0.077 | 0.023 | 1.027 | 3.379 | 0.001 |
| | Weight | -0.105 | 0.030 | -2.163 | -3.472 | 0.001 |
| | Body mass index | 0.226 | 0.080 | 1.520 | 2.835 | 0.005 |
| | Diastolic blood pressure | 0.006 | 0.002 | 0.148 | 3.341 | 0.001 |
| | Pulse pressure | -0.003 | 0.001 | -0.105 | -2.259 | 0.024 |
| LDL-C | Cigarette smoking | -0.318 | 0.062 | -0.250 | -5.164 | 0.000 |
| | Waist circumference | 0.007 | 0.004 | 0.094 | 1.974 | 0.049 |
| ApoA1 | Cigarette smoking | 0.075 | 0.019 | 0.190 | 3.930 | 0.000 |
| | Alcohol consumption | 0.047 | 0.018 | 0.128 | 2.629 | 0.009 |
| | Height | 0.040 | 0.014 | 0.928 | 2.851 | 0.005 |
| | Weight | -0.057 | 0.019 | -2.026 | -3.035 | 0.003 |
| | Body mass index | 0.126 | 0.050 | 1.456 | 2.534 | 0.012 |
| ApoB | Genotype | -0.049 | 0.018 | -0.115 | 2.729 | 0.007 |
| | Alcohol consumption | 0.034 | 0.013 | 0.116 | 2.580 | 0.010 |
| | Waist circumference | 0.009 | 0.002 | 0.355 | 4.234 | 0.000 |
| | Diastolic blood pressure | 0.004 | 0.001 | 0.230 | 5.262 | 0.000 |
| | Glucose | 0.028 | 0.006 | 0.222 | 5.027 | 0.000 |
| ApoA1/ApoB | Cigarette smoking | 0.111 | 0.031 | 0.164 | 3.545 | 0.000 |
| Han/female | | | | | | |
| TC | Age | 0.019 | 0.005 | 0.246 | 4.139 | 0.000 |
| TG | Waist circumference | 0.047 | 0.014 | 0.266 | 3.288 | 0.001 |
| | Diastolic blood pressure | 0.020 | 0.006 | 0.168 | 3.293 | 0.001 |
| HDL-C | Body mass index | -0.034 | 0.007 | -0.233 | -5.149 | 0.000 |
| | Genotype | 0.080 | 0.028 | 0.128 | 2.831 | 0.005 |
| LDL-C | Body mass index | 0.059 | 0.013 | 0.203 | 4.495 | 0.000 |
| | Age | 0.020 | 0.004 | 0.314 | 5.284 | 0.000 |
| ApoA1 | Height | -0.029 | 0.013 | -0.642 | -2.120 | 0.035 |
| | Body mass index | -0.106 | 0.049 | -1.296 | -2.168 | 0.031 |
| ApoB | Genotype | 0.034 | 0.017 | 0.088 | 2.029 | 0.043 |
| | Age | 0.002 | 0.001 | 0.130 | 2.262 | 0.024 |
| | Waist circumference | 0.005 | 0.002 | 0.200 | 2.631 | 0.009 |
| | Glucose | 0.015 | 0.007 | 0.099 | 2.153 | 0.032 |
| ApoA1/ApoB | Age | -0.011 | 0.004 | -0.133 | -2.886 | 0.004 |
| | Waist circumference | -0.024 | 0.010 | -0.156 | -2.386 | 0.017 |

SPTY2D1 rs17579600 polymorphism and serum lipid traits

| | | | | | | |
|--------------|--------------------------|--------|-------|--------|--------|-------|
| Mulao/male | | | | | | |
| TC | Age | 0.014 | 0.004 | 0.160 | 3.348 | 0.001 |
| | Cigarette smoking | 0.174 | 0.075 | 0.106 | 2.330 | 0.020 |
| | Glucose | -0.146 | 0.064 | -0.105 | -2.280 | 0.023 |
| TG | Waist circumference | 0.019 | 0.008 | 0.150 | 2.337 | 0.020 |
| HDL-C | Age | 0.004 | 0.002 | 0.095 | 2.044 | 0.042 |
| | Alcohol consumption | 0.144 | 0.030 | 0.215 | 4.807 | 0.000 |
| | Body mass index | -0.216 | 0.108 | -1.187 | -1.992 | 0.047 |
| | Glucose | -0.069 | 0.029 | -0.105 | -2.335 | 0.020 |
| LDL-C | Age | 0.008 | 0.003 | 0.118 | 2.440 | 0.015 |
| | Glucose | -0.100 | 0.049 | -0.095 | -2.039 | 0.042 |
| ApoA1 | Gender | -0.083 | 0.038 | -0.098 | -2.182 | 0.030 |
| | Age | 0.006 | 0.002 | 0.164 | 3.519 | 0.000 |
| | Alcohol consumption | 0.145 | 0.025 | 0.256 | 5.731 | 0.000 |
| | Diastolic blood pressure | 0.006 | 0.003 | 0.110 | 2.342 | 0.020 |
| ApoB | Cigarette smoking | 0.091 | 0.039 | 0.105 | 2.340 | 0.020 |
| | Pulse pressure | 0.011 | 0.003 | 0.180 | 3.960 | 0.000 |
| ApoA1/ApoB | Cigarette smoking | -0.158 | 0.050 | -0.141 | -3.166 | 0.002 |
| | Alcohol consumption | 0.145 | 0.044 | 0.148 | 3.316 | 0.001 |
| | Body mass index | -0.339 | 0.157 | -1.282 | -2.151 | 0.032 |
| | Diastolic blood pressure | 0.011 | 0.005 | 0.113 | 2.400 | 0.017 |
| Mulao/female | | | | | | |
| TC | Genotype | -0.283 | 0.104 | -0.123 | -2.725 | 0.007 |
| | Age | 0.020 | 0.004 | 0.231 | 4.660 | 0.000 |
| | Pulse pressure | 0.013 | 0.005 | 0.119 | 2.562 | 0.011 |
| TG | Gender | -0.164 | 0.057 | -0.129 | -2.884 | 0.004 |
| | Waist circumference | 0.015 | 0.006 | 0.179 | 2.319 | 0.021 |
| HDL-C | Waist circumference | -0.005 | 0.004 | -0.053 | -1.136 | 0.025 |
| | Genotype | 0.167 | 0.043 | 0.182 | 3.874 | 0.000 |
| LDL-C | Genotype | -0.200 | 0.070 | -0.124 | -2.855 | 0.005 |
| | Age | 0.018 | 0.003 | 0.290 | 6.080 | 0.000 |
| | Pulse pressure | 0.007 | 0.003 | 0.089 | 2.001 | 0.046 |
| ApoA1 | Genotype | -0.072 | 0.034 | -0.098 | -2.100 | 0.036 |
| | Age | 0.003 | 0.001 | 0.101 | 1.981 | 0.048 |
| ApoB | Genotype | -0.091 | 0.045 | -0.093 | -2.043 | 0.042 |
| | Waist circumference | 0.012 | 0.005 | 0.185 | 2.343 | 0.020 |
| | Glucose | 0.072 | 0.029 | 0.115 | 2.470 | 0.014 |
| ApoA1/ApoB | Body mass index | -0.036 | 0.008 | -0.206 | -4.460 | 0.000 |
| | Cigarette smoking | 0.345 | 0.112 | 0.144 | 3.093 | 0.002 |
| | Systolic blood pressure | -0.002 | 0.001 | -0.070 | -1.373 | 0.017 |
| | Age | -0.008 | 0.002 | -0.197 | -3.765 | 0.000 |
| | Genotype | 0.057 | 0.032 | -0.080 | 1.754 | 0.008 |

TC, total cholesterol; TG, triglyceride; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; ApoA1, apolipoprotein A1; ApoB, apolipoprotein B; ApoA1/ApoB, the ratio of apolipoprotein A1 to apolipoprotein B.

serum lipid levels [41-45]. Although rice and corn are the staple foods for both ethnic groups; Mulao peoples have a typical habit of eating cold foods along with acidic and spicy dishes,

local bean soy sauce, pickled vegetables and animal offal's which contain abundant saturated fatty acid. Therefore, it is possible that the difference in dietary habit between Han and

SPTY2D1 rs17579600 polymorphism and serum lipid traits

Mulao ethnic groups partly contribute variability in the effect of *SPTY2D1* rs17579600 SNP on serum lipid levels.

It has been well-known that the males had higher serum levels of bad cholesterols and lower levels of good cholesterols than the females, especially in women before menopause [46]. On gender subgroup analysis, the genotype frequencies between males and females were different both in Han and Mulao. The minor allele frequency was higher in males than females. Here, we found that the minor C allele of *SPTY2D1* rs17579600 SNP had higher serum ApoA1 level and lower serum TC, LDL-C and ApoB levels than the C allele non-carriers just in Mulao males. In other words, the subjects with the minor C allele have more favorable lipid profiles than those with the C allele non-carriers but the minor C allele frequency was higher in males than females. The reason for this discrepancy mainly attributed to the role of gonadal steroid hormones, estrogen especially [47-49]. To the best of our knowledge, this study is the first attempt to report the gender specific association of *SPTY2D1* rs17579600 SNP. Therefore, further studies with larger sample size are still needed to confirm this association.

Several environmental factors were also correlated with serum lipid levels in males and females of both Han and Mulao populations. In the present study, The Han has significantly higher levels of body weight, BMI, waist circumference, systolic blood pressure, diastolic blood pressure, pulse pressure, blood glucose and the levels of ApoB and lower the percentage of excessive alcohol consumption compared to the Han counterparts. Garcia-Palmieri et al. stated that diet and relative weight could account for up to 6% of the variability in serum cholesterol levels [41]. In particular, for every 1-kg decrease in body weight, TG decreased by 0.011 mmol/L and HDL-C increased by 0.011 mmol/L [50]. Rimm et al. documented that consuming of 30 g of ethanol per day increased the concentrations of HDL-C by 3.99 mg/dl, ApoA1 by 8.82 mg/dl, and TG by 5.69 mg/dl [51]. Yin et al. also showed that BMI and alcohol consumption could interact with certain lipid-related gene variants to modify the serum lipid levels in Bai Ku Yao and Han Chinese ethnic groups [52, 53]. Therefore, the results of exposure to different environmental factors may fur-

ther modify the effect of genetic variation on serum lipid levels in our study populations.

This study has some limitations. The sample size was relatively small compared to many GWAS and replication studies. Hence, further studies with larger sample sizes are needed to confirm our results. Secondly, we were not able to alleviate the effect of diet and several environmental factors during the statistical analysis. Thirdly, although we have detected the effects of *SPTY2D1* rs17579600 SNP on serum lipid levels in this study, there are still many lipid-related SNPs and the interactions of SNP-SNP and/or SNP-environmental factors. What's more, the relevance of this finding has to be defined in further high caliber of studies including incorporating the genetic information of *SPTY2D1* rs17579600 SNP and in vitro functional studies to confirm the impact of a variant on a molecular level.

Conclusion

In conclusion, the minor C allele frequency of the *SPTY2D1* rs17579600 SNP is higher in Mulao than in Han but lower in females than in males. The minor C allele carriers in both ethnic groups and gender subgroups have more favorable serum lipid profiles than the C allele non-carriers. These findings suggest that the association between the *SPTY2D1* rs17579600 SNP and serum lipid levels might have ethnic- and/or sex-specificity.

Acknowledgements

This study was supported by the National Natural Science Foundation of China (No: 30960130), and the Innovation Project of Guangxi Graduate Education.

Disclosure of conflict of interest

None.

Address correspondence to: Rui-Xing Yin, Department of Cardiology, Institute of Cardiovascular Diseases, The First Affiliated Hospital, Guangxi Medical University, Nanning 530021, Guangxi, China. E-mail: yinruixing@163.com

References

- [1] Yazdanyar A and Newman AB. The burden of cardiovascular disease in the elderly: morbidi-

SPTY2D1 rs17579600 polymorphism and serum lipid traits

- ty, mortality, and costs. *Clin Geriatr Med* 2009; 25: 563-577.
- [2] Pandya A, Gaziano TA, Weinstein MC and Cutler D. More Americans living longer with cardiovascular disease will increase costs while lowering quality of life. *Health Aff (Millwood)* 2013; 32: 1706-1714.
- [3] Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, de Ferranti S, Després JP, Fullerton HJ, Howard VJ, Huffman MD, Judd SE, Kissela BM, Lackland DT, Lichtman JH, Lisabeth LD, Liu S, Mackey RH, Matchar DB, McGuire DK, Mohler ER 3rd, Moy CS, Muntner P, Mussolino ME, Nasir K, Neumar RW, Nichol G, Palaniappan L, Pandey DK, Reeves MJ, Rodriguez CJ, Sorlie PD, Stein J, Towfighi A, Turan TN, Virani SS, Willey JZ, Woo D, Yeh RW, Turner MB; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2015 update: a report from the American Heart Association. *Circulation* 2015; 131: e29-322.
- [4] Matthan NR, Zhu L, Pencina M, D'Agostino RB, Schaefer EJ and Lichtenstein AH. Sex-specific differences in the predictive value of cholesterol homeostasis markers and 10-year cardiovascular disease event rate in Framingham Offspring Study participants. *J Am Heart Assoc* 2013; 2: e005066.
- [5] Lindman AS, Veierød MB, Tverdal A, Pedersen JI and Selmer R. Nonfasting triglycerides and risk of cardiovascular death in men and women from the Norwegian Counties Study. *Eur J Epidemiol* 2010; 25: 789-798.
- [6] Orekhov AN, Bobryshev YV, Sobenin IA, Melnichenko AA and Chistiakov DA. Modified low density lipoprotein and lipoprotein-containing circulating immune complexes as diagnostic and prognostic biomarkers of atherosclerosis and type 1 diabetes macrovascular disease. *Int J Mol Sci* 2014; 15: 12807-12841.
- [7] Furtado JD, Campos H, Appel LJ, Miller ER, Laranjo N, Carey VJ and Sacks FM. Effect of protein, unsaturated fat, and carbohydrate intakes on plasma apolipoprotein B and VLDL and LDL containing apolipoprotein C-III: results from the OmniHeart Trial. *Am J Clin Nutr* 2008; 87: 1623-1630.
- [8] Chapman MJ, Ginsberg HN, Amarenco P, Andreotti F, Borén J, Catapano AL, Descamps OS, Fisher E, Kovanen PT, Kuivenhoven JA, Lesnik P, Masana L, Nordestgaard BG, Ray KK, Reiner Z, Taskinen MR, Tokgözoğlu L, Tybjaerg-Hansen A, Watts GF; European Atherosclerosis Society Consensus Panel. Triglyceride-rich lipoproteins and high-density lipoprotein cholesterol in patients at high risk of cardiovascular disease: evidence and guidance for management. *Eur Heart J* 2011; 32: 1345-1361.
- [9] Hamon SC, Kardina SL, Boerwinkle E, Liu K, Klos KL, Clark AG and Sing CF. Evidence for consistent intragenic and intergenic interactions between SNP effects in the APOA1/C3/A4/A5 gene cluster. *Hum Hered* 2006; 61: 87-96.
- [10] Tamang HK, Timilsina U, Singh KP, Shrestha S, Raman RK, Panta P, Karna P, Khadka L and Dahal C. Apo B/Apo A-I Ratio is Statistically A Better Predictor of Cardiovascular Disease (CVD) than Conventional Lipid Profile: A Study from Kathmandu Valley, Nepal. *J Clin Diagn Res* 2014; 8: 34-36.
- [11] Scott R, Donoghoe M, Watts GF, O'Brien R, Pardy C, Taskinen MR, Davis TM, Colman PG, Manning P, Fulcher G, Keech AC; FIELD Study Investigators. Impact of metabolic syndrome and its components on cardiovascular disease event rates in 4900 patients with type 2 diabetes assigned to placebo in the FIELD randomised trial. *Cardiovasc Diabetol* 2011; 10: 102.
- [12] Vishram JK, Borglykke A, Andreasen AH, Jepsen J, Ibsen H, Jørgensen T, Palmieri L, Giampaoli S, Donfrancesco C, Kee F, Mancia G, Cesana G, Kuulasmaa K, Salomaa V, Sans S, Ferrieres J, Dallongeville J, Söderberg S, Arveiler D, Wagner A, Tunstall-Pedoe H, Drygas W, Olsen MH; MORGAM Project. Correction: Impact of Age and Gender on the Prevalence and Prognostic Importance of the Metabolic Syndrome and Its Components in Europeans. The MORGAM Prospective Cohort Project. *PLoS One* 2015; 10: e0128848.
- [13] Wald NJ, Simmonds M and Morris JK. Screening for future cardiovascular disease using age alone compared with multiple risk factors and age. *PLoS One* 2011; 6: e18742.
- [14] Goh LG, Dhaliwal SS, Welborn TA, Lee AH and Della PR. Ethnicity and the association between anthropometric indices of obesity and cardiovascular risk in women: a cross-sectional study. *BMJ Open* 2014; 4: e004702.
- [15] Stephanie EC and Nancy R. Lifestyle-Based Prediction Model for the Prevention of CVD: The Healthy Heart Score. *J Am Heart Assoc* 2014; 3: e000954.
- [16] Weihua ML, Yougang W and Jing W. Reduced or modified dietary fat for preventing cardiovascular disease. *J Cardiovasc Nurs* 2013; 28: 204-205.
- [17] Booth FW, Roberts CK and Laye MJ. Lack of exercise is a major cause of chronic diseases. *Compr Physiol* 2012; 2: 1143-1211.
- [18] Chum A and O'Campo P. Cross-sectional associations between residential environmental exposures and cardiovascular diseases. *BMC Public Health* 2015; 15: 438.

SPTY2D1 rs17579600 polymorphism and serum lipid traits

- [19] Thanassoulis G and Vasani RS. Genetic cardiovascular risk prediction: will we get there? *Circulation* 2010; 122: 2323-2334.
- [20] Thanassoulis G, Peloso GM, Pencina MJ, Hoffmann U, Fox CS, Cupples LA, Levy D, D'Agostino RB, Hwang SJ, O'Donnell CJ. A genetic risk score is associated with incident cardiovascular disease and coronary artery calcium: the Framingham Heart Study. *Circ Cardiovasc Genet* 2012; 5: 113-121.
- [21] Nelson TL, Vogler GP, Pedersen NL, Hong Y and Miles TP. Genetic and environmental influences on body fat distribution, fasting insulin levels and CVD: are the influences shared? *Twin Res* 2000; 3: 43-50.
- [22] Gehi AK, Lampert R, Veledar E, Lee F, Goldberg J, Jones L, Murrain N, Ashraf A and Vaccarino V. A twin study of metabolic syndrome and autonomic tone. *J Cardiovasc Electrophysiol* 2009; 20: 422-428.
- [23] Rankinen T, Sarzynski MA, Ghosh S and Bouchard C. Are there genetic paths common to obesity, cardiovascular disease outcomes, and cardiovascular risk factors? *Circ Res* 2015; 116: 909-922.
- [24] Elder SJ, Lichtenstein AH, Pittas AG, Roberts SB, Fuss PJ, Greenberg AS, McCrory MA, Bouchard TJ Jr, Saltzman E and Neale MC. Genetic and environmental influences on factors associated with cardiovascular disease and the metabolic syndrome. *J Lipid Res* 2009; 50: 1917-1926.
- [25] Becker F, van El CG, Ibarreta D, Zika E, Hogarth S, Borry P, Cambon-Thomsen A, Cassiman JJ, Evers-Kiebooms G, Hodgson S, Janssens AC, Kaariainen H, Krawczak M, Kristoffersson U, Lubinski J, Patch C, Penchaszadeh VB, Read A, Rogowski W, Sequeiros J, Tranebjaerg L, van Langen IM, Wallace H, Zimmern R, Schmidtke J and Cornel MC. Genetic testing and common disorders in a public health framework: how to assess relevance and possibilities. Background Document to the ESHG recommendations on genetic testing and common disorders. *Eur J Hum Genet* 2011; 19: S6-44.
- [26] Asselbergs FW, Guo Y, van Iperen EP, Sivapalaratnam S, Tragante V, Lanktree MB, Lange LA, Almqvister B, Appelman YE, Barnard J, Baumert J, Beitelshies AL, Bhangale TR, Chen YD, Gaunt TR, Gong Y, Hopewell JC, Johnson T, Kleber ME, Langaae TY, Li M, Li YR, Liu K, McDonough CW, Meijs MF, Middelberg RP, Musunuru K, Nelson CP, O'Connell JR, Padmanabhan S, Pankow JS, Pankratz N, Rafelt S, Rajagopalan R, Romaine SP, Schork NJ, Shaffer J, Shen H, Smith EN, Tischfield SE, van der Most PJ, van Vliet-Ostaptchouk JV, Verweij N, Volcik KA, Zhang L, Bailey KR, Bailey KM, Bauer F, Boer JM, Braund PS, Burt A, Burton PR, Buxbaum SG, Chen W, Cooper-Dehoff RM, Cupples LA, deJong JS, Delles C, Duggan D, Fornage M, Furlong CE, Glazer N, Gums JG, Hastie C, Holmes MV, Illig T, Kirkland SA, Kivimaki M, Klein R, Klein BE, Kooperberg C, Kottke-Marchant K, Kumari M, LaCroix AZ, Mallesla L, Murugesan G, Ordovas J, Ouwehand WH, Post WS, Saxena R, Scharnagl H, Schreiner PJ, Shah T, Shields DC, Shimbo D, Srinivasan SR, Stolk RP, Swerdlow DI, Taylor HA Jr, Topol EJ, Toshkala E, van Pelt JL, van Setten J, Yusuf S, Whittaker JC, Zwinderman AH; LifeLines Cohort Study, Anand SS, Balmforth AJ, Berenson GS, Bezzina CR, Boehm BO, Boerwinkle E, Casas JP, Caulfield MJ, Clarke R, Connell JM, Cruickshanks KJ, Davidson KW, Day IN, de Bakker PI, Doevendans PA, Dominiczak AF, Hall AS, Hartman CA, Hengstenberg C, Hillge HL, Hofker MH, Humphries SE, Jarvik GP, Johnson JA, Kaess BM, Kathiresan S, Koenig W, Lawlor DA, März W, Melander O, Mitchell BD, Montgomery GW, Munroe PB, Murray SS, Newhouse SJ, Onland-Moret NC, Poulter N, Psaty B, Redline S, Rich SS, Rotter JI, Schunkert H, Sever P, Shuldiner AR, Silverstein RL, Stanton A, Thorand B, Trip MD, Tsai MY, van der Harst P, van der Schoot E, van der Schouw YT, Verschuren WM, Watkins H, Wilde AA, Wolfenbutter BH, Whitfield JB, Hovingh GK, Ballantyne CM, Wijmenga C, Reilly MP, Martin NG, Wilson JG, Rader DJ, Samani NJ, Reiner AP, Hegele RA, Kastelein JJ, Hingorani AD, Talmud PJ, Hakonarson H, Elbers CC, Keating BJ and Drenos F. Large-scale gene-centric meta-analysis across 32 studies identifies multiple lipid loci. *Am J Hum Genet* 2012; 91: 823-838.
- [27] Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, Koseki M, Pirruccello JP, Ripatti S, Chasman DI, Willer CJ, Johansen CT, Fouchier SW, Isaacs A, Peloso GM, Barbalic M, Ricketts SL, Bis JC, Aulchenko YS, Thorleifsson G, Feitosa MF, Chambers J, Orho-Melander M, Melander O, Johnson T, Li X, Guo X, Li M, Shin Cho Y, Jin Go M, Jin Kim Y, Lee JY, Park T, Kim K, Sim X, Tzee-Hee Ong R, Croteau-Chonka DC, Lange LA, Smith JD, Song K, Hua Zhao J, Yuan X, Luan J, Lamina C, Ziegler A, Zhang W, Zee RY, Wright AF, Witteman JC, Wilson JF, Willemssen G, Wichmann HE, Whitfield JB, Waterworth DM, Wareham NJ, Waeber G, Vollenweider P, Voight BF, Vitart V, Uitterlinden AG, Uda M, Tuomilehto J, Thompson JR, Tanaka T, Surakka I, Stringham HM, Spector TD, Soranzo N, Smit JH, Sinisalo J, Silander K, Sijbrands EJ, Scuteri A, Scott J, Schlessinger D, Sanna S, Salomaa V, Saharinen J, Sabatti C, Ruukonen A, Rudan I, Rose LM, Roberts R, Rieder M, Psaty BM, Pramstaller PP, Pichler I, Perola M, Penninx BW, Pedersen NL, Pattaro C, Parker AN, Pare

SPTY2D1 rs17579600 polymorphism and serum lipid traits

- G, Oostra BA, O'Donnell CJ, Nieminen MS, Nickerson DA, Montgomery GW, Meitinger T, McPherson R, McCarthy MI, McArdle W, Masson D, Martin NG, Marroni F, Mangino M, Magnusson PK, Lucas G, Luben R, Loos RJ, Lokki ML, Lettre G, Langenberg C, Launer LJ, Lakatta EG, Laaksonen R, Kyvik KO, Kronenberg F, König IR, Khaw KT, Kaprio J, Kaplan LM, Johansson A, Jarvelin MR, Janssens AC, Ingelsson E, Igl W, Kees Hovingh G, Hottenga JJ, Hofman A, Hicks AA, Hengstenberg C, Heid IM, Hayward C, Havulinna AS, Hastie ND, Harris TB, Haritunians T, Hall AS, Gyllensten U, Guiducci C, Groop LC, Gonzalez E, Gieger C, Freimer NB, Ferrucci L, Erdmann J, Elliott P, Ejebe KG, Döring A, Dominiczak AF, Demissie S, Deloukas P, de Geus EJ, de Faire U, Crawford G, Collins FS, Chen YD, Caulfield MJ, Campbell H, Burt NP, Bonnycastle LL, Boomsma DI, Boehholdt SM, Bergman RN, Barroso I, Bandinelli S, Ballantyne CM, Assimes TL, Quertermous T, Altshuler D, Seielstad M, Wong TY, Tai ES, Feranil AB, Kuzawa CW, Adair LS, Taylor HA Jr, Borecki IB, Gabriel SB, Wilson JG, Holm H, Thorsteinsdottir U, Gudnason V, Krauss RM, Mohlke KL, Ordovas JM, Munroe PB, Kooner JS, Tall AR, Hegele RA, Kastelein JJ, Schadt EE, Rotter JI, Boerwinkle E, Strachan DP, Mooser V, Stefansson K, Reilly MP, Samani NJ, Schunkert H, Cupples LA, Sandhu MS, Ridker PM, Rader DJ, van Duijn CM, Peltonen L, Abecasis GR, Boehnke M and Kathiresan S. Biological, clinical and population relevance of 95 loci for blood lipids. *Nature* 2010; 466: 707-713.
- [28] Jeemon P, Pettigrew K, Sainsbury C, Prabhakaran D and Padmanabhan S. Implications of discoveries from genome-wide association studies in current cardiovascular practice. *World J Cardiol* 2011; 3: 230-247.
- [29] Novoseler M, Hershkovits G and Katcoff DJ. Functional domains of the yeast chromatin protein Sin1p/Spt2p can bind four-way junction and crossing DNA structures. *J Biol Chem* 2005; 280: 5169-5177.
- [30] Bhat W, Boutin G, Rufiange A and Nourani A. Casein kinase 2 associates with the yeast chromatin reassembly factor Spt2/Sin1 to regulate its function in the repression of spurious transcription. *Mol Cell Biol* 2013; 33: 4198-4211.
- [31] Osakabe A, Tachiwana H, Takaku M, Hori T, Obuse C, Kimura H, Fukagawa T and Kurumizaka H. Vertebrate Spt2 is a novel nucleolar histone chaperone that assists in ribosomal DNA transcription. *J Cell Sci* 2013; 126: 1323-1332.
- [32] Guo T, Yin RX, Chen X, Bin Y, Nie RJ and Li H. Sex-specific association of the SPTY2D1 rs7934205 polymorphism and serum lipid levels. *Int J Clin Exp Pathol* 2015; 8: 665-681.
- [33] Xu L, Deng QY, Li SF, Zhou LN, Gong JC and Wei BY. Genetic analysis of Mulao nationality using 15 short tandem repeats. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 2008; 25: 96-100.
- [34] Legato MJ. Gender and the heart: sex-specific differences in normal anatomy and physiology. *J Gend Specif Med* 2000; 3: 15-18.
- [35] Legato MJ. Postpartum depression and gender-specific justice. *J Gend Specif Med* 2002; 5: 7-8.
- [36] Guo T, Yin RX, Lin QZ, Wu J, Shen SW, Sun JQ, Shi GY, Wu JZ, Li H and Wang YM. Polymorphism of rs873308 near the transmembrane protein 57 gene is associated with serum lipid levels. *Biosci Rep* 2014; 34: e00095.
- [37] Guo T, Yin RX, Wu J, Lin QZ, Shi GY, Shen SW, Sun JQ, Li H, Lin WX and Yan DZ. Association of the angiopoietin-like protein 8 rs2278426 polymorphism and several environmental factors with serum lipid levels. *Mol Med Rep* 2015; 12: 3285-3296.
- [38] Guo T, Yin RX, Li H, Wang YM, Wu JZ and Yang DZ. Association of the Trp316Ser variant (rs1801690) near the apolipoprotein H (β -glycoprotein-I) gene and serum lipid levels. *Int J Clin Exp Pathol* 2015; 8: 7291-7304.
- [39] Guo T, Yin RX, Nie RJ, Chen X, Bin Y and Lin WX. Suppressor of cytokine signaling 3 A+930-->G (rs4969168) polymorphism is associated with apolipoprotein A1 and low-density lipoprotein cholesterol. *Int J Clin Exp Pathol* 2015; 8: 7305-7317.
- [40] Bermudez OI, Velez-Carrasco W, Schaefer EJ and Tucker KL. Dietary and plasma lipid, lipoprotein, and apolipoprotein profiles among elderly Hispanics and non-Hispanics and their association with diabetes. *Am J Clin Nutr* 2002; 76: 1214-1221.
- [41] Garcia-Palmieri MR, Tillotson J, Cordero E, Costas R Jr, Sorlie P, Gordon T, Kannel WB and Colon AA. Nutrient intake and serum lipids in urban and rural Puerto Rican men. *Am J Clin Nutr* 1977; 30: 2092-2100.
- [42] Solá R, Fitó M, Estruch R, Salas-Salvadó J, Corella D, de La Torre R, Muñoz MA, López-Sabater Mdel C, Martínez-González MA, Arós F, Ruiz-Gutierrez V, Fiol M, Casals E, Wärnberg J, Buil-Cosiales P, Ros E, Konstantinidou V, Lapeira J, Serra-Majem L and Covas MI. Effect of a traditional Mediterranean diet on apolipoproteins B, A-I, and their ratio: a randomized, controlled trial. *Atherosclerosis* 2011; 218: 174-180.
- [43] Valente EA, Sheehy ME, Avila JJ, Gutierrez JA, Delmonico MJ and Lofgren IE. The effect of the addition of resistance training to a dietary education intervention on apolipoproteins and diet quality in overweight and obese older adults. *Clin Interv Aging* 2011; 6: 235-241.

SPTY2D1 rs17579600 polymorphism and serum lipid traits

- [44] Watts GF, Jackson P, Burke V and Lewis B. Dietary fatty acids and progression of coronary artery disease in men. *Am J Clin Nutr* 1996; 64: 202-209.
- [45] Grundy SM and Denke MA. Dietary influences on serum lipids and lipoproteins. *J Lipid Res* 1990; 31: 1149-1172.
- [46] Predazzi IM, Sobota RS, Sanna S, Bush WS, Bartlett J, Lilley JS, Linton MF, Schlessinger D, Cucca F, Fazio S and Williams SM. Sex-Specific Parental Effects on Offspring Lipid Levels. *J Am Heart Assoc* 2015; 4: e001951.
- [47] Nicot A. Gender and sex hormones in multiple sclerosis pathology and therapy. *Front Biosci (Landmark Ed)* 2009; 14: 4477-4515.
- [48] Wang HH, Liu M, Clegg DJ, Portincasa P and Wang DQ. New insights into the molecular mechanisms underlying effects of estrogen on cholesterol gallstone formation. *Biochim Biophys Acta* 2009; 1791: 1037-1047.
- [49] Tomaszewski M, Charchar FJ, Maric C, Kuzniewicz R, Gola M, Grzeszczak W, Samani NJ and Zukowska-Szczechowska E. Association between lipid profile and circulating concentrations of estrogens in young men. *Atherosclerosis* 2009; 203: 257-262.
- [50] Yu-Poth S, Zhao G, Etherton T, Naglak M, Jonnalagadda S and Kris-Etherton PM. Effects of the National Cholesterol Education Program's Step I and Step II dietary intervention programs on cardiovascular disease risk factors: a meta-analysis. *Am J Clin Nutr* 1999; 69: 632-646.
- [51] Rimm EB, Williams P, Fosher K, Criqui M and Stampfer MJ. Moderate alcohol intake and lower risk of coronary heart disease: meta-analysis of effects on lipids and haemostatic factors. *BMJ* 1999; 319: 1523-1528.
- [52] Yin RX, Wu DF, Miao L, Aung LH, Cao XL, Yan TT, Long XJ, Liu WY, Zhang L and Li M. Several genetic polymorphisms interact with overweight/obesity to influence serum lipid levels. *Cardiovasc Diabetol* 2012; 11: 123.
- [53] Yin RX, Wu DF, Miao L, Htet Aung LH, Cao XL, Yan TT, Long XJ, Liu WY, Zhang L and Li M. Interactions of several single nucleotide polymorphisms and high body mass index on serum lipid traits. *Biofactors* 2013; 39: 315-325.