

## Original Article

# miR-204 inhibits invasion and epithelial-mesenchymal transition by targeting FOXM1 in esophageal cancer

Yurong Sun, Xiuwen Yu, Qingyang Bai

Department of Pathology, Qiqihar Medical University, Qiqihar 161006, Heilongjiang, China

Received August 17, 2015; Accepted September 23, 2015; Epub October 1, 2015; Published October 15, 2015

**Abstract:** MicroRNAs (miRNAs), endogenous noncoding small RNAs, have been reported to play crucial roles in epithelial-mesenchymal transition (EMT) in cancers. Deregulation of microRNA-204 (miR-204) has been documented in many cancers, but its role in the development of esophageal cancer (EC) has not been studied. Here, we reported the role of miR-204 in invasion and EMT in EC. We identified an inverse correlation between miR-204 expression level and the invasion and EMT phenotype of EC cells, and up-regulation of miR-204 inhibited invasion and EMT phenotype of EC cells. Furthermore, we showed that forkhead box protein M1 (FOXM1) was a direct target gene of miR-204, and miR-204 regulated invasion and EMT in EC by acting directly on the 3'UTR of FOXM1 mRNA and suppressing its protein expression. We also explored the anti-tumor effect of miR-204, and found that overexpression of miR-204 suppressed the growth of esophageal tumors *in vivo*. These findings suggest that miR-204 might be a suppressor of invasion and EMT in EC, which offers a novel potential therapeutic target for EC.

**Keywords:** Esophageal cancer, miR-204, forkhead box protein M1 (FOXM1), epithelial-mesenchymal transition, invasion

## Introduction

Esophageal cancer (EC) is one of the most frequent malignant tumors, which is the fifth and eighth most frequent cause of cancer-related death in male and female worldwide, respectively [1]. Despite the great advances achieved in diagnosis and multimodality therapies recently, the prognosis of EC is still poor with a 5-year survival rate of 26.2% [1, 2]. Besides, over 40% of esophageal carcinoma cases result in recurrence [3]. The high probability of metastasis and recurrence is the major cause of treatment failure, yet the precise molecular mechanism of metastatic dissemination is still not completely clear [4, 5]. Recently, many studies have demonstrated that epithelial-mesenchymal transition (EMT) plays a critical role in cancer metastasis [6]. Forkhead box protein M1 (FOXM1) is one of transcription factors, which plays an important role in the activation of EMT [7-9]. The EMT is a process whereby epithelial cells with a cobblestone morphology obtain mesenchymal cell features with a spindle-shaped fibroblast-like phenotype [9]. With the change of cell morphology, the epi-

thelial markers E-cadherin and  $\alpha$ 1-catenin decrease, and the mesenchymal markers N-cadherin, fibronectin and vimentin increase [5, 7, 8]. Besides, this process involves a disassembly of cell-cell junctions, which makes mesenchymal phenotypic cells have less cell adhesion capacity, and higher cell migration and invasion capacity, resulting in tumor aggressiveness [9].

MicroRNAs (miRNAs) are derived from characteristic hairpins in primary transcripts, and are a class of endogenous noncoding small RNAs with approximately 19-23 nucleotides, which regulate gene expression by binding to complementary sequences of targeted mRNAs, causing translational inhibition or degradation [10, 11]. miRNAs are involved in various biological processes, such as cellular proliferation, differentiation, apoptosis, aging, stress response, oncogenesis, and tumor suppression [12-15]. Mounting evidence indicates that aberrant expression of miRNAs occurs in many kinds of malignant tumors, some of which function as tumor oncogenes or suppressor genes [16, 17]. Furthermore, some studies have implied that

miRNAs function as critical modulators for EMT [18-20]. miR-204 has been reported to act as a tumor suppressor, and is obviously down-regulated in various types of solid malignant tumors [21-23]. Recent studies have found that the expression level of miR-204 in EC cells is decreased, but also directly related to the prognosis [24]. However, for miR-204, the related target genes and molecular mechanism in EC are still not well elucidated.

In this study, we detected the expression level of miR-204 in primary EC cases and cell lines by quantitative Real-Time PCR (qRT-PCR), and explored the association between miR-204 expression and invasive potential and EMT phenotype of EC cells. We further investigated the molecular mechanism by which miR-204 exerts regulatory effects on invasive potential and EMT phenotype in EC. Furthermore, we performed the animal experiments to explore the anticancer action of miR-204 *in vivo*. These findings will provide a novel potential therapeutic target for EC.

## Materials and methods

### *Clinical samples*

Esophageal cancer samples and the corresponding adjacent normal esophageal tissues were obtained from 21 patients with EC who underwent radical esophagectomy in the First Affiliated Hospital of Qiqihar Medical University. These 21 patients did not receive any treatment prior to surgery, and cases were confirmed by histopathological examination. All samples were stored in the refrigerator of -70°C after liquid nitrogen freezing. This study was approved by the Ethical Committee of Qiqihar Medical University, and all patients had written informed consent.

### *Cell lines and cell transfection*

Two EC cell lines EC109 and TE10 were obtained from American Type Culture Collection (ATCC, Manassas, VA). Human esophageal epithelial cells (HEEpiC) were purchased from Sciencell (Carlsbad, CA, USA). These cells were grown in RPMI 1640 medium (Hyclone, USA) with 10% fetal bovine serum (FBS, Hyclone, USA), 100 µg/ml of streptomycin, and 100 IU/ml of penicillin at 37°C in 5% CO<sub>2</sub>.

miR-204 mimics, negative control mimics (NC), miR-204 inhibitors (anti-miR-204 mimics), neg-

ative control inhibitors (anti-NC), and FOXM1 siRNAs were purchased from GenePharma Company (Shanghai, China). Cell transfection was carried out with Lipofectamine 2000 (Invitrogen, CA, USA) following the manufacturer's suggestions.

### *Quantitative real-time PCR (qRT-PCR)*


Total RNA samples were isolated from cultured cells or tissues using Trizol (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions, and both mRNA and miRNA were inversely transcribed to cDNA. The expression level of miR-204 was detected by TaqMan stem-loop RT-PCR method with a mirVana miRNA Detection Kit and the gene-specific primer (Invitrogen). Expression of FOXM1 mRNA was examined by SYBR Green real-time PCR (RT-PCR). RT-PCR was carried out by ABI 7500 Fast Real-Time PCR system (ABI, CA, USA). GAPDH was used for normalization.

### *Cell invasion assay*

Cell invasion assay was performed by using transwell chambers.  $2 \times 10^4$  cells were added into the top chamber with matrigel-coated membrane (24-well insert; 8 µm pore size; BD, San Jose, Biosciences, MA, USA). Top chamber contained serum-free medium, and full medium supplemented with 10% FBS was used as a chemoattractant in the bottom chamber. After 24 h incubation at 37°C, the cells which did not invade through the pores were carefully wiped out by a cotton-tipped swab. Cells on the lower surface of the membrane were immobilized with 4% paraformaldehyde, stained with 0.5% crystal violet, then rinsed with PBS and subjected to microscopic inspection. Invaded cells were counted (five high-power fields/chamber) using an inverted microscope (Olympus, Tokyo, Japan).

### *Western blot analysis*

Total proteins were extracted from cultured cells or tissues by using a RIPA buffer with 0.5% sodium dodecyl sulfate (SDS) in the presence of protease inhibitor cocktail (Beytime, Shanghai, China). Then the standard techniques polyacrylamide gel electrophoresis (PAGE), tank-based transfer to Immobilon Hybond-C membranes (Abcam, Cambridge, UK), and immunological detection were performed in turn. Primary antibodies against



**Figure 1.** The expression level of miR-204 in primary EC cases and cell lines. A. qRT-PCR shows that miR-204 was frequently up-regulated in 21 primary ES tissues compared with their adjacent normal tissues. B. Up-regulated miR-204 was detected in EC109 and TE10 cell line compared with human esophageal epithelial cells (HEEpiC). \*\* $P < 0.01$ .

E-cadherin (Genetex, San Antonio, Southern California USA),  $\alpha$ 1-catenin (Genmed Scientifics, Arlington, MA, USA), N-cadherin (Abcam), fibronectin (Abcam), vimentin (Sigma, St. Louis, MO, USA), FOXM1 (Sigma) and GAPDH (Abcam), and secondary antibody peroxidase-conjugated anti-IgG (Genmed Scientifics) were used in western blot analysis. Signals were visualized by a chemiluminescent detection system (Pierce ECL Substrate Western blot detection system, Thermo, Pittsburgh, PA) and then exposed in Molecular Imager ChemiDoc XRS System (Bio-Rad, Hercules, CA, USA). The integrated density of the band was quantified by Image J software (version 1.6 NIH).

#### Target prediction analysis

PicTar, TargetScan, and MicroRNA.org were used to perform bioinformatics-based target prediction analysis.

#### Luciferase reporter assay

The fragments including the 3'UTR regions (3'UTR-WT) or mutant 3'UTR regions (3'UTR-MUT) of FOXM1 were inserted into XhoI/NotI-digested vector psiCHECK-2 (Promega, Madison, WI) with a firefly and renilla luciferase reporter gene. Then, the psiCHECK-2 vectors including 3'UTR regions or mutant 3'UTR regions of FOXM1 were transfected into miR-204-overexpressing EC109 cells and control cells, separately. Twenty-four hours later, cells


were collected, and the firefly and renilla luciferase activities were detected using a dual-luciferase reporter assay system (Promega, USA) according to the manufacturer's suggestions.

#### In vivo tumor formation assay

Animal experiments were carried out complying with the United Kingdom Co-ordinating Committee on Cancer Research (UKCCCR) guidelines for the Welfare of Experimental Animals. For *in vivo* study, EC109 cells ( $1 \times 10^4$ ) transfected with miR-204 mimics or negative control mimics were subcutaneously injected into the dorsal flanks of nude mice. The volume of subcutaneous tumors was measured weekly by manual caliper measurements. Tumor volumes were calculated using the formula: Tumor volume = length  $\times$  width<sup>2</sup>/2 [25]. After 4 weeks, tested mice were sacrificed, and tumor weight was measured. In addition, the expression levels of E-cadherin,  $\alpha$ 1-catenin, N-cadherin, fibronectin, vimentin and FOXM1 protein in xenografts were detected by western blot analysis.

#### Statistical analysis

GraphPad Prism 5 (GraphPad Software, San Diego, CA, USA) was used for statistical analysis. Results were presented as mean  $\pm$  standard deviation (SD). Student's *t*-test was used to analyze the differences between groups. Statistical significance was presumed when  $P < 0.05$ .



**Figure 2.** miR-204 inhibited invasive and EMT of EC cell lines. A and B. EC109 and TE10 cells transfected with miR-204 mimics had a significantly lower invasion capacity than the control group. C and D. Compared with the control group, EC109 and TE10 cells transfected with miR-204 mimic had dramatic decreases in N-cadherin, fibronectin and vimentin expression and significant increases in E-cadherin and  $\alpha$ 1-catenin expression.  $**P < 0.01$ .

## Results

### *Down-regulation of miR-204 is frequently detected in EC*

We compared expression level of miR-204 between tumor and corresponding adjacent normal esophageal tissue in 21 EC patients. The result showed that the expression level of miR-204 in tumor tissues was significantly lower than that in their corresponding normal tissues (**Figure 1A**). Expression level of miR-204 in 2 human EC cell lines was also detected by qRT-PCR and the result showed that miR-204 was obviously down-regulated in EC109 and TE10 cell lines compared with that in HEEpiC (**Figure 1B**).

### *Exogenous expression of miR-204 inhibits invasion of EC cells*

To investigate the effect of miR-204 on the invasive capability in EC cells, we transfected


EC109 and TE10 cells with miR-204 mimics or negative control mimics, respectively, and then performed invasion assays by transwell methods. Results showed that EC109 and TE10 cells transfected with miR-204 mimics had a significantly lower invasion capacity than control, which indicated that miR-204 efficiently inhibits invasion of EC cells (**Figure 2A and 2B**).

### *miR-204 inversely regulates EMT phenotype of EC cells*

Epithelial-mesenchymal transition (EMT) has been identified to contribute to cancer metastasis by transformation of adherent epithelial cells into motile and invasive mesenchymal cells [6]. To investigate whether miR-204 could regulate EMT in EC cells, the expression of epithelial markers (*i.e.* E-cadherin and  $\alpha$ 1-catenin) and the mesenchymal markers (*i.e.* N-cadherin, fibronectin and vimentin) were determined in EC109 and TE10 cells with miR-204 mimics or negative control mimics. The results showed



## miR-204 inhibits invasion and EMT by FOXM1



**Figure 4.** miR-204 suppressed tumorigenesis of EC *in vivo*. A. In the miR-204 group, the tumor volume was significantly decreased than that in the control group. B. After 4 weeks, the tumor weight was significantly decreased in the miR-204 group than that in the control group. C. miR-204 inhibited the expression of FOXM1. D. miR-204 decreased N-cadherin, fibronectin and vimentin expression and increased E-cadherin, α1-catenin expression compared to the control group. \* $P < 0.05$ , \*\* $P < 0.01$ .

was a binding target of miR-204. The results showed that in the miR-204 group, the luciferase activity driven by FOXM1 3'UTR was significantly lowered compared with that in negative control cells, while miR-204-mediated inhibition of luciferase activity was abolished by the mutant presumptive binding site, which suggested that FOXM1 is indeed a direct target of miR-204 (Figure 3B). Furthermore, an inverse correlation was directly observed between miR-204 and FOXM1 mRNA ( $r = -0.861$ ;  $P < 0.0001$ ; Figure 3C) in EC samples. To confirm the regulatory effect of miR-204 on FOXM1, western blot assay was performed to detect the expression level of FOXM1 in response to the change of miR-204 expression in EC109 cells. As shown in Figure 3D, the results showed a negative regulatory effect of miR-204 on FOXM1. Up-regulated miR-204 in EC109 cells significantly decreased FOXM1 protein expression compared with cells transfected with negative control mimics. Relatively, down-regulation of miR-204 by inhibitors increased FOXM1 protein expression level. To fully explore the roles of

FOXM1 in EC, EC109 cells were transfected with FOXM1 siRNAs (Si-FOXM1) to silence the expression of FOXM1. The transwell invasion assay as illustrated in Figure 3E showed that in the Si-FOXM1 group, the number of migrated cells decreased significantly compared to the Si-control group, which indicated that knock-down of FOXM1 represses invasion of EC109 cells. Western blot analysis showed that Si-FOXM1 cells had dramatic decreases in N-cadherin, fibronectin and vimentin expression and significant increases in E-cadherin and α1-catenin expression compared to the Si-control group, which indicated that knock-down of FOXM1 inhibits EMT phenotype of EC109 cells (Figure 3F). Similar effect of Si-FOXM1 was observed in miR-204 mimics treated cells, suggesting that Si-FOXM1 could mimic the effect of miR-204. Taken together, these results suggested that FOXM1 is one of the direct targets of miR-204, and functional effect of miR-204 on EC cell lines is dependent on FOXM1.

## miR-204 inhibits invasion and EMT by FOXM1

### *Overexpression of miR-204 suppresses growth of esophageal tumors in vivo*

To explore the anti-tumor effect of miR-204, we subcutaneously injected EC109 cells ( $1 \times 10^4$ ) transfected with miR-204 mimics or negative control mimics into the dorsal flanks of nude mice. We measured the volume of subcutaneous tumors weekly and found that in the miR-204 group, the tumor growth was significantly inhibited than that in the control group (**Figure 4A**). After 4 weeks, tested mice were sacrificed, and tumor weight was measured. Accordingly, in the miR-204 group, the tumor weight was significantly lower than that in the control group (**Figure 4B**). In addition, we detected the expression of FOXM1, epithelial markers (*i.e.* E-cadherin and  $\alpha$ 1-catenin), and the mesenchymal markers (*i.e.* N-cadherin, fibronectin and vimentin) in xenografts by western blot. As shown in **Figure 4C**, compared with the control group, the miR-204 group had an obviously low expression of FOXM1. Furthermore, the miR-204 group had dramatic decreases in N-cadherin, fibronectin and vimentin expression and significant increases in E-cadherin and  $\alpha$ 1-catenin expression compared with the control group (**Figure 4D**). Taken together, these results indicated that overexpression of miR-204 suppresses growth of esophageal tumors *in vivo*.

### Discussion

EC is one of the common causes of cancer-related death worldwide. The main reason of cancer death is complications arising from metastasis, while EMT is the key process driving cancer metastasis [6, 26]. Losses of E-cadherin and  $\alpha$ 1-catenin expression and increases of N-cadherin, fibronectin and vimentin expression are identified as the most important molecular markers of EMT [5, 7, 27, 28]. Emerging evidences have showed that miRNAs play critical roles in the regulation of EMT of cancer cells [29]. A variety of miRNAs have been identified as the promoters for EMT. miR-23a inhibits E-cadherin expression and promotes EMT under the stimulation of TGF- $\beta$ 1 [30]. miR-31 represses the tumor suppressor expression and stimulates EMT [31]. miR-9 promotes tumor metastasis via inhibiting E-cadherin expression in esophageal squamous cell carcinoma [5]. Conversely, many miR-

NAs have been identified as the suppressors for EMT. Particularly, miR-149 and miR-134 inhibit non-small-cell lung cancer cells EMT by targeting FOXM1, and miR-140 and miR-625 repress EC cells EMT and invasion by targeting Slug and Sox2, respectively [7, 8, 32, 33].

Recent researches have demonstrated the essential role of miR-204 in the progression of various types of malignant tumors. miR-204 is down-regulated in some tumors, and functions as a tumor suppressor. For example, loss of miR-204 expression promotes glioma migration and stem cell-like phenotype, and miR-204 represses tumor growth through inhibition of MAP1LC3B-mediated autophagy in renal clear cell carcinoma [21, 23]. Recent studies have also found that miR-204 expression is decreased in EC cells, but also directly related to the prognosis [24]. However, the role of miR-204 in EC is still not well elucidated. Our study demonstrated that up-regulation of miR-204 inhibits invasion and EMT in EC.

FOXM1 is one of transcription factors characterized by the presence of a DNA-binding domain named the forkhead box or winged helix domain, which is detected in many tumor cell lines and regulates the expression of genes related to cell cycle [34]. Overexpression of FOXM1 has been found in various malignant tumors, such as esophagus, breast, lung, colon, and pancreas cancer [7, 35-37]. Interestingly, some studies on FOXM1 have implicated its participation in invasion and EMT of the tumor cells, the key processes contributing to cancer metastasis. Ke *et al.* and Li *et al.* found that FOXM1 stimulates invasive and EMT phenotype of non-small cell lung cancer cells [7, 8]. Bao *et al.* demonstrated that overexpression of FOXM1 leads to EMT and cancer stem cell phenotype in pancreatic cancer cells [9]. In this study, we obtained that FOXM1 is a potential target gene of hsa-miR-204 by bioinformatics analysis. Luciferase assay confirmed that miR-204 targets FOXM1 directly. An inverse correlation is also found between miR-204 and FOXM1 mRNA in clinical patients with EC. Furthermore, knockdown of FOXM1 by siRNA reverses invasion and EMT in EC cells. Li *et al.* observed a similar result that knockdown of FOXM1 by siRNA reversed EMT in non-small cell lung cancer cells [8]. These findings implied that miR-204 suppresses FOXM1 protein expression by

directly binding on the 3'UTR of FOXM1 mRNA to negatively regulate invasion and EMT in EC. We also explored the anti-tumor effect of miR-204 *in vivo*. We reached the conclusion that overexpression of miR-204 suppresses growth of esophageal tumors.

In summary, down-regulation of miR-204 is frequently detected in EC, and miR-204 inversely regulates invasion and EMT phenotype of EC cell. We also provided evidence to prove that miR-204 regulates invasion and EMT in EC by acting directly on the 3'UTR of FOXM1 mRNA. Due to the limit on the number of EC samples and cell types, further comprehensive study will be necessary for exploring the potential role of miR-204 in EC development. miR-204-FOXM1 pathway that we studied might be exploited by a therapeutic strategy for EC treatment in future.

#### Acknowledgements

Thank to all the members in our department for their help in experiment methods suggestion and data collection. This study was supported by the Science and Technology Research Project of Department of Education, Heilongjiang Province, China (12531804).

#### Disclosure of conflict of interest

None.

**Address correspondence to:** Dr. Qingyang Bai, Department of Pathology, Qiqihar Medical University, 333 Bukui North Street, Jianhua District, Qiqihar 161006, Heilongjiang, P. R. China. Tel: +86-452-2663483; E-mail: bqingyangedu@163.com

#### References

- [1] Yuequan J, Shifeng C, Bing Z. Prognostic factors and family history for survival of esophageal squamous cell carcinoma patients after surgery. *Ann Thorac Surg* 2010; 90: 908-913.
- [2] Yue D, Zhang Z, Li J, Chen X, Ping Y, Liu S, Shi X, Li L, Wang L, Huang L, Zhang B, Sun Y, Zhang Y. Transforming growth factor-beta1 promotes the migration and invasion of sphere-forming stem-like cell subpopulations in esophageal cancer. *Exp Cell Res* 2015; 336: 141-149.
- [3] Aminian A, Panahi N, Mirsharifi R, Karimian F, Meysamie A, Khorgami Z, Alibakhshi A. Predictors and outcome of cervical anastomotic leakage after esophageal cancer surgery. *J Cancer Res Ther* 2011; 7: 448-453.
- [4] Enzinger PC, Mayer RJ. Esophageal cancer. *N Engl J Med* 2003; 349: 2241-2252.
- [5] Song Y, Li J, Zhu Y, Dai Y, Zeng T, Liu L, Li J, Wang H, Qin Y, Zeng M, Guan XY, Li Y. MicroRNA-9 promotes tumor metastasis via repressing E-cadherin in esophageal squamous cell carcinoma. *Oncotarget* 2014; 5: 11669-11680.
- [6] Pavelic SK, Sedic M, Bosnjak H, Spaventi S, Pavelic K. Metastasis: new perspectives on an old problem. *Mol Cancer* 2011; 10: 22.
- [7] Ke Y, Zhao W, Xiong J, Cao R. miR-149 inhibits non-small-cell lung cancer cells EMT by targeting FOXM1. *Biochem Res Int* 2013; 2013: 506731.
- [8] Li J, Wang Y, Luo J, Fu Z, Ying J, Yu Y, Yu W. miR-134 inhibits epithelial to mesenchymal transition by targeting FOXM1 in non-small cell lung cancer cells. *FEBS Lett* 2012; 586: 3761-3765.
- [9] Bao B, Wang Z, Ali S, Kong D, Banerjee S, Ahmad A, Li Y, Azmi AS, Miele L, Sarkar FH. Over-expression of FoxM1 leads to epithelial-mesenchymal transition and cancer stem cell phenotype in pancreatic cancer cells. *J Cell Biochem* 2011; 112: 2296-2306.
- [10] Garzon R, Pichiorri F, Palumbo T, Visentini M, Aqeilan R, Cimmino A, Wang H, Sun H, Volinia S, Alder H, Calin GA, Liu CG, Andreeff M, Croce CM. MicroRNA gene expression during retinoic acid-induced differentiation of human acute promyelocytic leukemia. *Oncogene* 2007; 26: 4148-4157.
- [11] Mack GS. MicroRNA gets down to business. *Nat Biotechnol* 2007; 25: 631-638.
- [12] Li Q, Gregory RI. MicroRNA regulation of stem cell fate. *Cell Stem Cell* 2008; 2: 195-196.
- [13] Callis TE, Chen JF, Wang DZ. MicroRNAs in skeletal and cardiac muscle development. *DNA Cell Biol* 2007; 26: 219-225.
- [14] Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, Visone R, Iorio M, Roldo C, Ferracin M, Prueitt RL, Yanaihara N, Lanza G, Scarpa A, Vecchione A, Negrini M, Harris CC, Croce CM. A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci U S A* 2006; 103: 2257-2261.
- [15] Dresios J, Aschrafi A, Owens GC, Vanderklish PW, Edelman GM, Mauro VP. Cold stress-induced protein Rbm3 binds 60S ribosomal subunits, alters microRNA levels, and enhances global protein synthesis. *Proc Natl Acad Sci U S A* 2005; 102: 1865-1870.
- [16] Calin GA, Croce CM. MicroRNA signatures in human cancers. *Nat Rev Cancer* 2006; 6: 857-866.
- [17] Kumar MS, Lu J, Mercer KL, Golub TR, Jacks T. Impaired microRNA processing enhances cellular transformation and tumorigenesis. *Nat Genet* 2007; 39: 673-677.



## miR-204 inhibits invasion and EMT by FOXM1

- [18] Gregory PA, Bert AG, Paterson EL, Barry SC, Tsykin A, Farshid G, Vadas MA, Khew-Goodall Y, Goodall GJ. The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by targeting ZEB1 and SIP1. *Nat Cell Biol* 2008; 10: 593-601.
- [19] Dong P, Kaneuchi M, Watari H, Hamada J, Sudo S, Ju J, Sakuragi N. MicroRNA-194 inhibits epithelial to mesenchymal transition of endometrial cancer cells by targeting oncogene BMI-1. *Mol Cancer* 2011; 10: 99.
- [20] Kumarswamy R, Mudduluru G, Ceppi P, Muppala S, Kozlowski M, Niklinski J, Papotti M, Allgayer H. MicroRNA-30a inhibits epithelial-to-mesenchymal transition by targeting Snai1 and is downregulated in non-small cell lung cancer. *Int J Cancer* 2012; 130: 2044-2053.
- [21] Mikhaylova O, Stratton Y, Hall D, Kellner E, Ehmer B, Drew AF, Gallo CA, Plas DR, Biesiada J, Meller J, Czyzyk-Krzeska MF. VHL-regulated MiR-204 suppresses tumor growth through inhibition of LC3B-mediated autophagy in renal clear cell carcinoma. *Cancer cell* 2012; 21: 532-546.
- [22] Sacconi A, Biagioni F, Canu V, Mori F, Di Benedetto A, Lorenzon L, Ercolani C, Di Agostino S, Cambria A, Germoni S, Grasso G, Blandino R, Panebianco V, Ziparo V, Federici O, Muti P, Strano S, Carboni F, Mottolese M, Diodoro M, Pescarmona E, Garofalo A, Blandino G. miR-204 targets Bcl-2 expression and enhances responsiveness of gastric cancer. *Cell Death Dis* 2012; 3: e423.
- [23] Ying Z, Li Y, Wu J, Zhu X, Yang Y, Tian H, Li W, Hu B, Cheng SY, Li M. Loss of miR-204 expression enhances glioma migration and stem cell-like phenotype. *Cancer Res* 2013; 73: 990-999.
- [24] Alder H, Taccioli C, Chen H, Jiang Y, Smalley KJ, Fadda P, Ozer HG, Huebner K, Farber JL, Croce CM, Fong LY. Dysregulation of miR-31 and miR-21 induced by zinc deficiency promotes esophageal cancer. *Carcinogenesis* 2012; 33: 1736-1744.
- [25] Zhu X, Li Y, Shen H, Li H, Long L, Hui L, Xu W. miR-137 inhibits the proliferation of lung cancer cells by targeting Cdc42 and Cdk6. *FEBS Lett* 2013; 587: 73-81.
- [26] Gavert N, Ben-Ze'ev A. Epithelial-mesenchymal transition and the invasive potential of tumors. *Trends Mol Med* 2008; 14: 199-209.
- [27] Lin C, Tang X, Zhu Z, Liao X, Zhao R, Fu W, Chen B, Jiang J, Qian R, Guo D. The rhythmic expression of clock genes attenuated in human plaque-derived vascular smooth muscle cells. *Lipids Health Dis* 2014; 13: 14.
- [28] Lee JM, Dedhar S, Kalluri R, Thompson EW. The epithelial-mesenchymal transition: new insights in signaling, development, and disease. *J Cell Biol* 2006; 172: 973-981.
- [29] Lin PY, Yu SL, Yang PC. MicroRNA in lung cancer. *Br J Cancer* 2010; 103: 1144-1148.
- [30] Cao M, Seike M, Soeno C, Mizutani H, Kitamura K, Minegishi Y, Noro R, Yoshimura A, Cai L, Gemma A. MiR-23a regulates TGF- $\beta$ -induced epithelial-mesenchymal transition by targeting E-cadherin in lung cancer cells. *Int J Oncol* 2012; 41: 869-875.
- [31] Liu X, Sempere LF, Ouyang H, Memoli VA, Andrew AS, Luo Y, Demidenko E, Korc M, Shi W, Preis M, Dragnev KH, Li H, Drenzo J, Bak M, Freemantle SJ, Kauppinen S, Dmitrovsky E. MicroRNA-31 functions as an oncogenic microRNA in mouse and human lung cancer cells by repressing specific tumor suppressors. *J Clin Invest* 2010; 120: 1298-1309.
- [32] Li W, Jiang G, Zhou J, Wang H, Gong Z, Zhang Z, Min K, Zhu H, Tan Y. Down-regulation of miR-140 induces EMT and promotes invasion by targeting Slug in esophageal cancer. *Cell Physiol Biochem* 2014; 34: 1466-1476.
- [33] Wang Z, Qiao Q, Chen M, Li X, Wang Z, Liu C, Xie Z. miR-625 down-regulation promotes proliferation and invasion in esophageal cancer by targeting Sox2. *FEBS Lett* 2014; 588: 915-921.
- [34] Wang Y, Wen L, Zhao SH, Ai ZH, Guo JZ, Liu WC. FoxM1 expression is significantly associated with cisplatin-based chemotherapy resistance and poor prognosis in advanced non-small cell lung cancer patients. *Lung Cancer* 2013; 79: 173-179.
- [35] Hui MK, Chan KW, Luk JM, Lee NP, Chung Y, Cheung LC, Srivastava G, Tsao SW, Tang JC, Law S. Cytoplasmic Forkhead box M1 (FoxM1) in esophageal squamous cell carcinoma significantly correlates with pathological disease stage. *World J Surg* 2012; 36: 90-97.
- [36] Millour J, Constantinidou D, Stavropoulou AV, Wilson MS, Myatt SS, Kwok JM, Sivanandan K, Coombes RC, Medema RH, Hartman J, Lykkesfeldt AE, Lam EW. FOXM1 is a transcriptional target of ER $\alpha$  and has a critical role in breast cancer endocrine sensitivity and resistance. *Oncogene* 2010; 29: 2983-2995.
- [37] Koo CY, Muir KW, Lam EW. FOXM1: From cancer initiation to progression and treatment. *Biochim Biophys Acta* 2012; 1819: 28-37.