

Erratum

Versican isoforms modulate expression and function of nicotinic acetylcholine receptors: Int J Physiol Pathophysiol Pharmacol. 2009; 1(1): 64-75

Yao-Jiong Wu^{1,2}, Wang Sheng^{1,2}, Hai-Heng Dong^{1,3}, David Lapierre^{1,2}, Yu-Di Wan^{1,3}, Wei-Yang Lu^{1,3}, Burton B Yang^{1,2}

¹Sunnybrook Research Institute, Sunnybrook Health Sciences Centre, Toronto, Canada; ²Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Canada; ³Department of Anaesthesia, University of Toronto, Toronto, Canada

Received July 14, 2025; Accepted August 25, 2026; Epub August 25, 2026; Published August 30, 2026

In this article, an image in **Figure 2A** was uploaded with an error. We have now used another group of results generated at the same time to replace the original image. We sincerely apologize for this oversight and any confusion it may have caused. The corrected images **Figure 2** is shown below.

Address correspondence to: Drs. Wei-Yang Lu and Burton B Yang, Research Building, Sunnybrook Health Sciences Centre, 2075 Bayview Avenue, Toronto M4N 3M5, Canada. Tel: 416-480-4823; E-mail: weiyang.lu@sri.utoronto.ca (WYL); Tel: 416-480-5874; E-mail: byang@sri.utoronto.ca (BBY)




Figure 2. Versican alters the expression of nicotinic receptors. A. PC12 cells stably transfected with vector or V1 and V2 constructs were incubated with NGF (50 ng/ml). Four days after treatment photographs of differentiation were acquired. B. Western blot analyses of AChR subunits $\alpha 3$, $\alpha 4$, $\alpha 7$ and $\alpha 2$ in the presence or absence of NGF. Expression of $\alpha 3$ and $\alpha 7$ was up-regulated by V1- and V2-transfection, while expression of $\alpha 4$ was down-regulated. In the presence of NGF, expression of $\alpha 3$ and $\alpha 4$ exhibited complementary patterns. C. The intensities of the protein bands were scanned, and the densitometry results were semiquantified ($n = 3$, $*P < 0.05$).